

Occasional Papers of the Museum of Natural Science, Louisiana State University

Volume 1 | Number 68

Article 1

7-15-1994

Type specimens of birds in the Museum of Natural Science, Louisiana State University

Steven W. Cardiff

J. V. Remsen, Jr.

Follow this and additional works at: <https://digitalcommons.lsu.edu/opmns>



Part of the [Genetics and Genomics Commons](#), and the [Population Biology Commons](#)

Recommended Citation

Cardiff, Steven W. and Remsen, Jr., J. V. (1994) "Type specimens of birds in the Museum of Natural Science, Louisiana State University," *Occasional Papers of the Museum of Natural Science, Louisiana State University* : No. 68 , Article 1.

DOI: 10.31390/opmns.068

Available at: <https://digitalcommons.lsu.edu/opmns/vol1/iss68/1>



OCCASIONAL PAPERS OF THE
MUSEUM OF NATURAL SCIENCE

LOUISIANA STATE UNIVERSITY
BATON ROUGE, LOUISIANA 70803

TYPE SPECIMENS OF BIRDS IN THE MUSEUM OF
NATURAL SCIENCE, LOUISIANA STATE UNIVERSITY

By Steven W. Cardiff and J. V. Remsen, Jr.

The Museum of Natural Science, Louisiana State University (LSUMNS), holds 55 holotypes of birds. A large proportion of these are species and distinctive subspecies discovered in the last 30 years in Peru, including four new genera (*Conioptilon*, *Nephelornis*, *Wetmorethraupis*, and *Xenoglaux*). A synopsis of these type specimens has not been published. Herein, we summarize details on LSUMNS holotypes and their paratypes and, where appropriate, provide information on recent range extensions of the Peruvian taxa. We do not list LSUMNS paratypes of species for which LSUMNS does not possess the holotype. Although we have not attempted to analyze independently the validity of all taxa, we note which taxa have been recognized or synonymized by recent authors, and we provide taxonomic comments on many taxa. Catalog numbers of specimens at LSUMNS (formerly Louisiana State University Museum of Zoology) are still preceded by the acronym "LSUMZ."

FALCONIFORMES
FAMILY ACCIPITRIDAE

Buteogallus subtilis rhizophorae Monroe, 1963, Occas. Pap. Mus. Zool. Louisiana State Univ. (26): 1.

HOLOTYPE: LSUMZ 28923, "4 mi. SW San Lorenzo, Department of Valle, Honduras." Adult male collected by J. Alan Feduccia on 5 October 1962. Original number JAF 706.

PARATYPE: LSUMZ 28922, type locality and date, female collected by B. L. Monroe, Jr.

COMMENTS: The complete locality on the holotype label reads "HONDURAS: Valle: 4 mi. SW San Lorenzo, s.l. [=sea level]." Blake (1977) and Stresemann and Amadon (1979) recognized *rhizophorae*.

Buteogallus anthracinus micronyx van Rossem and Hachisuka, 1937, Trans. San Diego Soc. Nat. Hist. 8: 361.

HOLOTYPE: LSUMZ 39222, "Arivaipa Creek, Graham County, Arizona." Adult male collected by W. J. Sheffler on 3 June 1936. Original number WJS 1477.

PARATYPE: LSUMZ 61058, female, other data same as holotype.

COMMENTS: "Arivaipa" in the type description, and "Arivipa" on the holotype label are spelled incorrectly; the correct spelling is "Aravaipa." Hellmayr and Conover (1949) did not believe that *micronyx* was recognizable, but this was based on their examination of only two specimens. Friedmann (1950), Friedmann et al. (1950), Brown and Amadon (1968), Blake (1977), and Stresemann and Amadon (1979) evidently followed Hellmayr and Conover in considering *micronyx* to be a synonym of the nominate subspecies. We are not aware of any thorough examination of the validity of this taxon.

Urubitornis solitaria sheffleri van Rossem, 1948, Proc. Biol. Soc. Washington 61: 67.

HOLOTYPE: LSUMZ 39225, "extreme southeastern Sonora near the Chihuahua boundary." Adult female collected by W. J. Sheffler on 7 June 1949. Original number WJS 2666.

PARATYPE: LSUMZ 39224, Algodones, southeastern Sonora, Mexico, 2500 ft., male collected by WJS on 1 March 1948.

COMMENTS: The complete locality on the holotype label reads "near Chihuahua boundary, Mirasol, 16 mi. NE of Guiracoba, Sonora [Mexico]." Currently placed in *Harpyhaliaetus*. Friedmann (1950), Friedmann et al. (1950), and Brown and Amadon (1968) recognized *sheffleri*; the latter authors suspected that *sheffleri* applied to all Mexican birds north of the Isthmus of Tehuantepec. Blake (1977) suspected that *sheffleri*, then applied only to the isolated population of northern Mexico, was not a valid taxon. Stresemann and Amadon (1979) considered all Middle American populations referable to *sheffleri*.

GALLIFORMES FAMILY CRACIDAE

Crax estudilloi Allen, 1977, Game Bird Breeders Avicult. Zoo. Conserv. Gaz. 26: 6.

HOLOTYPE: LSUMZ 140000 [complete skeleton + some feathers], "Bolivia on the eastern slopes of the Andes near the village of Ixima [sic] in an indian hut at the shore of Tequije [sic] River that flows into the Rio Benii

[sic]." Adult male captured by local indians as a 5-6 week old chick in August or September of 1974 or 1975 and raised in captivity by Jesus Estudillo Lopez. Specimen prepared by Steven W. Cardiff. Original number SWC 5156.

COMMENTS: The capture site is in the depto. La Paz, Prov. Iturralde. Correct spellings: "Ixima"="Ixiamas;" "Tequije"="Tequeje;" "Benii"="Beni." A controversial "species" (see Vuilleumier and Mayr 1987, Remsen and Traylor 1989), based on a photograph and informal description, in captivity, of the only known individual. The bird died about 1989. Through the efforts of Stuart D. Strahl, the carcass was donated to LSUMNS, but it was so badly decomposed when received that it could only be prepared as a skeleton (with some feathers saved as well).

CHARADRIIFORMES
FAMILY LARIDAE

Sterna albifrons athalassos Burleigh and Lowery, 1942, Occas. Pap. Mus. Zool. Louisiana State Univ. (10): 173.

HOLOTYPE: LSUMZ 5443, "St. Francisville, West Feliciana Parish, Louisiana." Adult male collected by Thomas D. Burleigh on 27 May 1941. Original number TDB 10599.

PARATYPES (11): LSUMZ 4193, 4207, Rosedale, Bolivar Co., Mississippi, females collected by G. H. Lowery on 17 July 1940; LSUMZ 5757, 6 mi. S Rosedale, male collected by Sam M. Ray on 6 June 1941; LSUMZ 5453, 5455, Vicksburg (Eagle Lake), Warren Co., Mississippi, males collected by TDB on 20 June 1941; LSUMZ 3396, Mississippi-Tennessee state line at Mud Lake below Memphis, Shelby Co., Tennessee, male collected by E. Wallace on 27 August 1939; LSUMZ 4194, 4195, Baton Rouge, East Baton Rouge Par., Louisiana, female and male collected by GHL on 6 August 1940; LSUMZ 6374, Southwest Pass, Plaquemines Par., Louisiana, female collected by TDB on 13 August 1941; LSUMZ 5382, New Orleans, Orleans Par., Louisiana, female collected by TDB on 12 May 1941; LSUMZ 5890, Pass a Loutre, Plaquemines Par., Louisiana, female collected by TDB on 12 August 1941.

COMMENTS: Several other paratypes have been exchanged. New World populations of *S. albifrons* are now considered to represent a separate species, *S. antillarum*.

Recently, Thompson et al. (1992), using plumage, morphometric, and genetic characters, disputed the validity of *athalassos*. Flaws in their methodology, including failure to control for age and seasonal wear in examination of plumage and no attempt to remove migrant *athalassos* from their sample of coastal populations, prevent their analyses from being considered a proper assessment of the validity of *athalassos*. A detailed critique of their analysis, and a reanalysis of plumage criteria that show that *athalassos* is a valid, recognizable taxon, will be published

elsewhere (Remsen and Cardiff, in prep.). When specimens of comparable age and season are compared, *athalassos* may be distinguished by its darker gray back and, especially, upper tail coverts and rectrices.

PSITTACIFORMES
FAMILY PSITTACIDAE

Amazona ochrocephala magna Monroe and Howell, 1966, Occas. Pap. Mus. Zool. Louisiana State Univ. (34): 5.

HOLOTYPE: LSUMZ 5156, "Victoria, Tamaulipas, México." Male collected by T. D. Burleigh on 27 April 1941. Original number TDB 10489.

PARATYPES (6): LSUMZ 5155, type locality and date, female collected by G. H. Lowery; LSUMZ 16782-83, 10.5 mi. W Ebano, San Luis Potosí, male and female collected by R. Graber and J. Graber, respectively, on 11 April 1951; LSUMZ 10969, Hacienda Limón, San Luis Potosí, male collected by Herbert Stern, Jr. on 25 February 1948; LSUMZ 10968, 2 mi. N Tamuín, San Luis Potosí, female collected by Charles Shaw on 18 September 1946; LSUMZ 23890, 9 mi. N Balancán, Tabasco, male collected by E. T. Armstrong on 14 May 1961.

COMMENTS: The complete locality on the holotype label reads "Mexico: Victoria, near Güemes on Río Corona." Forshaw (1989) did not mention *magna* and apparently considered it to be a synonym of *A. ochrocephala oratrix*. The AOU (1983) treated populations from the Gulf-Caribbean slope of Mexico as a separate species, *A. oratrix*.

Amazona ochrocephala belizensis Monroe and Howell, 1966, Occas. Pap. Mus. Zool. Louisiana State Univ. (34): 6.

HOLOTYPE: LSUMZ 21811, "Hill Bank, Orange Walk Dist., British Honduras." Adult male collected by Stephen M. Russell on 20 November 1956. Original number SMR 1129.

PARATYPES (2): LSUMZ 20985, 22545, type locality, female and male collected by SMR on 21 February 1956, and D. A. Lancaster on 5 May 1958, respectively.

COMMENTS: Forshaw (1989) recognized *belizensis* as applicable to the population in British Honduras [=Belize]. The AOU (1983) treated populations from Belize and the Gulf-Caribbean slope of Mexico, including those within the range of *magna* above, as a separate species, *A. oratrix*.

STRIGIFORMES
FAMILY STRIGIDAE

Otus seductus colimensis Hekstra, 1982, Bull. Zool. Mus. Univ. Amsterdam 9: 61.

HOLOTYPE: LSUMZ 39853, "7 miles south of Colima, Colima, [Mexico]." Adult male collected by W. J. Schaldach, Jr. on 24 December 1958.

Original number W. J. Schaldach 4808 (W. J. Sheffler Collection number 7001).

PARATYPES (20): LSUMZ 39854, type locality, male collected by Schaldach on 24 December 1958; LSUMZ 39855, 44201, type locality, males collected by Schaldach on 27 December 1958; LSUMZ 39851-52, 39856-62, 44195-97, 44200, 44202-05, within an 18 km radius of Colima, 12 males and five females collected by Schaldach and J. T. Marshall during 1958-59.

COMMENTS: Browning (1989) provided numerous comments on the lack of validity of many of Hekstra's new taxa of owls but did not comment on the validity of *colimensis*. Thus, given Hekstra's questionable record of describing new owl taxa, we concur with Browning's statement that "the remaining forms he named require further study."

Glaucidium minutissimum sanchezi Lowery and Newman, 1949, Occas. Pap. Mus. Zool. Louisiana State Univ. (22): 1.

HOLOTYPE: LSUMZ 11002, "Llano de Garzas, near Cerro Coneja [sic], San Luis Potosí, México; elevation 6,800 feet." Male collected by Robert J. Newman on 14 May 1947, prepared by Marcella Newman. Original number RJN 669.

PARATYPES (3): LSUMZ 11003, near Cerro La Mina, Cerro Coneja [=Cerro Conejo] Region, 6500 ft., male, 12 June 1948; LSUMZ 11004, near Cerro La Luz, Cerro Coneja [=Cerro Conejo] Region, 7300 ft., male, 14 May 1947; LSUMZ 11005, beyond Puerto de Loso, Xilitla Region, 5600 ft., female, 5 February 1947 (all collected by RJN).

COMMENTS: A fourth paratype (LSUMZ 11001) has been exchanged. Additional skins were obtained by L. Guerrero at "Cerro San Antonio (=Cerro Miramar)" on 14 July 1950 (LSUMZ 15036, male) and by J. Graber at "Cerro Conejo" on 15 June 1951 (LSUMZ 16796, female).

Xenoglaux loweryi O'Neill and Graves, 1977, Auk 94: 411.

HOLOTYPE: LSUMZ 84000 (skin and partial skeleton), "10 km by road northeast of Abra Patricia on road to Rioja, 05°46'S, 77°41'W, Depto. San Martín, Perú, elevation approximately 1890 m." Adult male collected by J. P. O'Neill on 23 August 1976. Original number JPO 5638.

PARATYPES (2): LSUMZ 84001 (skin and partial skeleton), female, other data same as holotype; LSUMZ 84002 (skin and partial skeleton), type locality, female collected by JPO on 22 August 1976.

COMMENTS: A new genus (Auk 94: 410) and species. Two additional skins (LSUMZ 87292-93, female, male) were obtained by T. S. Schulenberg, Morris D. Williams, and others on 15 October 1978 in the Cordillera Colán, southeast of La Peca, ca. 2350 m, depto. Amazonas, Peru, approximately 90 km west-northwest of the type locality.

APODIFORMES
FAMILY TROCHILIDAE

Metallura theresiae parkeri Graves, 1981, Auk 98: 382.

HOLOTYPE: LSUMZ 87535, "Cordillera Colán, (5°34'S, 78°19'W), Department of Amazonas, Peru, elevation 9,100 ft (ca. 2,775 m)." Adult male collected by Thomas S. Schulenberg on 23 August 1978. Original number TSS 962.

PARATYPES (13): LSUMZ 87531-34, 87536-44, type locality between 9550-10,000 ft., seven males and six females collected by TSS, L. J. Barkley, G. L. Graham, and Morris D. Williams on 20 August-14 September 1978.

COMMENTS: The complete locality on the holotype label reads "PERU: Dpto. Amazonas; Cordillera Colán, E La Peca (ridge W of peaks) 9600 [ft.]." Not mentioned in the type description are two fluid-preserved specimens (LSUMZ 89422-23) and four skeletal specimens (LSUMZ 89836-39) collected with the type series. Fjeldså and Krabbe (1990) recognized *parkeri*.

Metallura odomae Graves, 1980, Wilson Bull. 92: 3.

HOLOTYPE: LSUMZ 87547, "Cerro Chinguela, ca. 5 km northeast of Sapalache, 05°07'S, 79°23'W: Department of Piura, Peru, elevation approximately 2900 m (9500 ft)." Adult male netted and prepared by Richard D. Semba on 10 June 1978. Original number RDS 535.

PARATYPES (4): LSUMZ 87545-46, 87548-49, type locality, females collected on 6, 9, 16 (all prepared by RDS), and 24 June (collected by J. William Eley) 1976, respectively.

COMMENTS: Not mentioned in the type description is an unsexed skeleton (LSUMZ 89835) collected with the type series. Three study skins were subsequently obtained from the type locality 12-16 July 1980 by Michael J. Braun (LSUMZ 97615-17). This species is known only from Cerro Chinguela and from recent specimens from southern Ecuador (Collar et al. 1992).

PICIFORMES
FAMILY PICIDAE

Centurus carolinus perplexus Burleigh and Lowery, 1944, Occas. Pap. Mus. Zool. Louisiana State Univ. (17): 298.

HOLOTYPE: LSUMZ 3846, "Cape Sable, near Flamingo, Monroe County, Florida." Adult male collected by George H. Lowery, Jr. on 1 April 1940. Original number GHL 1837.

PARATYPES (4) : LSUMZ 3847, Key Largo, Monroe Co., Florida, male collected by GHL on 31 March 1940; LSUMZ 3848, northeast corner of Cape Sable, Monroe Co., other data same as holotype; LSUMZ 6249, Miami, Dade Co., Florida, female collected by T. D. Burleigh on 8

December 1929; LSUMZ 8360, collected by TDB, other data same as holotype.

COMMENTS: Currently placed in *Melanerpes*. Short (1982) considered *perplexus* to be the end-point in a cline rather than a valid taxon; however, he stated that "although slightly smaller and exhibiting tendencies towards whiter underparts, paler red nasal tufts, and less white in the tail than [*M. c.*] *carolinus*, [*perplexus*] seems insufficiently differentiated to merit recognition (it comes closest of these putative "races" to being recognizable nomenclatorially)."

Centurus aurifrons turneffensis Russell, 1963, Occas. Pap. Mus. Zool. Louisiana State Univ. (25): 1.

HOLOTYPE: LSUMZ 21054, "Calabash Cay, Turneffe Islands, British Honduras." Adult male collected by Stephen M. Russell on 10 April 1956, prepared by E. Barahona. Original number SMR 599.

PARATYPES (4): LSUMZ 21053, collected by D. Dennett, Jr., other data same as holotype; LSUMZ 21055-56, type locality, females collected by SMR on 13-14 April 1956, respectively; LSUMZ 21057, type locality, male collected by EB on 15 April 1956.

COMMENTS: The complete locality on the holotype label reads "BR. HONDURAS: Belize Dist.; Turneffe, Calabash Cay." Currently placed in *Melanerpes*. Short (1982) recognized *turneffensis*.

Sphyrapicus varius appalachiensis Ganier, 1954, Migrant 25: 40.

HOLOTYPE: LSUMZ 75956, "4400 feet altitude in the Unicoi Mountains, Monroe County, Tennessee." Adult male collected by Albert F. Ganier on 21 June 1946. Original number AFG 1426 (but see below).

PARATYPES (2): LSUMZ 75957-58, type locality, male and female collected by AFG on 23-24 June 1944, respectively.

COMMENTS: No paratypes were specifically designated in the type description, but we consider the above two specimens as valid paratypes because they were taken at the type locality prior to publication of the type description. The "holotype" at LSUMNS does not match the one in the type description; the original Ganier Collection specimen label indicates "type specimen-collected by Ganier" in ink, in Ganier's hand, but there are discrepancies in the specimen number (AFG 1411), date (18 June 1946), and elevation (4600 feet). Both AFG 1411 and 1426 are present in the photograph of a series of specimens that appeared in Ganier (1954).

Short (1982) did not recognize *appalachiensis*, "the supposedly darker Appalachian population being based on worn summer specimens (these tend to be blacker). The southern Appalachian birds may be a trifle darker and smaller than northern birds, but overlap is great."

PASSERIFORMES
FAMILY DENDROCOLAPTIDAE

Dendrocolaptes certhia sheffleri Binford, 1965, Occas. Pap. Mus. Zool. Louisiana State Univ. (30): 1.

HOLOTYPE: LSUMZ 24410, "19 mi. N Puerto Angel, Oaxaca, México, elevation 900 feet." Adult male collected by L. C. Binford on 4 May 1961. Original number LCB 400.

PARATYPE: LSUMZ 24409, type locality, female collected by L. L. Wolf on 5 May 1961.

COMMENTS: This rare, disjunct, distinctive taxon is endemic to the Sierra de Miahuatlán on the Pacific side of Oaxaca (Binford 1989).

FAMILY FURNARIIDAE

Schizoeaca fuliginosa plengei O'Neill and Parker, 1976, Bull. Brit. Orn. Club 96: 137.

HOLOTYPE: LSUMZ 74008, "Bosque Taprag above (NE) Acomayo, Departamento de Huánuco, Peru, elevation c. 3350 m." Adult male collected by Erika J. Tallman on 23 June 1973, prepared by Dan A. Tallman. Original number DAT 1328.

PARATYPES (10): LSUMZ 74007, S of Huaylaspampa, Bosque Cutirragra, ca. 9100 ft., male collected by DAT on 23 July 1973; LSUMZ 74009-10, Bosque Magrapampa below Zapatagocha, ca. 10,000 ft., male and female collected by J. P. O'Neill and DAT, respectively, on 19 June 1973; LSUMZ 74011, Huaylaspampa, ca. 8500 ft., male collected by ET on 10 July 1973; LSUMZ 75209-10, Bosque Zapatagocha above Acomayo, one sex? and one female collected by Manuel Villar on 26 November 1973; LSUMZ 80501, 80504-06, Unchog, pass between Churubamba and Hacienda Paty above Acomayo, ca 10,000-11,000 ft., three males and a female, respectively, collected by R. S. Kennedy, JPO, and MV on 12 June-19 July 1975.

COMMENTS: The locality "Bosque Taprag" is incorrectly given as "Bosque Tapra" on the holotype label. A fluid-preserved specimen (LSUMZ 75007) collected with the type series was not mentioned in the type description. Three other paratypes (LSUMZ 79684, 80502-03) have been exchanged.

Fieldwork by LSUMNS personnel in Peru has now extended the range of this taxon north about 260 km to central depto. San Martín, close to the southern limit of *S. f. peruviana*. New localities are: Puerta del Monte, ca. 30 km NE Los Alisos, 3250 m, depto. San Martín (n = 8); Quebrada La Caldera, 7 km NE Tayabamba, 3550 m, depto. La Libertad (n = 3); and Mashua, east of Tayabamba, on trail to Ongón, 3350 m, depto. La Libertad (n = 12). Recent LSUMNS fieldwork has also extended the range of *plengei* south about 80 km, south of the Río Huallaga canyon, to northern depto. Pasco: 1 km east of Huánuco border on Pozuzo-Panao trail, 3700 m (n = 2); 2 km NW Punta Saria on Pozuzo-Chaglla trail, 3100 m (n = 3); and

Millpo, east of Tambo de Vacas on Pozuzo-Chaglla trail, 3450 m (n = 13). Also, 18 additional specimens have been taken by LSUMNS personnel in depto. Huánuco, including a locality on the south side of the Río Huallaga. Fjeldså and Krabbe (1990) recognized *plengei*.

Schizoeaca harterti bejaranoi Remsen, 1981, Proc. Biol. Soc. Washington 94: 1068.

HOLOTYPE: LSUMZ 36040, "kilometer 104, Prov. Chapare, Dpto. Cochabamba, Bolivia, 3200 m elevation (about 55 km ENE of the city of Cochabamba)." Male collected by F. Steinbach on 3 September 1962. Original number FS 14,375.

PARATYPES (20): LSUMZ 36027-39, 36041, 37611-37615, type locality, 10 males, 2 male?, 5 females, 2 sex? collected by FS between 17 February 1960-4 September 1962; LSUMZ 37616, Choro, Prov. Ayopaya, 3500 m, male collected by FS on 13 September 1953.

COMMENTS: Fjeldså and Krabbe (1990) recognized *bejaranoi*.

FAMILY FORMICARIIDAE

Herpsilochmus parkeri Davis and O'Neill, 1986, Wilson Bull. 98: 338.

HOLOTYPE: LSUMZ 116908, "ca 15 km by trail northeast of Jirillo on the trail to Balsapuerto, 06°03'S, 76°44'W, 1350 m, Department of San Martín, Peru." Adult male collected by Tristan J. Davis on 20 November 1983. Original number TJD 3051.

PARATYPES (9): LSUMZ 116901-07 (skins), 117994 (fluid-preserved), 118245 (skeleton), type locality, five males and four females collected by TJD, A. Urbay T., and J. P. O'Neill on 27 October-20 November 1983.

COMMENTS: This species is still known only from the type locality. Vuilleumier et al. (1992) favored treatment of this taxon as a subspecies of *H. pileatus*; presumably they are following a traditional classification and would include *atricapillus* and *motacilloides* in this polytypic species, although they are not explicit on this point. Davis and O'Neill (1986) followed Naumberg (1935, 1939), who pointed out that the distribution of nominate *pileatus* was entirely surrounded by that of *atricapillus*, and hence species status for both was indicated. Recent field-work has confirmed that *pileatus* and *atricapillus* are locally syntopic and that they are clearly different species (T. S. Schulenberg, in litt.). The Andean taxa, *motacilloides* and *parkeri*, are allopatric to both *atricapillus* and *pileatus*, and differ greatly in habitat from the latter two. As noted by Davis and O'Neill, *motacilloides* and *parkeri* differ in plumage as much or more as several pairs of sympatric species of *Herpsilochmus*. Furthermore, the songs of *parkeri*, *motacilloides*, *atricapillus*, and *pileatus* are quite distinct (T. S. Schulenberg, pers. comm.), consistent with Davis and O'Neill's recognition of all four taxa as species.

Grallaria eludens Lowery and O'Neill, 1969, Auk 86: 1.

HOLOTYPE: LSUMZ 62312, "Balta (at the point where the streams known to the local Cashinahua Indians as the Xumuya and the Inuya enter the Río Curanja), 10°08'S, 71°13'W, elevation ca. 300 m, Depto. Loreto, Peru." Adult male collected by J. P. O'Neill on 11 July 1967. Original number JPO 2623.

PARATYPES (10): LSUMZ 52135, collected by JPO on 24 July 1966; LSUMZ 62308 (skeleton and partial skin), 62311, 62313-15, collected by JPO between 13-29 July 1967; LSUMZ 62316-17, collected by James J. Hebrard on 21 and 17 July 1967, respectively; LSUMZ 62818-19 (skeletons), collected by JPO on 19-20 July 1967, respectively. All are from the type locality, and all are males except 62314 (female?) and 62819 (sex?).

COMMENTS: The type locality is now in depto. Ucayali. Three other paratypes (LSUMZ 62309-10, 62318) have been exchanged. A male (LSUMZ) collected by A. S. Meyer on the southeastern slope of Cerro Tahuayo, ca. 65 km ENE Pucallpa, depto. Ucayali (about 370 km northwest of the type locality), on 24 July 1987, is the only record away from the type locality.

Grallaria carrikeri Schulenberg and Williams, 1982, Wilson Bull. 94: 105.

HOLOTYPE: LSUMZ 88044 (skin and partial skeleton), "Cordillera Colán, SE La Peca, ca. 5°34'S, 78°19'W, 2450 m, Dept. Amazonas, Peru." Adult male collected by Morris D. Williams on 15 October 1978. Original number MDW 2056.

PARATYPES (11): LSUMZ 88042-43 (males), 88045-46 (male and female), 89957 (female skeleton), type locality between 7700-8400 ft., collected by Manuel Sánchez S., T. S. Schulenberg, and MDW on 10-21 October 1978; LSUMZ 92456-61, Cumpang, above Utcubamba, on trail to Ongón, depto. La Libertad, 2750-2900 m, three males and three females collected by T. A. Parker, III, Mark B. Robbins, and D. Wiedenfeld on 12-28 October 1979.

COMMENTS: The nestlings from Cumpang mentioned in the type description are LSUMZ 91513-14 (fluid-preserved; Wiedenfeld 1982). No new locality records have been obtained since the type description.

Grallaria blakei Graves, 1987, Wilson Bull. 99: 314.

HOLOTYPE: LSUMZ 64228, "east slope of the Cordillera Carpish, near the Carretera Central, ca 2400 m, Dpto. Huánuco, Peru." Adult female collected by J. P. O'Neill on 17 August 1968. Original number JPO 3292.

PARATYPES (6): LSUMZ 74096, 74098, 75243, trail to Hacienda Paty below Carpish Pass, Cordillera Carpish, depto. Huánuco, 7000-7700 ft., female collected by C. S. O'Neill on 10 August 1973 and two males collected by Dan A. Tallman on 2 August and 22 November 1973, respectively; LSUMZ 88068, Cordillera Colán, depto. Amazonas, female collected by T. S. Schulenberg on 17 October 1978; LSUMZ 106081, Cumbre de Ollón, ca. 12 km E Oxapampa, depto. Pasco, 2500 m, female collected by D.

Wiedenfeld on 15 March 1982; LSUMZ 116973, 30 road km E Florida, depto. Amazonas, female collected by Tristan J. Davis on 29 November 1983.

COMMENTS: LSUMZ 75098 is listed as a paratype in the type description, but that number belongs to a specimen of *Calidris alba*; LSUMZ 106081 (above) should be listed instead.

Grallaricula ochraceifrons Graves, O'Neill, and Parker, 1983, Wilson Bull. 95: 1.

HOLOTYPE: LSUMZ 81998, "10 km by road below (NE) Abra Patricia, elev. approx. 1890 m (6200 ft) 5°46'S, 77°41'W; Dpto. San Martín, Peru." Adult male prepared by J. P. O'Neill on 30 August 1976. Original number JPO 5715.

PARATYPES (4): LSUMZ 81996-97, type locality, female and male collected by JPO on 26 August 1976; LSUMZ 88092-93, ca. 20 trail km E La Peca [Cordillera Colán], depto. Amazonas, Peru, 6400 and 6500 ft., males collected by G. L. Graham and T. S. Schulenberg on 17-18 August 1978, respectively.

COMMENTS: This species is still known only from the type and paratype localities.

FAMILY TYRANNIDAE

Uromyias agraphia squamigera O'Neill and Parker, 1976, Bull. Brit. Orn. Club 96: 140.

HOLOTYPE: LSUMZ 74301, "Bosque Cutirragra, south of Huaylaspampa, Departamento de Huánuco, Peru, elevation c 2775 m." Adult male collected by Erika Tallman on 21 July 1973, prepared by Dan A. Tallman. Original number DAT 1552.

PARATYPE: LSUMZ 79704, Punta de Esperanza, depto. Huánuco, male collected by T. Parker on 7 February 1975.

COMMENTS: A fluid-preserved specimen, LSUMZ 79602 (see Schulenberg and Graham 1981, but number mistakenly given as "76902" therein), taken with the paratype was not mentioned in the type description. Traylor (1979) recognized *squamigera*. *Uromyias* was placed by Traylor (1979) in *Anairetes*, but Lanyon's (1988) analysis of syringeal morphology and other characters indicates that *Uromyias* and *Anairetes* are not sister taxa.

Anairetes agraphia plengei Schulenberg and Graham, 1981, Bull. Brit. Orn. Club 101: 242.

HOLOTYPE: LSUMZ 88474 (skin and partial skeleton), "Cordillera Colán, NE La Peca, about 5°34'S, 78°19'W, Department of Amazonas, Peru, elevation 3025 m." Adult male collected by T. S. Schulenberg on 29 August 1978. Original number TSS 988.

PARATYPES (7): LSUMZ 88470 (male skin and partial skeleton), 88471-73 (female, two males), 90079-81 (skeletons; two males, sex ?), type locality,

collected by L. J. Barkley, G. L. Graham, TSS, and Morris D. Williams on 29 August-16 September 1978.

Rhynchocyclus brevirostris pallidus Binford, 1965, Occas. Pap. Mus. Zool. Louisiana State Univ. (30): 5.

HOLOTYPE: LSUMZ 33238, "Minitán, 22 road mi. S Pinotepa Nacional, Oaxaca, México, sea level." Male collected by L. C. Binford on 1 March 1964. Original number LCB 945.

PARATYPES (7): 33239, type locality, female collected by LCB on 28 February 1964; LSUMZ 33240, 33243, 3 mi. N Pluma Hidalgo, 5000 ft., males collected by LCB and John J. Morony, respectively, on 1 May 1964; LSUMZ 33241-42, 1 mi. E Putla, 2400 ft., males collected by LCB and JJM, respectively, on 21 May 1964; LSUMZ 33244-45, 18 road mi. N San Gabriel Mixtepec, 4900 ft., females collected by JJM on 8-9 May 1964, respectively.

COMMENTS: T aylor (1979) recognized *pallidus*.

FAMILY COTINGIDAE

Pipreola riefferii tallmanorum O'Neill and Parker, 1981, Bull. Brit. Orn. Club 101: 294.

HOLOTYPE: LSUMZ 74145, "PERU: Department of Huánuco; on the trail to Hacienda Paty below (NE) Carpish Pass, elevation c 2200 m." Adult male collected by Erika Tallman on 5 August 1973, prepared by D. A. Tallman. Original number DAT 1734.

PARATYPES (11): LSUMZ 74140-44, 74146-49, type locality, six males and three females collected by R. J. Louque, C. S. O'Neill, J. P. O'Neill, DAT, and ET on 2-11 August 1973; LSUMZ 75246, type locality, female collected by ET on 22 November 1973; LSUMZ 80608, type locality, male collected by Robert S. Kennedy on 25 June 1975.

COMMENTS: A fluid-preserved specimen (LSUMZ 75017) taken with the type series was not mentioned in the type description. Fjeldså and Krabbe (1990) recognized *tallmanorum*.

Conioptilon mcilhennyi Lowery and O'Neill, 1966, Auk 83: 3.

HOLOTYPE: LSUMZ 42781, "Balta (at the point where the streams known to the local Cashinahua Indians as the Xumuya and the Inuya enter the Río Curanja), lat. 10°08'S, long. 71°13'W, elevation approximately 300 meters, Depto. Loreto, Peru." Adult male collected by John P. O'Neill on 18 March 1965. Original number JPO 1452.

PARATYPES (8): LSUMZ 35228-29, 42784, 42788, 42866 (fluid preserved), 42870-71 (both fluid-preserved), 42914 (male skeleton), type locality, collected by JPO 30 December 1964-21 March 1965 (females unless otherwise noted).

COMMENTS: A new genus (Auk 83: 2) and species. Five additional paratypes (LSUMZ 42782-83, 42785-87) have been exchanged. Another paratype, LSUMZ 42900, could not be located in the LSUMNS collection.

Another four skins (LSUMZ 52146-49, two males, two females) and one fluid-preserved specimen (LSUMZ 63727) were collected at the type locality by JPO and J. J. Morony, 22 June-22 July 1966, and an additional four skins (LSUMZ 63796 [now exchanged], 64245-47, two males, two females), one skeleton (LSUMZ 65024, female), and one fluid-preserved specimen (LSUMZ 64793) were obtained there by JPO, 10 June-23 July 1968.

FAMILY HIRUNDINIDAE

Hirundo albifrons ganieri Phillips, 1986, The Known Birds of North and Middle America, p. 34.

LECTOTYPE: LSUMZ 76133, "Swallow Bluff, Decatur Co., Tennessee." Adult male collected by A. F. Ganier on 18 May 1941. Original number AFG 916.

PARALECTOTYPES (7): LSUMZ 76130-32, 76134, 76136-38, five males and two females, other data same as for lectotype.

COMMENTS: Phillips (1986) stated that the type was a male from "Swallow Bluff, Decatur Co., Tennessee, 18 May; Wg 105, T. 47" (no year given). However, he did not designate a holotype by museum or specimen number, either in the original description or on specimen labels themselves. The description is almost certainly based on a series of eight specimens (LSUMZ 76130-34; 76136-38) taken by A. F. Ganier at this locality on 18 May 1941; six are males. Two additional specimens (LSUMZ 76139-40, both marked "nesting bird") were taken on the same date in Decatur Co. but not specifically designated by locality. An additional specimen collected at the type locality on 18 May 1941 (formerly LSUMZ 76135) is now housed at the Museo Javier Prado, Lima, Peru. Because Phillips (1986) did not designate a specific specimen as the holotype, we hereby designate LSUMZ 76133 as the lectotype.

Most recent treatments use the name *Hirundo pyrrhonota* for this species. We are uncertain of the validity of *ganieri*. Phillips' (1986) entire description of this taxon consists only of "Forehead buffy to buffy whitish; cheeks and throat dark chestnut as in *albifrons*; Wg 102-108 (110)." The characters given by Phillips for geographically adjacent *H. a. albifrons* are extremely similar to those that he provided for *ganieri*, i.e., forehead "buffy to whitish" and male wing length "106.5-110.5." Phillips gave no sample sizes for any mensural data.

FAMILY TROGLODYTIDAE

- Cistothorus platensis potosinus* Dickerman, 1975, Amer. Mus. Novitates (2569): 4.
HOLOTYPE: LSUMZ 33575, "Laguna Manantiales Media Luna, San Luis Potosí, Mexico." Adult male collected by L. C. Binford on 9 February 1964. Original number LCB 863.
PARATYPES (13): LSUMZ 18656-60, 19838, 19846, 19987, 33572-74, 33576, 52524, type locality, seven males and six females collected by L. Guerrero, LCB, and R. W. Dickerman from 1952-1965.
COMMENTS: Phillips (1986) recognized *potosinus*.

- Cistothorus platensis russelli* Dickerman, 1975, Amer. Mus. Novitates (2569): 6.
HOLOTYPE: LSUMZ 21238, "Augustine, Cayo District, British Honduras." Adult male collected by Stephen M. Russell on 9 March 1956. Original number SMR 474.
PARATYPES (17): LSUMZ 20726-27, 21 mi. SSE Cayo on eastern branch of Belize River, Cayo Dist., male and female collected by SMR and D. A. Lancaster on 26 and 28 April 1955, respectively; LSUMZ 21239, 5 mi. W Baldy Beacon, Cayo Dist., male collected by SMR on 20 April 1956; LSUMZ 21240, Baldy Beacon, Cayo Dist., male collected by SMR on 21 April 1956; LSUMZ 31163-65, 18 mi. NW Monkey River Town, Toledo Dist., males collected by SMR on 18 June 1963; LSUMZ 31166, 31168-69, 17 mi. NW Monkey River Town, Toledo Dist., males collected by SMR on 19 June 1963; LSUMZ 31170, 16 mi. NW Monkey River Town, Toledo Dist., male collected by SMR on 23 June 1963; LSUMZ 31171-31174, Sibun Hill, Cayo Dist., two males and two females collected by A. Palmisano on 20 July 1963; LSUMZ 31176, Sibun Hill, male collected by AP on 21 July 1963; LSUMZ 31177, 1 mi. SSW Cooma Cairn, Cayo Dist., male collected by AP on 7 July 1963.
COMMENTS: Phillips (1986) recognized *russelli*.

- Cistothorus platensis graberi* Dickerman, 1975, Amer. Mus. Novitates (2569): 7.
HOLOTYPE: LSUMZ 32356, "Puerto Lempira, Department of Gracias a Dios, Honduras." Adult male collected by R. R. Graber on 12 April 1964.
PARATYPES (3): LSUMZ 29832, 32355, type locality, females collected by B. L. Monroe on 7 February 1963 and 14 April 1964, respectively; LSUMZ 32357, female, other data same as holotype.
COMMENTS: From the holotype label, the elevation of the type locality is sea level. A fourth paratype (LSUMZ 32358) has been exchanged. Phillips (1986) recognized *graber*.

- Thryothorus euophrys schulenbergi* Parker and O'Neill, 1985, Ornithol. Monogr. (36): 12.
HOLOTYPE: LSUMZ 88579, "Cordillera Colán, SE La Peca, ca. 5°34'S, 78°19'W, elevation 2713 m, Department of Amazonas, Peru." Adult male

collected by Thomas S. Schulenberg on 12 October 1978. Original number TSS 1196.

PARATYPES (4): LSUMZ 88577-78, Cordillera Colán, 8500-8700 ft., females collected by Morris D. Williams and TSS, respectively, on 8 October 1978; LSUMZ 104540-41, Puerta del Monte, ca. 30 km NE Los Alisos, depto. San Martín, ca. 3250 m, males collected by Cesar Salirrosas and Susan Allen-Stotz on 6 and 10 August 1981, respectively.

COMMENTS: Not mentioned in the type description are two skeletons (LSUMZ 90127-28, female, male) collected with the type series at the type locality. Fjeldså and Krabbe (1990) recognized *schulenbergi*.

Thryothorus eisenmanni Parker and O'Neill, 1985, Ornithol. Monogr. (36): 9.

HOLOTYPE: LSUMZ 78913, "San Luís on Ollantaitambo-Quillabamba road, above Huyro, 13°06'S, 72°25'W, elevation "9000 ft" [2744 m], Department of Cuzco, Peru." Adult male collected by John P. O'Neill on 4 August 1974. Original number JPO 4878.

PARATYPES (14): LSUMZ 78910-12, Machu Picchu, depto. Cuzco, 7000 ft., two males and one female collected by JPO, Gary Lester, and T. A. Parker, III, respectively, on 21 July 1974; LSUMZ 78914, 78916-19, type locality, two males and two females collected by GL, TAP, K. R. Thomas, and Reyes Rivera on 4-20 August 1974; LSUMZ 78915, below San Luís on Ollantaitambo-Quillabamba road, ca. 7800 ft., male collected by GL on 6 August 1974; LSUMZ 78920, Hacienda Huyro between Huayopata and Quillabamba, ca. 6000 ft., male collected by R. Thomas on 26 August 1974; LSUMZ 78921-24, Bosque Aputinye above Huyro, ca. 6000 ft., two males and two females collected by TAP on 27-28 August 1974.

COMMENTS: The "specimens examined" section of the type description should be corrected as follows. The three male paratypes (excluding holotype) from the vicinity of San Luís are LSUMZ 78915 and 78917-18 (not "78913-78915"). There are three (not four) females from San Luís and their numbers are LSUMZ 78914, 78916, and 78919 (not "78619-78919"). There are three (not two) specimens from Machu Picchu - two (not one) males, LSUMZ 78910-11 (not "78912"), and one female, LSUMZ 78912 (not "78913"). Not mentioned in the type description is a skeleton (LSUMZ 79847) collected with the type series at the type locality.

Vuilleumier et al. (1992), after listing several opinions to the contrary from those familiar with the taxa involved, stated that "We believe, however, that *eisenmanni* is a well-marked subspecies of "*euophrys*" They did not present factual reasons for their decision, nor did they address any of Parker and O'Neill's (1985) points in favor of species status for *eisenmanni*. We note that *Thryothorus eisenmanni* differs from other members of the *T. euophrys* group more so than certain other South American *Thryothorus* currently treated as species differ from each other, e.g., *T. coraya* and *T. genibarbis*, and *T. leucotis* and *T. guarayanus*. In the

absence of arguments to the contrary, we recommend treatment of *eisenmanni* as a species.

Thryothorus ludovicianus tropicalis Lowery and Newman, 1949, Occas. Pap. Mus. Zool. Louisiana State Univ. (22): 4.

HOLOTYPE: LSUMZ 11823, ".5 mi. E Naranjos, San Luis Potosí, México; elevation 1,000 feet." Adult male collected by Robert J. Newman on 20 March 1947, prepared by Marcella Newman. Original number RJN 563.

PARATYPES (2): LSUMZ 11824, female, other data same as holotype; LSUMZ 11825, 3 mi. W Naranjo on hwy., 0.6 mi. E of El Salto, male collected by George H. Lowery, Jr. on 13 February 1948.

COMMENTS: The complete locality on the holotype label reads "MEXICO: San Luis Potosí; Naranjo Region, Sabinito, 6 1/2 mi. E. at El Salto turn-off, 1/2 mi. E Naranjos, 1000 ft." Paynter (1960) and Phillips (1986) recognized *tropicalis*.

Thryothorus ludovicianus tabascensis Lowery and Berrett, 1963, Occas. Pap. Mus. Zool. Louisiana State Univ. (24): 1.

HOLOTYPE: LSUMZ 27239, "8 mi. SE Tenosique, Tabasco, México." Immature male collected by D. G. Berrett on 10 November 1961. Original number DGB 884.

PARATYPES (2): LSUMZ 27238, 27240, type locality, female and male collected by DGB and L. C. Binford on 7-8 November 1961, respectively.

COMMENTS: Phillips (1986) was uncertain of the validity of *tabascensis*, which may still be known only from the type series. Phillips (1986) stated that he examined the holotype and "the 3 Tabasco specimens"; because there were only two additional paratypes, we are uncertain whether an additional specimen exists of which we are not aware.

Phillips (1986) placed the southern populations of *T. ludovicianus*, including those of Tabasco, in a separate species, "*Thryothorus*?" *albinucha*, which he suspected belonged in a separate genus. A tape-recorded song of *albinucha* from Uxmal, Yucatán, by Coffey and Coffey (1989) sounds to us to be similar to that of Louisiana *T. ludovicianus* except that it is longer, ca. 9 notes instead of 4-5, and faster. It is not indicated on the recording whether the song was in response to playback; such songs are often longer and faster in many bird species. We have no data relevant to whether *albinucha* merits recognition as a species. We also hesitate to make any conclusions from a single recording. Nonetheless, the similarity in the songs is so great that we believe that *albinucha* is not only congeneric with but also the sister taxon to North American *ludovicianus*.

Henicorhina leucosticta costaricensis Dickerman, 1973, Condor 75: 362.

HOLOTYPE: LSUMZ 32665, "Turrialba, Cartago Province, Costa Rica." Adult(?) male collected by D. G. Huckaby on 9 November 1963. Original number DGH 67.

PARATYPES (3): LSUMZ 30701, type locality, male collected by K. A. Arnold on 7 December 1962; LSUMZ 32666, female, other data same as holotype; LSUMZ 35032, type locality, male collected by KAA on 23 September 1964.

COMMENTS: The complete locality on the holotype label reads "Costa Rica: , Prov. Cartago; Turrialba, IICA, 600 m." Phillips (1986) recognized *costaricensis*.

FAMILY MIMIDAE

Margarops fuscus hypenemus Buden, 1993, Bull. Brit. Orn. Club 113: 82.

HOLOTYPE: LSUMZ 142124, "1.5 mi. (2.4 km) SW Lodge, St. Mary Cayon Parish, St. Christopher (=St. Kitts), West Indies." Male collected by Albert Schwartz on 18 April 1962. Original number AS 4349.

PARATYPES (25): LSUMZ 142574-78 (AS 4526-30), west slope, Mt. Scenery, 1200 ft., Saba, two males and 3 females collected by AS and R. F. Klinikowski on 16 May 1962; LSUMZ 142579-81 (AS 4350-52), type locality, female and two males collected by AS and RFK on 18 April 1962; LSUMZ 142582 (AS 4480), 3 mi. SW Molyneux, 1600 ft., Christ Church Nichola Town Parish, St. Christopher, female collected by AS on 3 May 1962; LSUMZ 142583 (AS 4486), 2.5 mi. SW Molyneux, male collected by AS on 4 May 1962; LSUMZ 142584-85 (AS 4365-66), Golden Rock Estate, St. George Gingerland Parish, Nevis, males collected by AS on 20 April 1962; LSUMZ 142586-89 (AS 4415-18), Golden Rock Estate, two males and two females collected by AS on 23 April 1962; LSUMZ 142590 (AS 4110), Wallings Reservoir, St. Mary Parish, Antigua, male collected by RFK on 20 March 1962; LSUMZ 142591-92 (AS 4250,4269), west slope, South Soufriere, 1400 ft., St. Anthony's Parish, Montserrat, males collected by RFK on 5 & 7 April 1962; LSUMZ 142593-94 (AS 4278-79), 1 mi. N Salem, St. Peter's Par., Montserrat, female and male collected by RFK on 7 April 1962; LSUMZ 142595 (AS 5686), 1 km S Mahaudière, Guadeloupe, sex? collected by AS on 27 February 1963; LSUMZ 142596 (AS 5669), 0.5 km NE Grande Anse, Désirade, sex ? collected by R. Thomas on 28 February 1963; LSUMZ 142597-98 (AS 5661-62), 5 km SE Grand-Bourg, Marie-Galante, females collected by RT on 1 March 1963.

COMMENTS: Sometimes placed in the genus *Allenia*.

Margarops fuscus schwartzi Buden, 1993, Bull. Brit. Orn. Club 113: 83.

HOLOTYPE: LSUMZ 142125, "L'Hermitage, Dauphin Quarter, St. Lucia, West Indies." Male collected by Albert Schwartz on 18 January 1962. Original number AS 3532.

PARATYPES (22): LSUMZ 142626-30 (AS 3439-43), type locality, males collected by R. F. Klinikowski on 7 January 1962; LSUMZ 142631-32 (AS 3530-31), type locality, males collected by RFK on 18 January 1962; LSUMZ 142633, 142635 (AS 5730,5761), Petite Anse River, males collected

by AS on 3 & 8 April 1963, respectively; LSUMZ 142637-38 (AS 5729, 5768), 1.5 mi. NW Grande Anse, collected by R. Thomas and AS on 3 & 9 April 1963, respectively; LSUMZ 142639-40 (AS 5785, 5787), La Sorciere, north slope, 600 ft., males collected by AS and RT on 13 & 14 April 1963, respectively; LSUMZ 142641-42 (AS 5759-60), De Barra, males collected by AS and RT on 7 April 1963; LSUMZ 142643-44 (AS 5777, 5784), De Barra, males collected by AS and RT on 10 & 13 April 1963, respectively; LSUMZ 142645 (AS 3480), 3 mi. SE Canaries, Anse-la-Ray Quarter, male collected by RFK on 12 January 1962; LSUMZ 142646 (AS 3511), Desqaitiers, Choiseue Quarter, male collected by AS on 16 January 1962; LSUMZ 142647 (AS 5794), Venus Estate, Anse la Ray Quarter, female collected by AS on 16 April 1963; LSUMZ 142648 (AS 5793), 2.5 mi. SW Duraudeau, Anse la Ray Quarter, female collected by RT on 16 April 1963; LSUMZ 142649 (AS 5786), 3 mi. SW Réduit, Gros Islet Quarter, male collected by AS on 14 April 1963.

COMMENTS: Sometimes placed in the genus *Allenia*. Two unsexed specimens were not used as paratypes: LSUMZ 142634 (AS 5731, same data as LSUMZ 142633) and LSUMZ 142636 (AS 5762, same data as LSUMZ 142635).

FAMILY MUSCICAPIDAE
SUBFAMILY TURDINAE

Turdus grayi microrhynchus Lowery and Newman, 1949, Occas. Pap. Mus. Zool. Louisiana State Univ. (22): 5.

HOLOTYPE: LSUMZ 12008, "Santa María del Río, San Luis Potosí, México; elevation 5,500 feet." Adult male collected by Robert J. Newman on 13 March 1948, prepared by Marcella Newman. Original number RJN 1098.

PARATYPES (7): LSUMZ 12005-06, type locality, males collected by G. H. Lowery, Jr. on 30 August 1947; LSUMZ 12007, type locality, male collected by RJN on 30 August 1947; 12009-11, type locality, two females and one male, respectively, collected by RJN on 13 March 1948; LSUMZ 12012, Labor del Río, 5100 ft., male collected by RJN on 10 August 1947.

COMMENTS: Ripley (1964) and Phillips (1991) recognized *microrhynchus*.

SUBFAMILY MONARCHINAE

Chasiempis sandwichensis bryani Pratt, 1979, Bull. Brit. Orn. Club 99: 106.

HOLOTYPE: LSUMZ 81725, "Puu Laau, Hamakua District, Hawaii County, Hawaii, elevation c. 1950 m." Adult male collected by H. Douglas Pratt on 27 August 1976. Original number HDP 545.

PARATYPES (21): LSUMZ 81709-24, 81726-27, type locality, eleven males, five females, two sex? collected by HDP on 16-27 August 1976; LSUMZ 87023-25, type locality, male and two females collected by HDP on 5 February 1978.

COMMENTS: Watson et al. (1986) recognized *bryani*.

FAMILY PARIDAE

Penthestes hudsonicus cascadiensis Miller, 1943, Occas. Pap. Mus. Zool. Louisiana State Univ. (14): 262.

HOLOTYPE: LSUMZ 3270, "1 mile south of Monument 83 (long. 120° 38 1/2') on the United States-Canadian boundary, 6,000 feet, Okanogan Co., Washington." Adult male collected by A. W. Burdick on 23 August 1942. Original number AWB 371.

PARATYPES (2): LSUMZ 3267, 3269, collected on 11 July and 12 August 1942, other data same as holotype.

COMMENTS: Currently placed in *Parus*. Snow (1967) recognized *cascadiensis*. Phillips (1986) synonymized *cascadiensis* with *P. h. columbianus*; however, given Phillips' apparent lack of objectivity in evaluating subspecies described by A. H. Miller, we suggest that the validity of *cascadiensis* warrants independent evaluation.

FAMILY VIREONIDAE

Vireo crassirostris stalagmium Buden, 1985, Proc. Biol. Soc. Washington 98: 595.

HOLOTYPE: LSUMZ 70838, "Caicos Islands, North Caicos, along Flamingo Pond Road, east of Kew." Adult male collected by D. W. Buden on 29 February 1972. Original number DWB 41.

PARATYPES (5): LSUMZ 70369-70, Flamingo Pond Rd. between Kew and Flamingo Pond, males collected by DWB on 8 June 1971; LSUMZ 70371, trail between Kew and Sandy Point, male collected by DWB on 7 June 1971; LSUMZ 70372, 70839, Kew, male and sex?, collected by DWB on 7 June 1971 and 8 February 1972, respectively.

FAMILY EMBERIZIDAE SUBFAMILY EMBERIZINAE

Aimophila ruficeps tenuirostra Burleigh and Lowery, 1939, Occas. Pap. Mus. Zool. Louisiana State Univ. (6): 67.

HOLOTYPE: LSUMZ 3334, "Guadalupe Mountains, Culberson County, Texas; McKittrick Canyon, altitude 5,500 feet." Adult male collected by George H. Lowery, Jr. on 3 January 1939. Original number GHL 1293.

PARATYPES (3): LSUMZ 3511-12, 3514, "type locality," male, female, and unsexed juvenile collected by GHL on 1 January 1939, 8 October 1938, and 29 July 1938, respectively.

COMMENTS: The locality on the holotype label reads "Texas: Frijole, 5500 ft., McKittrick Canyon." Although the paratypes are from the type locality in the sense that they are from the Guadalupe Mountains, their specific localities are sufficiently different from the holotype to be noted here:

LSUMZ 3511, "Frijole, Pine Spring Canyon, 6000 ft.;" LSUMZ 3512, "Frijole, Pine Springs Canyon, 5800 ft.;" LSUMZ 3514, "Guadalupe Mts., Pine Canyon, 5600 ft." LSUMZ 3511, originally labeled as a female, is re-identified as a male for unexplained reasons. Paynter (1970) synonymized *tenuirostra* with *A. r. eremoeca*.

SUBFAMILY THRAUPINAE

Chlorospingus ophthalmicus hiaticolus O'Neill and Parker, 1981, Bull. Brit. Orn. Club 101: 296.

HOLOTYPE: LSUMZ 82222, "10 km (by road) below (NE) Abra Patricia on the road to Rioja, Departamento de San Martín, Peru, elevation c. 1890 m." Adult male collected by J. P. O'Neill on 24 August 1976. Original number JPO 5657.

PARATYPES (22): LSUMZ 64603-06, E. slope Cordillera Carpish, ca. 2400 m, depto. Huánuco, two females and two males collected by JPO on 22 & 21 August 1968, respectively; LSUMZ 74683, 75388, trail to Hacienda Paty below Carpish Pass, male and female collected by D. A. Tallman on 8 August and 22 November 1973, respectively; LSUMZ 82219-21, 82223-24, type locality, three males, two females, one sex? collected by Gary R. Graves and JPO on 22-26 August 1976; LSUMZ 88863-67, ca. 20 trail km E La Peca, Cordillera Colán, depto. Amazonas, 6550-6625 ft., males collected by G. L. Graham, David R. Hunter, and T. S. Schulenberg on 10-17 August 1978; LSUMZ 93226, 93228-29, 93231-33, Cumpang, depto. La Libertad, 2450-2800 m, two males, three females, and one sex? collected by J. William Eley, T. A. Parker, III, Mark B. Robbins, and D. Wiedenfeld on 8 October-5 November 1979.

COMMENTS: Two skeletons taken with the type series at the type locality (LSUMZ 84106-07, male, female), and another collected with the series from Cordillera Colán (LSUMZ 90200) were not mentioned in the type description. Three additional paratypes (LSUMZ 93225, 93227, 93230) have been exchanged.

The discovery and description of this taxon is a good example of how much progress has been made in the knowledge of Andean bird distribution in the last two decades. As recently as 1970 (Storer 1970), no form of *Chlorospingus ophthalmicus* was known in central Peru from the south side of the Marañón River south to depto. Junín, a distance of at least 725 km. Now, *hiaticolus* is known from throughout this range except in southern depto. Pasco, where the next subspecies to the south, *C. o. cinereocephalus*, recently has been recorded (Isler and Isler 1987).

As noted by Isler and Isler (1987) and Fjeldså and Krabbe (1990), a series of specimens from northern depto. Pasco is intermediate between *hiaticolus* and *cinereocephalus*; their breast bands are a paler shade of yellow than in typical *hiaticolus*, with one specimen as washed out as one of the LSUMNS specimens of *cinereocephalus* from southern depto. Pasco. The

locality of the intermediate specimens is "Playa Pampa, ca. 8 km NW Cushi on trail to Chaglla, 2100 m"; the seven specimens (LSUMZ) were collected by G. H. Rosenberg, K. V. Rosenberg, and T. S. Schulenberg from 14 June to 4 July 1985. The LSUMNS specimens (n = 16) of *cinereocephalus* from southern depto. Pasco mentioned by Isler and Isler (1987) were collected at "ca. 9 km SSE Oxapampa, 2050 m" from 13 February to 8 March 1982 by Manuel Sánchez S., TSS, Susan Allen-Stotz, and David A. Wiedenfeld, and at "Cumbre de Ollón, 12 km E Oxapampa, 2500 m" on 25 March 1982 by DAW.

Hemispingus rufosuperciliaris Blake and Hocking, 1974, Wilson Bull. 86: 321.

HOLOTYPE: LSUMZ 74727, "Bosque Huaylaspampa, Depto. Huánuco, Peru, approximately 8,400 ft." Adult male collected by D. A. Tallman on 18 July 1973. Original number DAT 1524.

PARATYPES (6): LSUMZ 74724, between Zapatagocha and Huaylaspampa, ca. 9500 ft., female collected by Erika Tallman on 29 June 1973; LSUMZ 74725, 75062 (fluid-preserved), Bosque Cutirragra, S of Huaylaspampa, ca. 9100 ft., collected by C. S. O'Neill and ET on 15 and 22 July 1973, respectively; LSUMZ 74729, Bosque Magrapampa, below Zapatagocha, ca. 8900 ft., female collected by DAT on 19 June 1973; LSUMZ 75391-92, Huaylaspampa, male and female collected by Manuel Villar on 18 November 1973.

COMMENTS: An unsexed skeleton (LSUMZ 75649) collected with the above paratypes was not mentioned in the type description. Two additional paratypes (LSUMZ 74726, 74728) have been exchanged.

As indicated by Isler and Isler (1987), subsequent LSUMZ fieldwork has extended the range of *H. rufosuperciliaris* north to dptos. La Libertad, San Martín, and Amazonas, more than 550 km northwest of the type locality. We suspect that the distribution of this species is continuous in elfin cloud forest from north of the Río Huallaga, depto. Huánuco, to just south of the Río Marañón in depto. Amazonas, mostly at 2900-3300 m. In 1979, five specimens (LSUMZ) were collected in depto. La Libertad at Mashua, east of Tayabamba, on the trail to Ongón, 3350 m, 13-24 September. In 1981, three (LSUMZ) were collected in depto. San Martín at Puerta del Monte, ca. 30 km NE Los Alisos, ca. 3250 m, 6-18 August. In 1978, six were collected at four localities in the Cordillera Colán, depto. Amazonas: 2 at 2990 m and 1 at 3050 m, northeast of La Peca, and 2 at 2685 m and 1 at 3050 m, east of La Peca, 30 August-11 September. In addition, new localities have been discovered in depto. Huánuco since Blake and Hocking's (1974) type description, all above Acomayo (all LSUMZ): (a) 1 at Bosque Cahuencho, 10 April 1974; (b) 2 at Quillacocha, ca. 3800 m, 18 June 1974; (c) 1 at Punta de Esperanza, on trail to Huaylaspampa, 6 February 1975; (d) 4 at Unchog, pass between Churubamba and Hacienda Paty, 3050-3350 m, 14 June and 17 July 1975, and 6 August 1977; and (e) 1 at Bosque Ishanca Ragra, in (?June) 1975.

This species may not belong in *Hemispingus*. Its skulking, secretive, slow-moving behavior (Isler and Isler 1987) seems unusual for the genus. The little-known *H. goeringi* of the Venezuelan Andes, proposed as the closest relative of *rufosuperciliaris* by Blake and Hocking, resembles *H. rufosuperciliaris* in color pattern, elevational distribution, foraging stratum and microhabitat (near ground in dense undergrowth), habitat (timberline humid forest), and rarity (information from Isler and Isler 1987, Ridgely and Tudor 1989).

Wetmorethraupis sterrhopteron Lowery and O'Neill, 1964, Auk 81: 126.

HOLOTYPE: LSUMZ 31457, "Chávez Valdivia, near confluence of the Río Comaina and the Río Cenepa, 4°26'S, 78°11'W, Depto. Amazonas, Peru." Unsexed skin collected by Aguaruna Indians in July 1963. Obtained indirectly by Millie Larson and John P. O'Neill.

COMMENTS: A new genus (Auk 81: 125) and species. The original description was based on the unique holotype. Subsequently, 53 specimens (LSUMZ) were collected, primarily by native hunters, in the upper Marañón Valley at approximately 600-800 m, all in depto. Amazonas and mostly from April to August 1964, with three in July 1978. These localities are: Kusú, Río Comaina (n = 4), Quebrada Achunts, near Kusú on Río Comaina (2), Chiangkus (2), Chicais (4), Chipi (1), Tutinum, Río Cenepa (6), Suwa, Río Cenepa (1), vicinity of Pagat on Río Cenepa (1), Río Cenepa (1), 2 mi. west of Urakusa, Río Marañón (2), Nazaret, Río Marañón (2), Bashuim, Río Kagka (2), and uncertain locality within same region (25).

Isler and Isler (1987) and Ridgely and Tudor (1989) suggested that *Wetmorethraupis* was most closely related to the *Bangsia* tanagers of the Chocó region and southern Central America based on plumage pattern and behavior. We add to this that they also share remarkably restricted geographical and elevational distributions.

Euphonia affinis olmecorum Dickerman, 1981, Occas. Pap. Mus. Zool. Louisiana State Univ. (59): 4.

HOLOTYPE: LSUMZ 16068, "1.3 miles beyond Ajinche on road to Oviedo in the vicinity of Ebano, San Luis Potosí." Adult female collected by Robert J. Newman on 15 March 1950, prepared by Marcella Newman. Original number RJN 1636.

PARATYPES (7): LSUMZ 12655, 2.7 mi. E of Tamuin, collected by RJN on 5 June 1948; LSUMZ 12658, Ebano, collected by RJN on 7 November 1946; LSUMZ 16064, Hacienda Capulin, collected by RJN on 16 June 1950; LSUMZ 17558, El Nacimiento del Coy, 300 ft., collected by R. Graber on 28 April 1951; LSUMZ 17559, Puente de Dios, 300 ft., collected by J. Graber on 31 May 1951; LSUMZ 18727, El Sol [Tamazunchale], collected by L. Guerrero on 20 July 1952 (all preceding from San Luis Potosí); LSUMZ

24904, 1 mi. SW (not SE) Valle Nacional, Oaxaca, collected by L. C. Binford on 27 March 1961 (all females).

Tangara meyerdeschauenseei Schulenberg and Binford, 1985, Wilson Bull. 97: 413.

HOLOTYPE: LSUMZ 98917, "2 km northeast of Sandia, ca. 2175 m, 14°17'S, 69°26'W, Department of Puno, Peru." Adult male collected by Laurence C. Binford on 2 November 1980. Original number LCB 1495.

PARATYPE: LSUMZ 98918, female, other data same as holotype.

COMMENTS: This species is still known only from four specimens from two localities in depto. Puno, from sight records at a third locality in depto. Puno (Schulenberg and Binford 1985), and from a possible sighting at Calabatea in adjacent depto. La Paz, Bolivia (Parker and Bailey 1991; initially thought to be *meyerdeschauenseei* and published as "*meyerdeschauenseei* ?," but Parker (pers. comm.) later decided that *T. cayana* could not be ruled-out).

Nephelornis oneilli Lowery and Tallman, 1976, Auk 93: 420.

HOLOTYPE: LSUMZ 81114, "Bosque Unchog, on pass between Churubamba and Hacienda Paty above Acomayo, 09°41'S, 76°07'W, elevation approximately 3592 meters, Depto. Huánuco, Perú." Adult male collected by Theodore Parker III on 16 July 1975. Original number TP 922.

PARATYPES (38): LSUMZ 74730-31 (skins and partial skeletons), Bosque Cahuincho above Acomayo, 11,200 ft., males collected by Manuel Villar and R. S. Louque, respectively, on 15 June 1973; LSUMZ 80112-80118, Bosque Quiullacocha above (NE) Acomayo, ca. 12,500 ft., five males and two females collected by MV on 14-18 June 1974; LSUMZ 80119-21, Bosque Quiullacocha, ca. 9800 ft., female? and two females collected by TP on 27-28 January 1975; LSUMZ 81091-93, 81095, 81097-104, 81106-07, 81110-12, 81115-81119, 81120 (skeleton), 81121-22 (both fluid-preserved), 81358 (skeleton), type locality, ca. 10,000-11,500 ft., 13 males, 11 females, two unsexed, collected by R. S. Kennedy, TP, and MV on 12 June-17 July 1975.

COMMENTS: A new genus (Auk 93: 416) and species. The elevation given on the holotype label is "ca. 11,500 ft.," which is actually ca. 3500 m (not "ca. 3592 m"). An additional eight paratypes (LSUMZ 77654, 80122, 81094, 81096, 81105, 81108-09, 81113) have been exchanged.

Subsequent LSUMNS fieldwork has extended its range north to dptos. La Libertad and San Martín, about 400 km north of the type locality, as well as south to depto. Pasco, about 75 km southwest of the type locality. In 1979, 20 specimens were collected in depto. La Libertad at Mashua, east of Tayabamba, on trail to Ongón, 3350 m, 12 September - 1 October. In 1981, seven were collected in depto. San Martín at Puerta del Monte, ca. 30 km NE Los Alisos, ca. 3250 m, 11-22 August. In 1983, three were collected at Mascarón, 3710 m, uppermost Bosque San Marcos, ca. 18 km west of

Panao, depto. Huánuco, 28 June 1983; these were the first records south of the Río Huallaga canyon. In 1983 and 1984, 17 additional specimens were collected at the type locality. In 1985, 42 specimens were collected in depto. Pasco at Millpo, east of Tambo de Vacas on Pozuzo-Chaglla trail, 25 at 3450 m and 17 at 3600 m, 16 July-8 August. Three other specimens were collected in depto. Pasco at 1 km east of Huánuco border on Pozuzo-Panao trail, 3700 m, 25 July 1985. Isler and Isler (1987) summarized the information obtained from this recent fieldwork on the natural history of this unusual species, now considered to be a tanager.

FAMILY ICTERIDAE

Cacicus koepckeae Lowery and O'Neill, 1965, Occas. Pap. Mus. Zool. Louisiana State Univ. (33): 1.

HOLOTYPE: LSUMZ 42827, "Balta (at the point where the streams known to local Cashinahua Indians as the Xumuya and the Inuya enter the Río Curanja), lat. 10° 08' S, long. 71° 13' W, elevation approximately 300 meters, Depto. Loreto, Perú." Adult male collected by J. P. O'Neill on 22 March 1965. Original number JPO 1487.

PARATYPE: LSUMZ 31429, Río Curanja, male collected by JPO on 23 July 1963.

COMMENTS: This species is still known only from the holotype and paratype; a sight record from southeastern Peru (in Terborgh et al. 1974) is considered not well-documented (D. F. Stotz, pers. comm.).

Lowery and O'Neill (1965) considered *Cacicus chrysopterus* to be the closest relative of *C. koepckeae*. Subsequent authors have agreed with Lowery and O'Neill either explicitly (Mayr 1971) or implicitly by placing *koepckeae* next to *chrysopterus* in their linear sequence (Blake 1968, Ridgely and Tudor 1989). Sibley and Monroe (1990) formally designated *koepckeae* as an allospecies of *chrysopterus*. As pointed out to us about ten years ago by T. A. Parker, the closest relative of *koepckeae* is most likely *C. sclateri*. These two species share the following: (a) virtually identical body size, more similar to each other than either is to *chrysopterus*; (b) similar distribution, namely lowlands of western Amazonia; and (c) rarity, both known from just a few specimens and localities (in contrast to *C. chrysopterus*, which is widely distributed and locally common; Ridgely and Tudor 1989). The plumage of both *sclateri* and *koepckeae* is also a slightly deeper black than that of *chrysopterus*. Lowery and O'Neill (1965) noted that *C. koepckeae* differs from *C. chrysopterus* in having a more graduated tail, relatively larger feet, and a slightly swollen, broader-based culmen. They did not note, however, that *C. koepckeae* and *C. sclateri* are identical in these features. In fact, *C. koepckeae* and *C. sclateri* differ only in a single plumage feature, namely the presence of a yellow rump patch in *koepckeae*.

Agelaius phoeniceus nelsoni Dickerman, 1965, Occas. Pap. Mus. Zool. Louisiana State Univ. (31): 2.

HOLOTYPE: LSUMZ 42674 [skin and partial skeleton], "Lago Coatetelco, Morelos, México." Adult male collected by Robert W. Dickerman on 18 November 1964. Original number RWD 12176.

PARATYPES (11): LSUMZ 14648, 42027, Acatlipa km 86, 4000 ft., Morelos, males collected by R. J. Newman and J. S. Rowley on 1 June 1949 and 14 May 1959, respectively; LSUMZ 48691, type locality, female collected by RWD on 30 November 1963; LSUMZ 42022-23, 5 mi. N Izucar de Matamoros, 4350 ft., Puebla, males collected by Chester C. Lamb on 18 and 16 March 1956; LSUMZ 42024-26, 48010-12, 4 mi. N Izucar de Matamoros, 4350 ft., four males and two females collected by CCL on 27-29 July 1957.

COMMENTS: Blake (1968) recognized *nelsoni*.

Agelaius phoeniceus brevirostris Monroe, 1963, Occas. Pap. Mus. Zool. Louisiana State Univ. (26): 6.

HOLOTYPE: LSUMZ 30249, "4 miles north of Río Lindo, Department of Cortés, Honduras." Adult male collected by Burt L. Monroe on 29 November 1962. Original number BLM 3393.

PARATYPES (3): LSUMZ 30250, female, other data same as holotype; LSUMZ 30251, Jaral, 2100 ft., Dept. of Cortés, female collected by BLM on 12 January 1963; LSUMZ 30252, 1 mi. W Jaral, 2200 ft., male collected by BLM on 21 August 1962.

COMMENTS: From the holotype label, the elevation of the type locality is 400 ft. Blake (1968) recognized *brevirostris*.

Cassidix mexicanus prosopidicola Lowery, 1938, Occas. Pap. Mus. Zool. Louisiana State Univ. (1): 1.

HOLOTYPE: LSUMZ 1568, "Brownsville, Cameron County, Texas." Adult female collected by George H. Lowery, Jr. on 25 October 1937. Original number GHL 619.

PARATYPES (5): LSUMZ 1578, 1632, males, other data same as holotype; LSUMZ 1637, type locality, male collected by GHL on 24 October 1937; LSUMZ 1565, 1633, Tivoli, Refugio Co., males collected by GHL on 27 October 1937.

COMMENTS: The complete locality on the holotype label reads "Texas: Cameron County, 6 mi. W Brownsville." Placed in *Quiscalus* by most recent authors. Blake (1968) recognized *prosopidicola*.

Molothrus ater buphilus Oberholser, 1938, The bird life of Louisiana. Louisiana Dept. Conservation Bull. 28: 611.

HOLOTYPE: LSUMZ 14745 ("old number 1-31633"), "Marsh Island, Louisiana." Adult male collected by John D. Mizelle on 16 March 1933.

PARATYPES (4): LSUMZ 1230, Baton Rouge, East Baton Rouge Par., male collected by G. H. Lowery on 4 March 1937; LSUMZ 1650, Shreveport, Caddo Par., male collected by GHL on 16 May 1937; LSUMZ 1655, Grand Isle, Jefferson Par., male collected by Si Ward on 2 August 1937; LSUMZ 1669, Montgomery, Grant Par., male collected by GHL on 10 June 1937.

COMMENTS: The type locality is in Iberia Parish. Blake (1968) synonymized *buphilus* with *M. a. ater*, but no reasons were given.

ACKNOWLEDGMENTS

We thank T. S. Schulenberg and R. W. Storer for helpful comments on the manuscript.

LITERATURE CITED

- ALLEN, L. 1977. New species of curassow discovered. *Game Bird Breeders Avicult. Zoo. Conserv. Gaz.* 26: 6-7.
- AMERICAN ORNITHOLOGISTS' UNION. 1983. Check-list of North American Birds. Sixth edition. American Ornithologists' Union, Lawrence, KS.
- BINFORD, L. C. 1965. Two new subspecies of birds from Oaxaca, Mexico. *Occas. Pap. Mus. Zool. Louisiana State Univ.* (30): 1-6.
- BINFORD, L. C. 1989. A distributional survey of the birds of the Mexican state of Oaxaca. *Ornithological Monogr.* (43): 1-418.
- BLAKE, E. R. 1968. Family Icteridae. Pp. 138-202 *in*: Check-list of Birds of the World, Vol. XIV (Paynter, R. A., Jr., ed.). Museum of Comparative Zoology, Cambridge, MA.
- BLAKE, E. R. 1977. *Manual of Neotropical Birds*, Vol. 1. Univ. Chicago Press, Chicago, IL.
- BLAKE, E. R., AND P. HOCKING. 1974. Two new species of tanager from Peru. *Wilson Bull.* 86: 321-324.
- BROWN, L. AND D. AMADON. 1968. *Eagles, Hawks and Falcons of the World*, Vol. 2. McGraw-Hill, New York, NY.
- BROWNING, M. R. 1989. The type specimens of Hekstra's owls. *Proc. Biol. Soc. Washington* 102: 515-519.

- BUDEN, D. W. 1985. New subspecies of Thick-billed Vireo (Aves: Vireonidae) from the Caicos Islands, with remarks on the taxonomic status of other populations. *Proc. Biol. Soc. Washington* 98: 591-597.
- BUDEN, D. W. 1993. Geographic variation in the Scaly-breasted Thrasher *Margarops fuscus* with descriptions of three new subspecies. *Bull. Brit. Orn. Club* 113: 75-84.
- BURLEIGH, T. D., AND G. H. LOWERY, JR. 1939. Description of two new birds from western Texas. *Occas. Pap. Mus. Zool. Louisiana State Univ.* (6): 67-68.
- BURLEIGH, T. D., AND G. H. LOWERY, JR. 1942. An inland race of *Sterna albifrons*. *Occas. Pap. Mus. Zool. Louisiana State Univ.* (10): 173-177.
- BURLEIGH, T. D., AND G. H. LOWERY, JR. 1944. Geographical variation in the Red-bellied Woodpecker in the southeastern United States. *Occas. Pap. Mus. Zool. Louisiana State Univ.* (17): 293-301.
- COFFEY, B. B., JR., AND L. C. COFFEY. 1989. Songs of Mexican Birds, Cassette No. 2. ARA Records, Gainesville, FL.
- COLLAR, N. J., L. P. GONZAGA, N. KRABBE, A. MADROÑO NIETO, L. G. NARANJO, T. A. PARKER, III, AND D. G. WEGE. 1992. Threatened Birds of the Americas. The ICBP/IUCN Red Data Book. Third edition, part 2. International Council Bird Preservation, Cambridge, UK.
- DAVIS, T. J., AND J. P. O'NEILL. 1986. A new species of antwren (Formicariidae: *Herpsilochmus*) from Peru, with comments on the systematics of other species of the genus. *Wilson Bull.* 98: 337-352.
- DICKERMAN, R. W. 1965. The nomenclature of the Red-winged Blackbird (*Agelaius phoeniceus*) of south-central Mexico. *Occas. Pap. Mus. Zool. Louisiana State Univ.* (31): 1-6.
- DICKERMAN, R. W. 1973. A review of the White-breasted Woodwrens of México and Central America. *Condor* 75: 361-363.
- DICKERMAN, R. W. 1975. Revision of the Short-billed Marsh Wren (*Cistothorus platensis*) of Mexico and Central America. *Amer. Mus. Novitates* (2569): 1-8.
- DICKERMAN, R. W. 1981. Geographic variation in the Scrub Euphonia. *Occas. Pap. Mus. Zool. Louisiana State Univ.* (59): 4-5.

- FJELDSÅ, J. AND N. KRABBE. 1990. Birds of the High Andes. Zoological Museum, Univ. of Copenhagen, Denmark.
- FORSHAW, J. M. 1989. Parrots of the World. Third {revised} edition. Weldon Publishing, Willoughby, Australia.
- FRIEDMANN, H. 1950. The Birds of North and Middle America. Part XI. U. S. Natl. Mus. Bull. 50: 1-793.
- FRIEDMANN, H., L. GRISCOM, AND R. T. MOORE. 1950. Distributional check-list of the birds of Mexico, Part I. Pacific Coast Avifauna 29: 1-202.
- GANIER, A. F. 1954. A new race of the Yellow-bellied Sapsucker. Migrant 25: 37-41.
- GRAVES, G. R. 1980. A new species of metaltail hummingbird from northern Peru. Wilson Bull. 92: 1-7.
- GRAVES, G. R. 1981. A new subspecies of Coppery Metaltail (*Metallura theresiae*) from northern Peru. Auk 98: 382.
- GRAVES, G. R. 1987. A cryptic new species of antpitta (Formicariidae: *Grallaria*) from the Peruvian Andes. Wilson Bull. 99: 313-321.
- GRAVES, G. R., J. P. O'NEILL, AND T. A. PARKER, III. 1983. *Grallaricula ochraceifrons*, a new species of antpitta from northern Peru. Wilson Bull. 95: 1-6.
- HEKSTRA, G. P. 1982. Descriptions of twenty-four new subspecies of American *Otus* (Aves, Strigidae). Bull. Zool. Mus. Univ. Amsterdam 9: 49-63.
- HELLMAYR, C. E., AND B. CONOVER. 1949. Catalogue of Birds of the Americas, Part I, Number 4. Zoological Series, Field Museum of Natural History, Chicago, IL.
- LANYON, W. E. 1988. A phylogeny of the thirty-two genera in the *Elaenia* assemblage of tyrant flycatchers. Amer. Mus. Novitates No. (2914): 1-57.
- LOWERY, G. H. 1938. A new grackle of the *Cassidix mexicanus* group. Occas. Pap. Mus. Zool. Louisiana State Univ. (1): 1-11.
- LOWERY, G. H., JR., AND D. G. BERRETT. 1963. A new Carolina Wren (Aves: Troglodytidae) from southern Mexico. Occas. Pap. Mus. Zool. Louisiana State Univ. (24): 1-3.

- LOWERY, G. H., JR., AND R. J. NEWMAN. 1949. New birds from the state of San Luis Potosi and the Tuxtla Mountains of Veracruz, Mexico. Occas. Pap. Mus. Zool. Louisiana State Univ. (22): 1-10.
- LOWERY, G. H., JR., AND J. P. O'NEILL. 1964. A new genus and species of tanager from Peru. Auk 81: 125-131.
- LOWERY, G. H., JR., AND J. P. O'NEILL. 1965. A new species of *Cacicus* (Aves: Icteridae) from Peru. Occas. Pap. Mus. Zool. Louisiana State Univ. (33): 1-5.
- LOWERY, G. H., JR., AND J. P. O'NEILL. 1966. A new genus and species of cotinga from eastern Peru. Auk 83: 1-9.
- LOWERY, G. H., JR., AND J. P. O'NEILL. 1969. A new species of antpitta from Peru and a revision of the subfamily Grallariinae. Auk 86: 1-12.
- LOWERY, G. H., JR., AND D. A. TALLMAN. 1976. A new genus and species of nine-primaried oscine of uncertain affinities from Peru. Auk 93: 415-428.
- MILLER, A. H. 1943. A new race of Brown-headed Chickadee from northern Washington. Occas. Pap. Mus. Zool. Louisiana State Univ. (14): 261-263.
- MONROE, B. L., JR. 1963. Three new subspecies of birds from Honduras. Occas. Pap. Mus. Zool. Louisiana State Univ. (26): 1-7.
- MONROE, B. L., JR., AND T. R. HOWELL. 1966. Geographic variation in Middle American parrots of the *Amazona ochrocephala* complex. Occas. Pap. Mus. Zool. Louisiana State Univ. (34): 1-18.
- NAUMBERG, E. M. B. 1935. Gazetteer and maps showing stations visited by Emil Kaempfer in eastern Brazil and Paraguay. Bull. Amer. Mus. Nat. Hist. 68: 449-470.
- NAUMBERG, E. M. B. 1939. Studies of birds from eastern Brazil and Paraguay, based on a collection made by Emil Kaempfer. Bull. Amer. Mus. Nat. Hist. 76: 231-276.
- OBERHOLSER, H. C. 1938. The bird life of Louisiana. Louisiana Dept. Conserv. Bull. 28: 611.
- O'NEILL, J. P., AND G. R. GRAVES. 1977. A new genus and species of owl (Aves: Strigidae) from Peru. Auk 94: 409-416.

- O'NEILL, J. P., AND T. A. PARKER, III. 1976. New subspecies of *Schizoeaca fuliginosa* and *Uromyias agraphia* from Peru. Bull. Brit. Orn. Club 96: 136-141.
- O'NEILL, J. P., AND T. A. PARKER, III. 1981. New subspecies of *Pipreola riefferii* and *Chlorospingus ophthalmicus* from Peru. Bull. Brit. Orn. Club 101: 294-299.
- PARKER, T. A., III, AND J. P. O'NEILL. 1985. A new species and a new subspecies of *Thryothorus* wren from Peru. Neotropical Ornithology, Ornithol. Monogr. (36): 9-15.
- PARKER, T. A., III, AND B. BAILEY (EDS.). 1991. A Biological Assessment of the Alto Madidi Region and Adjacent Areas of Northwest Bolivia, May 18-June 15, 1990. Rapid Assessment Program Working Papers No. 1. Conservation International, Washington, DC.
- PAYNTER, R. A., JR. 1960. Family Troglodytidae. Pp. 379-440 in: Check-list of Birds of the World, Vol. IX (Mayr, E. and J. C. Greenway, eds.). Museum of Comparative Zoology, Cambridge, MA.
- PAYNTER, R. A., JR. 1970. Subfamily Emberizinae. Pp. 3-214 in: Check-list of Birds of the World, Vol. XIII (Paynter, R. A., Jr., ed.). Museum of Comparative Zoology, Cambridge, MA.
- PHILLIPS, A. R. 1986. The Known Birds of North and Middle America, Part I. Published by the author, Denver, CO.
- PHILLIPS, A. R. 1991. The Known Birds of North and Middle America, Part II. Published by the author, Denver, CO.
- PRATT, H. D. 1979. A new subspecies of the Elepaio, *Chasiempis sandwichensis*, from the Island of Hawaii. Bull. Brit. Orn. Club 99: 105-108.
- REMSEN, J. V., JR. 1981. A new subspecies of *Schizoeaca harterti* with notes on taxonomy and natural history of *Schizoeaca* (Aves: Furnariidae). Proc. Biol. Soc. Washington 94: 1068-1075.
- REMSEN, J. V., JR. AND M. A. TRAYLOR, JR. 1989. Annotated List of the Birds of Bolivia. Buteo Books, Vermilion, SD.
- RIDGELEY, R. S. AND G. TUDOR. 1989. The Birds of South America, Vol I. Univ. of Texas Press, Austin, TX.

- RIPLEY, S. D. 1964. Subfamily Turdinae. Pp. 13-227 *in*: Check-list of birds of the World, Vol. X (Mayr, E. and R. A. Paynter, Jr., eds.). Museum of Comparative Zoology, Cambridge, MA.
- RUSSELL, S. M. 1963. A new race of *Centurus aurifrons* (Aves: Picidae) from British Honduras. Occas. Pap. Mus. Zool. Louisiana State Univ. (25): 1-3.
- SCHULENBERG, T. S., AND L. C. BINFORD. 1985. A new species of tanager (Emberizidae: Thraupinae, *Tangara*) from southern Peru. Wilson Bull. 97: 413-420.
- SCHULENBERG, T. S., AND M. D. WILLIAMS. 1982. A new species of antpitta (*Grallaria*) from northern Peru. Wilson Bull. 94: 105-113.
- SCHULENBERG, T. S., AND G. L. GRAHAM. 1981. A new subspecies of *Anairetes agraphia* (Tyrannidae) from northern Peru. Bull. Brit. Orn. Club 101: 241-243.
- SHORT, L. L. 1982. Woodpeckers of the World. Delaware Museum of Natural History, Greenville, DE.
- SIBLEY, C. G., AND B. L. MONROE, JR. 1990. Distribution and Taxonomy of Birds of the World. Yale Univ. Press, New Haven, CT.
- SNOW, D. W. 1967. Family Paridae. Pp. 70-124 *in*: Check-list of Birds of the World, Vol. XII (Paynter, R. A., Jr., ed.) Museum of Comparative Zoology, Cambridge, MA.
- STORER, R. W. 1970. Subfamily Thraupinae. Pp. 246-408 *in*: Check-list of Birds of the World, Vol. XIII (Paynter, R. A., Jr., ed.). Museum of Comparative Zoology, Cambridge, MA.
- STRESEMANN, E. AND D. AMADON. 1979. Order Falconiformes. Pp. 271-425 *in*: Check-list of Birds of the World, Vol. I (Mayr, E. and G. W. Cottrell, eds.). Museum of Comparative Zoology, Cambridge, MA.
- TERBORGH, J. W., J. W. FITZPATRICK, AND L. EMMONS. 1984. Annotated checklist of bird and mammal species of Cocha Cashu Biological Station, Manu National Park, Peru. Fieldiana (Zool.) 21: 1-29.
- THOMPSON, B. C., M. E. SCHMIDT, S. W. CALHOUN, D. C. MORIZOT, AND R. D. SLACK. 1992. Subspecific status of Least Tern populations in Texas: North American implications. Wilson Bull. 104: 244-262.

- TRAYLOR, M. A., Jr. 1979. Subfamily Elaeniinae. Pp. 1-112 *in*: Check-list of Birds of the World, Vol. VIII (Traylor, M. A., Jr., ed.). Museum of Comparative Zoology, Cambridge, MA.
- VAN ROSSEM, A. J., AND M. HACHISUKA. 1937. A northwestern race of the Mexican Black Hawk. *Trans. San Diego Soc. Nat. Hist.* 8: 361-362.
- VAN ROSSEM, A. J. 1948. A race of *Urubitornis solitaria* from northwestern Mexico. *Proc. Biol. Soc. Washington* 61: 67-68.
- VUILLEUMIER, F., AND E. MAYR. 1987. New species of birds described from 1976 to 1980. *Jour. für Ornith.* 128: 137-150.
- VUILLEUMIER, F., M. LECROY, AND E. MAYR. 1992. New species of birds described from 1981 to 1990. *Bull. Brit. Ornith. Club Centenary Suppl.* 112A: 267-309.
- WATSON, G. E., M. A. TRAYLOR, JR., AND E. MAYR. 1986. Subfamily Monarchinae. Pp. 465-526 *in*: Check-list of Birds of the World, Vol. XI (Mayr, E. and C. W. Cottrell, eds.). Museum of Comparative Zoology, Cambridge, MA.
- WIEDENFELD, D. A. 1982. A nest of the Pale-billed Antpitta (*Grallaria carrikeri*) with comparative remarks on antpitta nests. *Wilson Bull.* 94: 580-582.