1-1-2014

Gastrointestinal helminths in two species of geckos, gekko vittatus, gehyra oceanica, and five species of skinks, caledoniscincus atropunctatus, Emoia cyanogaster, Emoia erronan, Emoia nigra, Emoia sanfordi (Squamata) from the republic of Vanuatu, Oceania

Stephen R. Goldberg
Whittier College

Charles R. Bursey
Penn State Shenango

Alison M. Hamilton
Smithsonian Institution

Christopher C. Austin
Louisiana State University

Follow this and additional works at: https://digitalcommons.lsu.edu/biosci_pubs

Recommended Citation

Goldberg, S., Bursey, C., Hamilton, A., & Austin, C. (2014). Gastrointestinal helminths in two species of geckos, gekko vittatus, gehyra oceanica, and five species of skinks, caledoniscincus atropunctatus, Emoia cyanogaster, Emoia erronan, Emoia nigra, Emoia sanfordi (Squamata) from the republic of Vanuatu, Oceania. *Comparative Parasitology*, 81 (1), 67-74. <https://doi.org/10.1654/4658.1>

This Article is brought to you for free and open access by the Department of Biological Sciences at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Gastrointestinal Helminths in Two Species of Geckos, *Gekko vittatus*, *Gehyra oceanica*, and Five Species of Skinks, *Caledoniscincus atropunctatus*, *Emoia cyanogaster*, *Emoia erronan*, *Emoia nigra*, *Emoia sanfordi* (Squamata) from the Republic of Vanuatu, Oceania

Authors: Goldberg, Stephen R., Bursey, Charles R., Hamilton, Alison M., and Austin, Christopher C.

Source: Comparative Parasitology, 81(1) : 67-74

Published By: The Helminthological Society of Washington

URL: <https://doi.org/10.1654/4658.1>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Gastrointestinal Helminths in Two Species of Geckos, *Gekko vittatus*, *Gehyra oceanica*, and Five Species of Skinks, *Caledoniscincus atropunctatus*, *Emoia cyanogaster*, *Emoia erronan*, *Emoia nigra*, *Emoia sanfordi* (Squamata) from the Republic of Vanuatu, Oceania

STEPHEN R. GOLDBERG,^{1,5} CHARLES R. BURSEY,² ALISON M. HAMILTON,³ AND CHRISTOPHER C. AUSTIN⁴

¹Department of Biology, Whittier College, P.O. Box 634, Whittier, California 90608, U.S.A.

(e-mail: sgoldberg@whittier.edu),

²Department of Biology, Pennsylvania State University, Shenango Campus, Sharon, Pennsylvania 16146, U.S.A.

(e-mail: cxb13@psu.edu),

³Division of Amphibians and Reptiles, Smithsonian Institution, P.O. Box 37012, MRC 162, Washington, DC 20013, U.S.A.

(e-mail: HamiltonAH@si.edu), and

⁴Department of Biological Sciences, 119 Foster Hall, Louisiana State University, Baton Rouge, Louisiana 70803, U.S.A.

(e-mail: ccaustin@lsu.edu)

ABSTRACT: Two species of geckos, *Gekko vittatus*, *Gehyra oceanica* and 5 species of skinks, *Caledoniscincus atropunctatus*, *Emoia cyanogaster*, *Emoia erronan*, *Emoia nigra*, and *Emoia sanfordi* from the Republic of Vanuatu were examined for helminths. One species of Digenea (*Mesocoelium microom*), 3 species of Cestoda (*Gekkotaenia novaeguineensis*, *Cylindrotaenia hickmani*, *Ophiotaenia greeri*), and 7 species of Nematoda, gravid individuals of *Hedruris hanleyae*, *Maxvachonia chabaudi*, *Parapharyngodon maplestoni*, *Spauligodon gehyrai*, *Spauligodon zweifeli*, juveniles assigned to the Filarioidea, and larvae of *Physocephalus* sp. in cysts were found. Eighteen new host records and 7 new locality records are reported. The lizards of Vanuatu are infected by generalist helminths that occur in a variety of lizards throughout Oceania.

KEY WORDS: Cestoda, Digenea, Nematoda, Gekkonidae, Scincidae, Oceania.

The Republic of Vanuatu is located in the South Pacific Ocean (17°45'S; 168°18'E), situated some 1,750 km east of northern Australia and 500 km northeast of New Caledonia (Fig. 1). This oceanic archipelago is comprised of 83 islands with a total land mass of approximately 18,000 km² (Hamilton et al., 2009; Goldberg et al., 2011). The majority of the islands of Vanuatu are small: only 2 islands in this archipelago are greater than 1,000 km² and only a single island has an elevation greater than 1,000 m (Hamilton et al., 2009).

Prior research has examined the helminth fauna of 2 species of gecko (*Nactus multicarinatus* and *N. pelagicus*) and 5 species of skink (*Emoia caeruleocauda*, *Emoia cyanogaster*, *Emoia impar*, *Emoia nigromarginata*, and *Emoia sanfordi*) from Vanuatu (Goldberg et al., 2005, 2011; Bursey et al., 2010). These studies have generated a faunal list for Vanuatu consisting of 7 species of helminths (Table 1). Here we present data on the endoparasites from an additional 2 species of geckos (*Gekko vittatus* and *Gehyra oceanica*) and 3 species of skinks (*Caledoniscincus atropunctatus*, *Emoia erronan*, and *Emoia*

nigra). We also add to the previously reported endoparasite records from 2 other species of skink from Vanuatu: *E. cyanogaster* and *E. sanfordi*.

Both gecko species (family Gekkonidae) have relatively broad distributions in the Pacific region. *Gekko vittatus* occurs in Indonesia, Palau, New Guinea, the Solomon Islands, and Vanuatu (McCoy, 2006). Within Vanuatu, *G. vittatus* has a relatively limited distribution, occurring only in the Banks and Torres island groups on northern Vanuatu (Hamilton, personal observations). *Gehyra oceanica* is widespread in the islands of the Pacific basin and occurs from the Caroline Islands, the Mariana Islands, Papua New Guinea, the Solomon Islands, and Vanuatu eastward to the Marquesas Islands, the Society Islands, and the Tuamotu Archipelago (Fisher, 1997). Two of the skink species examined in this study also have broad distributions. *Emoia cyanogaster* occurs throughout the Vanuatu archipelago (Hamilton, personal observations) as well as the Solomon Islands and the islands of the Bismarck Archipelago of Papua New Guinea (McCoy, 2006). *Emoia nigra* ranges from the Bismarck Archipelago through the Solomon Islands and Vanuatu eastward to Samoa and Tonga (Brown, 1991). Like *Gekko vittatus*, *E. nigra* has a limited distribution within

⁵Corresponding author.

Figure 1. Map of Vanuatu Archipelago. Islands from which specific lizards were collected are indicated in the text.

Table 1. Previous reports of helminths from lizards of Vanuatu (Goldberg et al., 2005; Goldberg, Bursey, Hamilton et al., 2011; Bursey et al., 2010). Number of helminths, prevalence (%), mean intensity $\pm$ 1 SD, and range for 7 helminth species in 2 gekkonid species (*Nactus multicarinatus* and *Nactus pelagicus*) and 5 scincid species (*Emoia caeruleocauda*, *Emoia cyanogaster*, *Emoia impar*, *Emoia nigromarginata*, and *Emoia sanfordi*). Values calculated from the original data sets.

Digenea	Gekkonidae					
	<i>Nactus multicarinatus</i>		<i>Nactus pelagicus</i>		<i>Emoia caeruleocauda</i>	
	No. and prevalence (%)	Mean intensity $\pm$ SD (range)	No. and prevalence (%)	Mean intensity $\pm$ SD (range)	No. and prevalence (%)	Mean intensity $\pm$ SD (range)
<i>Mesocoelium microon</i>	2; 2/265 (1)	1 $\pm$ 0 (0)	—		—	
Cestoda						
<i>Oochoristica javaensis</i>	67; 32/165 (19)	2.1 $\pm$ 1.7 (1–7)	4; 3/172 (2)	1.3 $\pm$ 0.6 (1–2)	—	
Nematoda						
<i>Falcaustra tannaensis</i>	—		19; 7/172 (4)	3.0 $\pm$ 2.8 (1–8)	—	
<i>Hedruris hanleyae</i>	137; 63/165 (38)	2.2 $\pm$ 2.0 (1–11)	91; 18/172 (10)	5.0 $\pm$ 3.2 (1–14)	18; 8/8 (100)	2.3 $\pm$ 1.5 (1–5)
<i>Parapharyngodon maplestoni</i>	460; 120/125 (73)	3.8 $\pm$ 2.9 (1–17)	448; 127/172 (74)	3.5 $\pm$ 2.4 (1–11)	—	
<i>Physocephalus</i> sp. (larvae)	82; 20/165 (12)	4.1 $\pm$ 5.3 (1–24)	223; 7/172 (4)	31.8 $\pm$ 79.4 (1–212)	—	
Filarioidae gen. sp.	3; 2/165 (1)	1.5 $\pm$ 0.7 (1–2)	—		—	

Vanuatu, occurring in the northern islands of the archipelago, as well as Ambae, Maewo, and northern Espiritu Santo (Hamilton, personal observation). The other 3 skinks examined in this study have more restricted distributions. *Caledoniscincus atropunctatus* occurs on New Caledonia and southern Vanuatu (Bauer and Sadlier, 2000; Hamilton et al., 2010). The remaining 2 species are endemic to the Vanuatu archipelago. *Emoia sanfordi* is distributed in northern and central Vanuatu but absent from the islands south of Cheesman's line, a name proposed by Hamilton et al. (2010) to designate faunal and floral discontinuity between northern and southern islands of Vanuatu. *Emoia erronan* is known only from the small islands of Futuna and Aniwa in southern Vanuatu (Brown, 1991; Hamilton, personal observations). In this study we establish an initial helminth list for *C. atropunctatus* and *E. erronan* and add to the previously reported helminth faunas of *E. cyanogaster*, *E. nigra*, *E. sanfordi*, *Gehyra oceanica*, and *Gekko vittatus*.

METHODS

We examined a total of 30 lizards comprising 7 species collected from 7 islands in the Republic of Vanuatu, Oceania during 2005 and deposited them in the herpetology collection of Louisiana State University, Baton Rouge, Louisiana, U.S.A. (Table 2). Lizards were collected by hand, preserved in 10% neutral buffered formalin, and stored in 70% ethanol. Visceral organs were removed and sent to Whittier College for helminthological examination. The esophagus, stomach, and large and small intestines were examined under a dissecting microscope. Digeneans and

cestodes were regressively stained in hematoxylin, mounted in Canada balsam, and examined with a compound microscope. Nematodes were cleared in glycerol for examination. Parasite terminology is in accordance with Bush et al. (1997). Voucher helminths were deposited in the United States National Parasite Collection, Beltsville, Maryland, U.S.A. (Table 3).

RESULTS

A total of 369 individual endoparasites was collected from the 18 skinks and 12 geckos. A total of 11 species of endoparasites was detected in this sample. Found were 1 species of Digenea (*Mesocotyle microon* [Nicoll, 1914]); 3 species of Cestoda (*Cylindrotaenia hickmani* Jones, 1985; *Gekkotaenia novaeguineensis* Bursey, Goldberg, and Kraus 2005; *Ophiotaenia greeri* [Bursey et al., 2006]); and 6 species of Nematoda (*Hedruris hanleyae* Bursey and Goldberg, 2000; *Maxvachonia chabaudi* Mawson, 1972; *Parapharyngodon maplestoni* Chatterji, 1933; *Spauligodon gehyrai* Bursey and Goldberg, 1996; *Spauligodon zweifeli* Bursey, Goldberg, and Kraus, 2005; Filarioidea gen. sp. [juveniles]; *Physocephalus* sp. [larvae in cysts]). The 2 lizard species for which we examined the largest number of specimens contained the most diverse parasite communities. *Emoia sanfordi* ($N = 12$) harbored 8 species of helminths; the greatest diversity of endoparasites in our study, including digeneans, cestodes, and nematodes, were found in specimens of this species (Table 4). *Gehyra oceanica* ($N = 9$) also had a relatively diverse assemblage of 6 species

Table 1. Extended.

Gekkonidae		Scincidae					
<i>Emoia cyanogaster</i>		<i>Emoia impar</i>		<i>Emoia nigromarginata</i>		<i>Emoia sanfordi</i>	
No. and prevalence (%)	Mean intensity $\pm$ SD (range)	No. and prevalence (%)	Mean intensity $\pm$ SD (range)	No. and prevalence (%)	Mean intensity $\pm$ SD (range)	No. and prevalence (%)	Mean intensity $\pm$ SD (range)
—	—	—	—	—	—	—	—
—	—	—	—	—	—	—	—
—	—	—	—	—	—	—	—
2; 2/2 (100)	1 $\pm$ 0 (0)	6; 4/4 (100)	1.5 $\pm$ 1.0 (1–3)	27; 10/10 (100)	2.7 $\pm$ 0.9 (1–4)	1; 1/1 (100)	0 (0)
—	—	—	—	—	—	—	—
—	—	—	—	—	—	—	—
—	—	—	—	—	—	—	—

Table 2. Host specimens examined for parasites in this study. Specimens are deposited in the herpetology collection of the Louisiana State University Museum of Natural Science (LSUMZ), Baton Rouge, Louisiana.

Host taxon	N	Locality and voucher number
Gekkonidae		
<i>Gehyra oceanica</i>	9	Aniwa: LSUMZ 91309; Espiritu Santo: LSUMZ 91311, 91315; Malekula: LSUMZ 91298; Mota Lava: LSUMZ 91317, 91318, 91322, 91325; Tanna: LSUMZ 91304
<i>Gekko vittatus</i>	3	Mota Lava: LSUMZ 91222, 91223, 91230
Scincidae		
<i>Caledoniscincus atropunctatus</i>	1	Tanna: LSUMZ 90073
<i>Emoia cyanogaster</i>	1	Malekula: LSUMZ 91529
<i>Emoia erronan</i>	1	Aniwa: LSUMZ 91772
<i>Emoia nigra</i>	1	Mota Lava: LSUMZ 90107, 90109, 90110
<i>Emoia sanfordi</i>	12	Ambrym: LSUMZ 90171; Epi: LSUMZ 89895; Espiritu Santo: LSUMZ 90909; Malekula: LSUMZ 90135, 90136, 90145, 90146; Mota Lava: LSUMZ 89904, 89906, 89913, 89915, 89916

representing digeneans, cestodes, and nematodes (Table 4). Number of helminths, prevalence (%), mean intensity $\pm$ 1 SD, and range for these 11 species of helminths by host species are given in Table 4.

DISCUSSION

Dronen et al. (2012) revised the Mesocoelidae Dollfus, 1929 and recognized 43 valid species. In prior revisions, Freitas (1963) established 19 species, which Nasir and Diaz (1971) reduced to 4. Previous Vanuatu helminthological studies have followed the keys of Nasir and Diaz (1971). Following the key provided by Dronen et al. (2012), we have assigned our specimens to *M. microon* (vitelline fields and ceca moderately long and of similar length, genital pore prebifurcal and median, gonads overlapping

acetabulum), a species originally described from the hylid frog species *Litoria caerulea* and *Litoria gracilentia* (reported as *Hyla caerulea*, *Hyla gracilentia*) and *Tiliqua scincoides* (Scincidae) collected in Queensland, Australia by Nicoll (1914). Given the recent revision of the Mesocoelidae, specimens from the gecko *N. multicaudatus* collected in Vanuatu (Goldberg et al., 2011) and assigned to *Mesocoelium monas* should be re-evaluated. *Gehyra oceanica*, *E. nigra*, and *E. sanfordi* represent new host records for *M. microon*; Vanuatu is a new locality record.

Gekkotaenia novaeguineensis was described from *Cyrtodactylus louisianensis* (Gekkonidae) collected in Papua New Guinea (Bursey et al., 2005b). *Gehyra oceanica* represents a new host record and is only the second host record for *G. novaeguineensis*. Vanuatu is a new locality record.

Table 3. Voucher helminths from lizards from Vanuatu deposited in the U.S. National Parasite Collection, (USNPC), Beltsville, Maryland, U.S.A.

Host taxon	N	Helminth and voucher number
Gekkonidae		
<i>Gehyra oceanica</i>	9	<i>Mesocoelium microon</i> (USNPC 106897); <i>Gekkotaenia novaeguineensis</i> (USNPC 106818); <i>Hedruris hanleyae</i> (USNPC 106819); <i>Parapharyngodon maplestoni</i> (USNPC 106820); <i>Spauligodon gehyrai</i> (USNPC 106821); <i>Spauligodon zweifeli</i> (USNPC 106822)
<i>Gekko vittatus</i>	3	<i>Hedruris hanleyae</i> (USNPC 106823); <i>Maxvachonia chabaudi</i> (USNPC 106824)
Scincidae		
<i>Caledoniscincus atropunctatus</i>	1	<i>Parapharyngodon maplestoni</i> (USNPC 106825)
<i>Emoia cyanogaster</i>	1	<i>Hedruris hanleyae</i> (USNPC 106826)
<i>Emoia erronan</i>	1	<i>Cylindrotaenia hickmani</i> (USNPC 106827); <i>Hedruris hanleyae</i> (USNPC 106828); <i>Parapharyngodon maplestoni</i> (USNPC 106829)
<i>Emoia nigra</i>	1	<i>Mesocoelium microon</i> (USNPC 106898); <i>Ophiotaenia greeri</i> (USNPC 106830, 106831); <i>Hedruris hanleyae</i> (USNPC 106832)
<i>Emoia sanfordi</i>	12	<i>Mesocoelium microon</i> (USNPC 106899); <i>Cylindrotaenia hickmani</i> (USNPC 106833, 106834, 106835); <i>Hedruris hanleyae</i> (USNPC 106836); <i>Maxvachonia chabaudi</i> (USNPC 106837); <i>Parapharyngodon maplestoni</i> (USNPC 106838); <i>Spauligodon zweifeli</i> (USNPC 106839); <i>Physocephalus</i> sp. (USNPC 106840); Filarioidae gen. sp. (USNPC 106841)

Cylindrotaenia hickmani was originally described as *Baerietta hickmani* from Australian populations of 5 species of scincid lizards, *Lampropholis challengeri*, *Lampropholis delicata*, *Lampropholis guichenoti*, *Lampropholis mustelina*, and *Nannoscincus maccayi* (Jones, 1985) and subsequently reassigned to *Cylindrotaenia* by Jones (1987). It has been reported from the skink species *Emoia jakati*, *Emoia longicauda*, *Emoia obscura*, *Emoia pallidiceps*, *Emoia popei*, *Sphenomorphus aignanus*, and *Sphenomorphus lousiadensis* from Papua New Guinea (Goldberg et al., 2008, 2009) as well as *E. caeruleocauda* from the Marianas (Goldberg, Bursey, Hamilton et al., 2011). This cestode was found in two skink species in our sample, and *E. erronan* and *E. sanfordi* represent new host records for *C. hickmani*. Vanuatu is a new locality record.

Ophiotaenia greeri was originally described as *Adenobrechmos greeri* from *Sphenomorphus aignanus* (Scincidae) collected in Papua New Guinea (Bursey et al., 2006), but was reassigned to *Ophiotaenia* by de Chambrier et al. (2009). It has also been reported from the Papua New Guinean populations of 6 additional skink species: *Carlia eothern*, *Carlia luctuosa*, *E. jakati*, *E. obscura*, *E. pallidiceps*, and *S. lousiadensis* (Goldberg et al., 2008, 2009, 2010a). *Ophiotaenia greeri* was present in only 1 species examined and this species, *E. nigra* (Scincidae), represents a new host record. Vanuatu is a new locality record.

Hedruris hanleyae was described from the gecko *Hemidactylus garnotii* collected in the Cook Islands (Bursey and Goldberg, 2000) and is common in lizards (Scincidae and Gekkonidae) from Oceania and Papua New Guinea (Goldberg et al., 2005, 2008). This nematode is now known from 15 species of skinks from 2 genera (*Carlia* and *Emoia*) and 10 species of geckos from 5 genera (*Gehyra*, *Gekko*, *Hemidactylus*, *Lepidodactylus*, and *Nactus*); a complete host list will be found in Goldberg et al. (2010a). *Hedruris hanleyae* has the broadest taxonomic distribution of hosts for the endoparasite species found in our sample, and is present in 6 of the 7 host species examined (Table 4). This nematode was previously reported from Vanuatu in *N. multicarinatus* and *N. pelagicus* (Goldberg et al., 2011). *Emoia erronan* and *G. vittatus* represent a new host record for *H. hanleyae*.

Maxvachonia chabaudi was described from Australian lizards by Mawson (1972) and has been reported from gekkonid hosts from Oceania (*Gehyra mutilata*, *G. oceanica*, *Lepidodactylus lugubris*, and

L. paurolepis) and Papua New Guinea (*Cyrtodactylus sermowaiensis* and *Gekko vittatus*). This nematode was found in 2 species in our sample: *G. vittatus* (Gekkonidae) and *E. sanfordi* (Scincidae). *Gekko vittatus* as a host was previously reported by Goldberg et al. (2010b). *Emoia sanfordi* represents a new host record for *M. chabaudi* and is the first non-gekkonid host species identified. Vanuatu is a new locality record.

Parapharyngodon maplestoni was described from the lizard *Calotes versicolor* (Agamidae) collected in Burma by Chatterji (1933). This endoparasite has previously been reported from hosts representing 4 families of squamate reptiles (Agamidae: *Bronchocela* and *Calotes*; Anguidae: *Pseudopus*; Gekkonidae: *Cyrtodactylus*, *Gekko*, *Hemidactylus*, and *Nactus*; Scincidae: *Emoia*, *Eremiascincus*, and *Sphenomorphus*) from Asia the Australo-Papuan region, and Oceania, as well as the Palearctic (Goldberg et al., 2009). More detailed information regarding previously reported hosts can be found in Bursey et al. (2005a) and Goldberg et al. (2009, 2010a, 2010b, 2011). *Parapharyngodon maplestoni* was previously reported from Vanuatu in the gekkonids *N. multicarinatus* and *N. pelagicus* (Goldberg et al., 2011) but was not previously reported from scincid hosts within Vanuatu. The skinks *Caledoniscincus atropunctatus*, *E. erronan*, and *E. sanfordi* and the gecko *Gehyra oceanica* represent new host records for *P. maplestoni*.

Spauligodon gehyae was originally described from *G. oceanica* collected on Guam and subsequent research indicates that this is a common helminth in *G. oceanica* from widely scattered populations across Oceania (Bursey and Goldberg, 1996). It has also been reported from *Emoia trossula* and *L. lugubris* (Goldberg and Bursey, 2002; Goldberg et al., 2005). *Spauligodon gehyae* was present in 4 of the 9 *G. oceanica* examined and was not present in any other host species examined in this study. Vanuatu is a new locality record.

Spauligodon zweifeli was described from *Lepidodactylus novaeguineae* (Gekkonidae) from Papua New Guinea by Bursey et al. (2005b). Two species of lizard from Vanuatu were infected with *S. zweifeli*. *Gehyra oceanica* is the second gecko host reported. *Emoia sanfordi* (Scincidae) is the third host species (and first non-gekkonid host) reported for *S. zweifeli*. Both *E. sanfordi* and *G. oceanica* represent new host records for *S. zweifeli*. Vanuatu is a new locality record.

Juvenile nematodes assigned to Filarioidea have previously been reported from *N. multicarinatus* from

Table 4. Number of helminths, prevalence (%), mean intensity $\pm$ 1 SD, and range for 11 species of helminths in 2 gekkonid species (*Gehyra oceanica* and *Gekko vittatus*) and 5 scincid species (*Caledoniscincus atropunctatus*, *Emoia cyanogaster*, *Emoia erronan*, *Emoia nigra*, and *Emoia sanfordi*) from Vanuatu, Oceania.

	Gekkonidae					
	<i>Gehyra oceanica</i>		<i>Gekko vittatus</i>		<i>Caledoniscincus atropunctatus</i>	
	No. and prevalence (%)	Mean intensity $\pm$ SD (range)	No. and prevalence (%)	Mean intensity $\pm$ SD (range)	No. and prevalence (%)	Mean intensity $\pm$ SD (range)
Digenea						
<i>Mesocoelium microon</i>	1; 1/9 (11)*	0 (0)	—	—	—	—
Cestoda						
<i>Cylindrotaenia hickmani</i>	—	—	—	—	—	—
<i>Gekkotaenia novaequinaensis</i>	1; 1/9 (11)*	0 (0)	—	—	—	—
<i>Ophiotaenia greeri</i>	—	—	—	—	—	—
Nematoda						
<i>Hedruris hanleyae</i>	50; 3/9 (33)	16.7 $\pm$ 10.0 (9–28)	21; 2/3 (67)*	10.5 $\pm$ 6.4 (6–15)	—	—
<i>Maxvachonia chabaudi</i>	—	—	10; 2/3 (67)	5.0 $\pm$ 1.4 (4–6)	—	—
<i>Parapharyngodon maplestoni</i>	4; 1/9 (11)*	0 (0)	—	—	1; 1/1 (100)*	0 (0)
<i>Spauligodon gehyrae</i>	15; 4/9 (44)	3.8 $\pm$ 2.8 (2–7)	—	—	—	—
<i>Spauligodon zweifeli</i>	5; 1/9 (11)*	0 (0)	—	—	—	—
<i>Physocephalus</i> sp. (larvae)	—	—	—	—	—	—
Filarioidae gen. sp.	—	—	—	—	—	—

* New host records.

Vanuatu (Goldberg, Bursey, Hamilton et al., 2011). The gross morphology of these specimens is different from previously described Australo-Papuan species, and these specimens may represent an undescribed species. *Emoia sanfordi* represents a new host record for juvenile Filarioidea.

Physocephalus sp. (larvae in cysts) have previously been found in both *N. multicarinatus* and *N. pelagicus* from Vanuatu (Goldberg et al., 2011). Adults of *Physocephalus* typically occur in the stomachs of pigs, and dung beetles serve as intermediate hosts (Anderson, 2000). Lizards likely become infected with *Physocephalus* by ingesting infected insects. Since development beyond the larval stage does not occur, lizards are likely transport hosts. *Emoia sanfordi* represents a new host record for larvae of *Physocephalus* sp.

Twenty-seven species of lizards have been recorded for Vanuatu (Hamilton et al. 2009; Goldberg et al., 2011; Ineich, 2008). With the addition of this study, 392 individuals representing nearly half of the recognized species (12 species, or 44%) have now been examined for helminths (although for some species, only a few individuals have been examined). From the data collected to date, it appears that with the exception of *Falcaustra tannaensis*, the lizards of

Vanuatu are infected by generalist helminths (helminths that infect more than a single host species). The lizards of Vanuatu are infected by helminth species previously known from Asia, Australia, and New Guinea as well as species commonly found in lizards from the islands of Oceania. Of the cestodes and nematodes found in Vanuatu lizard hosts in this work and previous studies that are identifiable to species level, 50% have a broad “pan-Oceania” distribution, occurring in Papua New Guinea and ranging north and northeast across many of the islands of the Pacific (*Hedruris hanleyae*, *M. chabaudi*, *Oochoristica javanensis*, *Parapharyngodon maplestoni*, and *S. gehyrae*). We postulate that they reached Vanuatu and islands in other regions of the Pacific Ocean by rafting and their establishment was fortuitous. An additional species (*Cylindrotaenia hickmani*) is also a cestode of the Pacific islands, but its range is much narrower; this species has not been reported from any island groups east of Vanuatu. A significant portion of the helminth fauna known from the lizards of Vanuatu is absent from the isolated island groups of the eastern Pacific. Thirty percent of the cestodes and nematodes found in our study were previously known only from Papua New Guinea (*Gekkotaenia novaequinaensis*, *Ophiotaenia greeri*,

Table 4. Extended.

Gekkonidae		Scincidae					
<i>Emoia cyanogaster</i>		<i>Emoia erronan</i>		<i>Emoia nigra</i>		<i>Emoia sanfordi</i>	
No. and prevalence (%)	Mean intensity $\pm$ SD (range)	No. and prevalence (%)	Mean intensity $\pm$ SD (range)	No. and prevalence (%)	Mean intensity $\pm$ SD (range)	No. and prevalence (%)	Mean intensity $\pm$ SD (range)
—	—	—	—	1; 1/3 (33)*	0; 0	1; 1/12 (8)*	0 (0)
—	—	2; 1/1 (100)*	0 (0)	—	—	8; 6/12 (50)*	1.3 $\pm$ 0.5 (1–2)
—	—	—	—	—	—	—	—
—	—	—	—	2; 2/3 (67)*	1 $\pm$ 0 (0)	—	—
1; 1/1 (100)	0 (0)	8; 1/1 (100)*	0 (0)	78; 3/3 (100)	0 (0)	135; 7/12 (58)	26.0 $\pm$ 28.8 (6–59)
—	—	—	—	—	—	1; 1/12 (8)*	0 (0)
—	—	2; 1/1 (100)*	0 (0)	—	—	3; 2/12 (17)*	1.5 $\pm$ 0.7 (1–2)
—	—	—	—	—	—	—	—
—	—	—	—	—	—	16; 3/12 (25)*	5.3 $\pm$ 4.5 (1–10)
—	—	—	—	—	—	1; 1/12 (8)*	0 (0)
—	—	—	—	—	—	1; 1/12 (8)*	0 (0)

and *S. zweifeli*). To date, only a single species of nematode, *F. tannaensis*, is unique to the lizard fauna of Vanuatu.

Bursey et al. (2001) introduced the concept of importance (*I*), an estimate of the fitness (relative survival and reproductive success) of a species within a community calculated as $I = \text{relative prevalence} + \text{relative abundance} \times 100$. By this measure, the most important parasites to the supracommunity are *P. maplestoni* ($I = 73$) and *H. hanleyae* ($I = 63$); remaining helminth species have an *I* value less than 20.

Although the life cycles of most of the helminth species found in this study have not been examined, life cycles of congeners are available. Species assigned to Pharyngodonidae, i.e., *Parapharyngodon maplestoni*, *S. gehyae*, and *S. zweifeli*, have direct (monoxenous) life cycles and infection most probably occurs when contaminated substrate is ingested. Members of the Cosmoceroidea, i.e., *M. chabaudi*, also have direct life cycles and infect either orally or by skin penetration (Anderson, 2000). The other species reported in this study require intermediate or transport hosts. Given the identity of the two species with high importance values, it would appear that both diet (the intake of infected food items enabling infection by *H.*

hanleyae) and habitat (substrate licking for environmental clues leading to infection by *P. maplestoni*) are important for the establishment of helminth infection in Vanuatu lizards.

ACKNOWLEDGMENTS

We thank Mr. Ernest Bani and Ms. Donna Kalfatak of the Environment Unit of the Republic of Vanuatu for permits to conduct research on the reptiles of Vanuatu and for export permits for tissues and voucher specimens, numerous chiefs and villagers for assistance with fieldwork in Vanuatu, and E. Klein, E. Hartfield, K. Blaha, M. Eckstut, and K. Grazyck for assistance for fieldwork in Vanuatu. Funding for this project was provided to A.M.H. and C.C.A. by the National Science Foundation (DEB 0408010, DEB 0445213, DEB 1146033, and DBI 04009797), Graduate Women in Science, the American Society of Ichthyologists and Herpetologists, and the Society for the Study of Amphibians and Reptiles. Research was conducted under Louisiana State University Institutional Animal Care and Use Committee Protocol #03-121. We also thank Peggy Firth for the map of Vanuatu (Fig. 1) and Sarah Goldsberry and Tenzing Doleck (Whittier College) for assistance with dissections.

LITERATURE CITED

- Anderson, R. C.** 2000. Nematode Parasites of Vertebrates. Their Development and Transmission, 2nd ed. CABI Publishing, Wallingford, U.K.
- Bauer, A. M., and R. A. Sadlier.** 2000. The Herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles. Ithaca, New York. 310 pp.
- Brown, W. C.** 1991. Lizards of the genus *Emoia* (Scincidae) with observations on their evolution and biogeography. *Memoirs of the California Academy of Science* 15:1–94.
- Bursey, C. R., and S. R. Goldberg.** 1996. *Spauligodon gehyrae* n. sp. (Nematoda: Pharyngodonidae) from *Gehyra oceanica* (Sauria: Gekkonidae) from Guam, Mariana Islands, Micronesia. *Journal of Parasitology* 82:962–964.
- Bursey, C. R., and S. R. Goldberg.** 2000. *Hedruris hanleyae* n. sp. (Nematoda: Hedruidae) from *Hemidactylus garnotii* (Sauria: Gekkonidae) from the Cook Islands. *Journal of Parasitology* 86:556–559.
- Bursey, C. R., S. R. Goldberg, A. M. Hamilton, and C. C. Austin.** 2010. A new species of *Falcaustra* (Nematoda: Kathliniidae) in *Nactus pelagicus* (Squamata: Gekkonidae) from Tanna Island, Vanuatu. *Journal of Parasitology* 96: 968–971.
- Bursey, C. R., S. R. Goldberg, and F. Kraus.** 2005a. New genus, new species of Cestoda (Anoplocephalidae), new species of Nematoda (Cosmocercidae) and other helminths in *Cyrtodactylus louisiadensis* (Sauria: Gekkonidae) from Papua New Guinea. *Journal of Parasitology* 91:882–889.
- Bursey, C. R., S. R. Goldberg, and F. Kraus.** 2005b. New species of *Spauligodon* (Nematoda: Pharyngodonidae) in *Lepidodactylus novaeguineae* (Sauria: Gekkonidae) from Papua New Guinea. *Journal of Parasitology* 91:324–328.
- Bursey, C. R., S. R. Goldberg, and F. Kraus.** 2006. New genus, new species of Cestoda (Proteocephalidae) from the lizard *Sphenomorphus aignanus* (Squamata: Scincidae) from Papua New Guinea. *Comparative Parasitology* 73:184–187.
- Bursey, C. R., S. R. Goldberg, and J. R. Parmelee.** 2001. Gastrointestinal helminths of 51 species of anurans from Reserva Cuzco Amazónico, Peru. *Comparative Parasitology* 68:21–35.
- Bush, A. O., K. D. Lafferty, J. M. Lotz, and A. W. Shostak.** 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology* 83:575–583.
- Chatterji, R. C.** 1933. On a new nematode, *Parapharyngodon maplestoni* gen. nov., sp. nov., from a Burmese lizard. *Annals of Tropical Medicine and Parasitology* 27:131–134.
- de Chambrier, A., S. C. Coquille, J. Mariaux, and V. Tkach.** 2009. Redescription of *Testudotaenia testudo* (Magath, 1924) (Eucestoda: Proteocephalidae), a parasite of *Apalone spinifera* (Le Sueur) (Reptilia: Trionychidae) and *Amia calva* L. (Pisces: Amiidae) in North America and erection of the Testudotaeniinae n. subfam. *Systematic Parasitology* 73:49–64.
- Dronen, N. O., D. M. Calhoun, and S. R. Simcik.** 2012. *Mesocoelium* Odhner, 1901 (Digenea: Mesocoeliidae) revisited; a revision of the family and re-evaluation of species composition in the genus. *Zootaxa* 3387:1–96.
- Fisher, R. N.** 1997. Dispersal and evolution of the Pacific basin gekkonid lizards, *Gehyra oceanica* and *Gehyra mutilata*. *Evolution* 51:906–921.
- Freitas, J. F. T.** 1963. Revisão de família Mesocoeliidae Dollfus, 1933 (Trematoda). *Memórias do Instituto Oswaldo Cruz* 61:177–311.
- Goldberg, S. R., and C. R. Bursey.** 2002. Gastrointestinal helminths of seven gekkonid species (Sauria: Gekkonidae) from Oceania. *Journal of Natural History* 36:2249–2264.
- Goldberg, S. R., C. R. Bursey, and R. N. Fisher.** 2005. Helminth records from eleven species of *Emoia* (Sauria: Scincidae) from Oceania. *Pacific Science* 59:609–614.
- Goldberg, S. R., C. R. Bursey, A. M. Hamilton, and C. C. Austin.** 2011. Gastrointestinal helminth communities of two gekkonid lizard species, *Nactus multicarinatus* and *Nactus pelagicus* (Squamata: Gekkonidae), from the Republic of Vanuatu, Oceania. *Journal of Natural History* 45:1983–1993.
- Goldberg, S. R., C. R. Bursey, and F. Kraus.** 2008. Gastrointestinal helminths of eleven species of *Emoia* (Squamata: Scincidae) from Papua New Guinea. *Journal of Natural History* 42:1923–1935.
- Goldberg, S. R., C. R. Bursey, and F. Kraus.** 2009. Endoparasites in 12 species of *Sphenomorphus* (Squamata: Scincidae) from Papua New Guinea. *Comparative Parasitology* 76:58–83.
- Goldberg, S. R., C. R. Bursey, and F. Kraus.** 2010a. Metazoan endoparasites of 14 species of skinks (Squamata: Scincidae) from Papua New Guinea. *Journal of Natural History* 44:447–467.
- Goldberg, S. R., C. R. Bursey, and F. Kraus.** 2010b. Helminths of ten species of geckos (Squamata: Gekkonidae) from Papua New Guinea, with comparisons between immigrant and endemic geckos. *Pacific Science* 64:131–139.
- Goldberg, S. R., C. R. Bursey, and F. Kraus.** 2011. Helminths of *Emoia caeruleocauda* (Squamata: Scincidae) from the Northern Mariana Islands, Micronesia. *Journal of Natural History* 45:497–503.
- Hamilton, A. M., J. H. Hartman, and C. C. Austin.** 2009. Island area and species richness in the southwest Pacific Ocean: Is the lizard fauna of Vanuatu depauperate? *Ecography* 32:247–258.
- Hamilton, A. M., E. R. Klein, and C. C. Austin.** 2010. Biogeographic breaks in Vanuatu, a nascent oceanic archipelago. *Pacific Science* 64:149–159.
- Ineich, I.** 2008. A new arboreal *Lepidodactylus* (Reptilia: Gekkonidae) from Espiritu Santo Island: from egg to holotype. *Zootaxa* 1918:26–38.
- Jones, M. K.** 1985. Morphology of *Baerietta hickmani* n. sp. (Cestoda, Nematotaeniidae) from Australian scincid lizards. *Journal of Parasitology* 74:4–9.
- Jones, M. K.** 1987. A taxonomic revision of the Nematotaeniidae Lühe, 1910 (Cestoda: Cyclophyllidae). *Systematic Parasitology* 10:165–245.
- Mawson, P. M.** 1972. The nematode genus *Maxvachonia* (Oxyurata: Cosmocercidae) in Australian reptiles and frogs. *Transactions of the Royal Society of South Australia* 96:101–108.
- McCoy, M.** 2006. Reptiles of the Solomon Islands. Pensoft, Sofia, Bulgaria. 147 pp.
- Nasir, P., and M. T. Diaz.** 1971. A redescription of *Mesocoelium monas* (Rudolphi, 1819) Freitas, 1958, and specific determination in genus *Mesocoelium* Odhner 1910 (Trematoda, Digenea). *Rivista di Parassitologia* 32:149–158.
- Nicoll, W.** 1914. The trematode parasites of North Queensland. I. *Parasitology* 6:331–350.