

10-1-2013

Two new species of *Eimeria* Schneider, 1875 (Apicomplexa: Eimeriidae) from emerald tree skinks, *Lamprolepis smaragdina* (Lesson) (Sauria: Scincidae) from Papua New Guinea and the Philippines

Chris T. McAllister
Eastern Oklahoma State College

R. Scott Seville
University of Wyoming

Donald W. Duszynski
The University of New Mexico

Sarah E. Bush
The University of Utah

Robert N. Fisher
United States Geological Survey

See next page for additional authors

Follow this and additional works at: https://digitalcommons.lsu.edu/biosci_pubs

Recommended Citation

McAllister, C., Seville, R., Duszynski, D., Bush, S., Fisher, R., & Austin, C. (2013). Two new species of *Eimeria* Schneider, 1875 (Apicomplexa: Eimeriidae) from emerald tree skinks, *Lamprolepis smaragdina* (Lesson) (Sauria: Scincidae) from Papua New Guinea and the Philippines. *Systematic Parasitology*, 86 (2), 165-171. <https://doi.org/10.1007/s11230-013-9442-7>

This Article is brought to you for free and open access by the Department of Biological Sciences at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Authors

Chris T. McAllister, R. Scott Seville, Donald W. Duszynski, Sarah E. Bush, Robert N. Fisher, and Christopher C. Austin

Published in final edited form as:

Syst Parasitol. 2013 October ; 86(2): 165–171. doi:10.1007/s11230-013-9442-7.

Two new species of *Eimeria* (Apicomplexa: Eimeriidae) from emerald tree skinks, *Lamprolepis smaragdina* (Sauria: Scincidae) from Papua New Guinea and the Philippines

Chris T. McAllister

Science and Mathematics Division, Eastern Oklahoma State College, Idabel, Oklahoma 74745, USA cmcallister@se.edu

R. Scott Seville

Department of Zoology and Physiology, University of Wyoming/Casper Center, Casper, Wyoming 82601, USA

Donald W. Duszynski

Department of Biology, University of New Mexico, Albuquerque, New Mexico 87131, USA

Sarah E. Bush

Department of Biology, University of Utah, Salt Lake City, Utah 84112, USA

Robert N. Fisher

U. S. Geological Survey, Western Ecological Research Center, San Diego Field Station, 4165 Spruance Road, Suite 200, San Diego, California 92101-0812, USA

Christopher C. Austin

Department of Biological Sciences and Museum of Natural Sciences, Louisiana State University, Baton Rouge, Louisiana 70803, USA

Abstract

Two new species of *Eimeria* Schneider, 1875, from emerald tree skinks, *Lamprolepis smaragdina* (Lesson) are described from specimens collected in Papua New Guinea (PNG) and the Philippines. Oöcysts of *Eimeria nuiailan* sp. n. from the only *L. smaragdina* from PNG are ovoidal, with a smooth, colourless, bilayered wall, measure $23.7 \times 19.1 \mu\text{m}$, and have a length/width (L/W) ratio of 1.3; both micropyle and oöcyst residuum are absent, but a fragmented polar granule is present. Sporocysts are ovoidal to ellipsoidal, $11.9 \times 7.0 \mu\text{m}$, L/W 1.7, and the wall is composed of 2 valves joined by a longitudinal suture; neither Stieda nor sub-Stieda bodies are present; a sporocyst residuum is present as a compact mass of granules. Sporozoites are elongate, $14.6 \times 2.6 \mu\text{m}$, and contain anterior and posterior refractile bodies with a nucleus between them. Oöcysts of *Eimeria auffenbergi* sp. n. from both *L. smaragdina* we collected in the Philippines are ovoidal, with a smooth, colorless, bilayered wall, measure $19.9 \times 15.8 \mu\text{m}$, L/W 1.3; both micropyle and oöcyst residuum are absent, but 1–4 polar granules are present. Sporocysts are ovoidal to ellipsoidal, $10.3 \times 5.8 \mu\text{m}$, L/W 1.8, and the wall is composed of 2 valves joined by a longitudinal suture; neither Stieda nor sub-Stieda bodies are present; a sporocyst residuum is composed of dispersed granules.

Keywords

Coccidia; *Eimeria*; *Lamprolepis smaragdina*; emerald tree skink; Papua New Guinea; Philippines; taxonomy

Introduction

The emerald tree skink, *Lamprolepis smaragdina* (Lesson) is a fairly common and widespread arboreal lizard that occurs from the Wallacea islands, the Philippines, New Guinea, and Melanesia, to islands of the western Pacific (Perry & Buden, 1999; Linkem et al., 2012). It is one of the most common reptiles in villages, coconut plantations, secondary vegetation and gardens, where it occurs in foliage of bushes and on tree trunks (Foufopoulos & Richards, 2007). The emerald tree skink feeds mainly on beetles, larval lepidopterans, and fruit (Auffenberg & Auffenberg, 1988).

Although information is available on the ecology of *L. smaragdina* (Auffenberg & Auffenberg, 1988; Perry & Buden, 1999; Linkem et al., 2012), little, to our knowledge, has been published on its parasites. For example, Austin & Perkins (2006) examined several *L. smaragdina* for hemoparasites, but did not find any, and Goldberg et al. (2010) examined 40 *L. smaragdina* from PNG and identified 5 nematode species (4 as adult, 1 as encysted larval parasites), but no flatworm or acanthocephalan species. To date, there are no reports of coccidian parasites from this skink. Here, we document the sporulated oöcysts of two coccidian parasites, presumably *Eimeria* species (see Remarks sections), from emerald tree skinks collected from PNG and the Philippines.

Materials and methods

During June, 1991, one adult *L. smaragdina* was collected by hand from New Ireland, PNG, and in June, 2009, two adult *L. smaragdina* were collected by hand from Barangay Zabali, Campus of the Aurora State College of Technology (ASCOT), Luzon, Philippines. Fresh faecal samples were collected from each individual and placed in individual vials containing 2.5% (w/v) aqueous potassium dichromate ($K_2Cr_2O_7$) (Gardner et al., 2012). They were later examined for coccidia by light microscopy after flotation in Sheather's sugar solution (specific gravity = 1.30). All three samples contained unsporulated and partially sporulated oöcysts and they were placed in Petri dishes containing 2.5 % $K_2Cr_2O_7$ and allowed to complete sporulation for five days at room temperature (~23° C). Still later, samples were again examined after flotation and measurements were taken on at least 25 oöcysts using a calibrated ocular micrometer or Olympus® cellSens 1.7 digital imaging software, and reported in micrometres (µm), with means followed by the ranges in parentheses. Photographs were taken using Nomarski interference-contrast (DIC) optics. Oöcysts were between 300 (PNG) and ~1,095 (Philippines) days old when measured and photographed. Descriptions of oöcysts and sporocysts follow guidelines of Wilber et al. (1998). A symbiotype voucher specimen of *L. smaragdina* from PNG was accessioned into the Texas Natural History Collection (TNHC), Austin, Texas, USA, and those from the Philippines were accessioned into Natural History Museum, University of Kansas (KUMNH), Lawrence, Kansas, USA. Photosyntype vouchers of sporulated oöcysts were accessioned into the United States National Parasite Collection (USNPC), Beltsville, Maryland, USA. Lizard taxonomy follows the TIGR reptile database (Uetz 2013) and Zug (2013).

Results

All three *L. smaragdina* from PNG and the Philippines were found to be passing oöcysts of what we believe are two undescribed coccidians; their descriptions follow:

Eimeria nuailan n. sp

Type-host: Emerald treeskink, *Lamprolepis smaragdina* (Lesson) (Sauria: Scincidae), symbiotype TNHC 51465 (field number CCA 754) collected June 1991.

Type-locality: Kavieng, New Ireland Province, PNG (2.566466°S, 150.798547°E), at sea level.

Type-material: Photosyntype (see Duszynski, 1999) as USNPC No. 106965.

Prevalence: 1/1 in the type host.

Sporulation time: Unknown. Specimens were collected in the field, stored in K₂Cr₂O₇ solution, and not examined until 300 days later.

Site of infection: Unknown. Oöcysts were passed in faeces and host tissues were not collected or preserved for histological sectioning.

Prepatent and patent periods: Unknown.

Pathogenicity: Unknown.

Etymology: The specific epithet is given for the Province of New Ireland in Tok Pisin, the official language of PNG. It is a form of Melanesian Pidgin English that was developed in the early 1800's.

Description (Figs. 1, 2, 5)

Sporulated oöcyst: Oöcyst (n = 25): colourless, smooth, ovoidal, 23.7 × 19.1 (21–26 × 17–22); length/width (L/W) ratio 1.3 (1.2–1.4). Wall bi-layered, ×1.2–1.4; outer layer, ~0.8–1.0, inner is ×0.4–0.6. Micropyle absent, oöcyst residuum absent; fragmented polar granule present.

Sporocyst: Sporocysts (n = 20) four, colourless, smooth, ovoidal to ellipsoidal, 11.9 × 7.0 (10–14 × 6–8); L/W ratio 1.7 (1.4–2.1); wall single-layered, ×0.6, composed of 2 valves joined by a longitudinal suture. Stieda body, sub-Stieda and para-Stieda body absent; sporocyst residuum (n = 10) spheroidal to subspheroidal, 4.4 × 3.2 (3–6 × 3–4) composed of compact mass of granules.

Sporozoite: Sporozoite (n = 10), two, elongate, 14.6 × 2.6 (13–17 × 2–3) *in situ*, lying along one side of sporocyst; single ellipsoidal posterior refractile body (n = 5), 3.6 × 2.4 (3–5 × 2–3) spheroidal anterior refractile body (n = 5), 2.0 (1.6–2.2), with nucleus between them.

Remarks—Sporulated oöcysts of eimerians previously reported from skinks (see Modrý & Jirk, 2006) that are most similar to *E. nuiailan* are those of *Eimeria ablephari* Cannon, 1967, from the snake-eyed skink, *Cryptoblepharus* (= *Ablepharus*) *boutonii* (?) (species uncertain) from Australia with oöcysts that measured 23.1 × 17.7, with L/W ratio of 1.3 (Cannon, 1967); oöcysts of *Eimeria beyerae* Ovezmukhammedov, 1977, from the desert lidless skink, *Ablepharus deserti* from northern Turkmenistan (Ovezmukhammedov, 1977) that measured 23.1 × 16.2, L/W ratio 1.4; and oöcysts of *Eimeria zugi* McAllister, Duszynski, Fisher and Austin, 2013, from *Carlia eother* and Admiralty brown skink, *Carlia ailanpalai*, both from PNG, that measured 25.1 × 15.5, L/W ratio 1.6 (McAllister et al., 2013c). However, we believe the oöcysts of *E. nuiailan* can be differentiated from the other three as follows: those of *E. nuiailan* possess a polar granule which both *E. ablephari* and *E. beyerae* lack, and sporocysts of *E. nuiailan* are much larger than those of both *E. ablephari* and *E. beyerae* (11.9 × 7.0 vs. 8.9 × 6.6 and 9.4 × 6.6, respectively), with a larger L/W ratio (1.7 vs. 1.3 and 1.4). Our new species has oöcysts that differ from those of *E. zugi* by having a much smaller oöcyst L/W ratio (1.3 vs. 1.6).

Generic classification of the coccidia, historically, has been based on the structure of the sporulated oocysts recovered from faecal material since this is the stage most readily available to investigators doing parasite survey work. The genus *Eimeria*, characterized by sporulated oocysts with four sporocysts, each containing two sporozoites, has been a “catch-all” for such descriptions, but the tremendous diversity of such forms has prompted recent investigators to look more carefully at those coccidians parasitizing reptiles. Paperna and Landsberg (1989) noted that certain *Eimeria*-like oocysts had specific location and/or developmental patterns within their reptile hosts and erected two new genera: *Acroeimeria*—for those species that develop immediately beneath the brush-border of the intestinal epithelium and later have endogenous stages that extrude to form a layer on the surface of the gut mucosa, and *Choleoeimeria*—for those species that develop exclusively in the biliary tract of reptiles and almost always produce elongate-ellipsoidal oocysts. Sporocysts of species placed in both genera lack both Stieda and sub-Stieda bodies and usually are composed of two valves joined by a longitudinal suture presenting a “bi-valved” appearance. It is certainly possible that the morphotype we describe and name here as *E. nuiailan* may be found by future investigators to have endogenous developmental characteristics to place it in the *Acroeimeria*; however, since we were unable to examine intestinal tissues for endogenous stages nor do we have DNA sequences, we think it is prudent to name this form as a new eimerian until future studies can prove otherwise.

***Eimeria auffenbergi* n. sp**

Type-host: Emerald tree skink, *Lamprolepis smaragdina* (Sauria: Scincidae), Symbiotype KUMNH 3232324, collected 3 June 1999; a second *L. smaragdina*, as KUMNH 323232, collected same date and locality.

Type-locality: Barangay Zabali, Campus of the Aurora State College of Technology (ASCOT), Luzon, Philippines (15°44.59'N, 121°34.54'E), elevation = 50 m.

Type-material: Photosyntype (see Duszynski, 1999) as USNPC No. 106966 a second *L. smaragdina*, as KUMNH 323232, collected by SEB on 3 June 1999 from same locality.

Prevalence: 2/2 of the type host.

Other host and locality: Another *L. smaragdina* (KUMNH 323232) collected on same date and locality as symbiotype.

Sporulation time: Unknown. Specimens were collected in the field, stored in K₂Cr₂O₇ solution, and not examined until 1,095 days later.

Site of infection: Unknown. Oocysts were passed in the faeces and host tissues were not collected or preserved for histological sectioning.

Etymology: The specific epithet is given as a patronym in honor of the late Walter Auffenberg (1928–2004), University of Florida, for his numerous studies on the systematics and biology of reptiles, including skinks from the Philippines.

Description (Figs. 3–4, 6)

Sporulated oocyst: Oocyst (n = 31): colourless, smooth, ovoidal, 19.9 × 15.8 (17–22 × 14–18); length/width (L/W) ratio 1.3 (1.1–1.4). Wall bi-layered, ~0.8–1.2; outer layer, ~0.3–0.7, inner is ~0.2–0.5. Micropyle absent, oocyst residuum absent; polar granules (1–4) present.

Sporocyst: Sporocyst (n = 21) four, colourless, smooth, ellipsoidal, 10.3×5.8 ($9-13 \times 3-7$); L/W ratio 1.8 (1.4–2.4); wall single-layered, ~ 0.5, composed of 2 valves joined by a longitudinal suture. Stieda body, sub-Stieda body and paraStieda body absent; sporocyst residuum is spheroidal to subspheroidal, composed of dispersed granules.

Sporozoite: Sporozoite (not measured), two, elongate, lying along one side of sporocyst; single ellipsoidal posterior refractile body and spheroidal anterior refractile body, with nucleus between them.

Remarks—The following eimerians from skinks are most similar to *E. auffenbergi*: *Eimeria lampropholidis* Cannon, 1967 from the pale-flecked garden sunskink, *Lampropholis guichenoti* from Australia, and *Eimeria leiopismatis* Cannon, 1967 from the orange-tailed shadeskink, *Saproscincus challengerii* from Australia (Cannon, 1967). However, sporulated oöcysts of *E. auffenbergi* can be differentiated from their oöcysts as follows: *E. auffenbergi* has polar granules, which both *E. lampropholidis* and *E. leiopismatis* lack; the sporocysts of *E. auffenbergi* are larger, with a greater L/W ratio, than those of *E. lampropholidis* (10.3×5.8 , L/W 1.8 vs. 7.7×6.5 , L/W 1.2, respectively), and they have smaller sporocysts, with a larger L/W ratio than those of *E. leiopismatis* (10.3×5.8 , L/W 1.8 vs. 11.5×8.1 , L/W 1.4, respectively).

There are at least two other factors we think merit consideration. First, the oöcysts available to us were > 3 yrs old when examined and, in older sporocysts with sutures, it is not uncommon for the walls to disintegrate along the suture-lines, significantly changing their appearance. Thus, in some views of the oöcysts/sporocysts we had to examine, it appeared there may be a tiny Stieda body on the end of some sporocysts; however, we attribute those observations as light refraction of the collapsed sporocyst wall. Our second point is the possibility that the oöcysts we describe and name here as *E. auffenbergi* may be found by future investigators to have endogenous developmental characteristics to place it in the genus *Acroeimeria* (*sensu* Paperna and Landsberg, 1989); however, since we were unable to examine intestinal tissues for endogenous stages and do not have DNA sequences, we think it is prudent to name this form as a new eimerian until empirical evidence is available to prove otherwise.

Discussion

For the last two decades we have been collecting skinks representing eight genera (Carlia, Cryptoblepharus, Emoia, Lipinia, Lamprolepis, Papuascincus, Prasinohaema, Sphenomorphus) from PNG and other South Pacific islands and examining their feces for coccidia. These results are just now being published (McAllister et al., 2013a, b, c, 2014) and it appears that each skink species sampled has discharged oöcysts of at least one coccidian that is unique to it. Herein we have described two new species of *Eimeria* from a single, widely distributed host species (*L. smaragdina*). When compared to other genera examined above, these new eimerians were found only in emerald tree skinks, and each (*E. nuiailan* and *E. auffenbergi*) were also geographically restricted to PNG and the Philippines, respectively. We suggest that the host and geographic specificity of these parasites promises a wealth of undiscovered coccidian diversity that we are just now learning about.

Acknowledgments


We thank the late Steve J. Upton (Kansas State University, Manhattan, KS, USA) for technical assistance, and Scott L. Gardner (Manter Parasite Collection, Lincoln, Nebraska, USA) for parasitological training of RNF. Further appreciation is extended to Patricia A. Piliitt (USNPC) for expert curatorial assistance. The PNG Department of Environment and Conservation and the Protected Areas and Wildlife Bureau (PAWB) of the Philippine Department of Environment and Natural Resources (DENR) facilitated the exportation of lizard and parasite specimens. This

work was funded in part by IBN 9311139 and DEB 1146033 to CCA, by the National Science Foundation BS&I grant DEB 0743491 to R. M. Brown, SEB, D. H. Clayton, and R. G. Moyle, and by grants from the National Center for Research Resources (P20RR016474) and the National Institute of General Medical Sciences (P20GM103432) from the National Institutes of Health to RSS. SEB also thanks V. Tkach and R.M. Brown for assistance in the field. The use of trade, product, or firm names in this publication does not imply endorsement by the U.S. Government and the content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References


- Auffenberg W, Auffenberg T. Resource partitioning in a community of Philippine skinks (Sauria: Scincidae). *Bulletin of the Florida State Museum of Biological Sciences*. 1988; 32:151–219.
- Austin CC, Perkins SL. Parasites in a biodiversity hotspot: A survey of hematozoa and a molecular phylogenetic analysis of *Plasmodium* in New Guinea skinks. *Journal of Parasitology*. 2006; 92:770–777. [PubMed: 16995395]
- Cannon LRG. New coccidia from Australian lizards. II. *Eimeria*. *Parasitology*. 1967; 57:237–250.
- Duszynski DW. Critical comment: revisiting the code: clarifying name-bearing types for photomicrographs of Protozoa. *Journal of Parasitology*. 1999; 85:769–770. [PubMed: 10461969]
- Foufopoulos J, Richards S. Amphibians and reptiles of the New Britain Island, Papua New Guinea: Diversity and conservation status. *Hamadryad*. 2007; 31:176–201.
- Gardner, SL.; Fisher, RN.; Berry, SJ. Chapter 7. Voucher specimens: Collecting and preserving parasites during reptile biodiversity surveys. In: McDiarmid, R.; Foster, M.; Guyer, C.; Gibbons, JE.; Chernoff, N., editors. *Reptile Biodiversity: Standard Methods of Inventory and Monitoring*. University of California Press; Berkeley: 2012. p. 114–121.
- Goldberg SR, Bursey CR, Kraus F. Metazoan endoparasites of fourteen species of skinks (Squamata: Scincidae) from Papua New Guinea. *Journal of Natural History*. 2010; 44:447–467.
- Linkem CW, Brown RM, Siler CD, Evans BJ, Austin CC, Iskandar DT, Diesmos AC, Supriatna J, Andayani N, McGuire JA. Stochastic faunal exchanges drives diversification in widespread Wallacean and Pacific island lizards (Squamata: Scincidae: *Lamprolepis smaragdina*). *Journal of Biogeography*. 2012; 40:507–520.
- McAllister CT, Duszynski DW, Fisher RN. Two new species of *Isospora* (Apicomplexa: Eimeriidae) from skinks, *Emoia* spp. (Sauria: Scincidae) from Fiji and Papua New Guinea. *Journal of Parasitology*. 2013a; 99:677–679. [PubMed: 23517292]
- McAllister CT, Duszynski DW, Fisher RN. New host and distributional records for *Cryptosporidium* sp. (Apicomplexa: Cryptosporidiidae) from lizards (Sauria: Gekkonidae, Scincidae) from the Cook Islands and Vanuatu, South Pacific. *Comparative Parasitology*. 2013b; 80:297–300.
- McAllister CT, Duszynski DW, Fisher RN, Austin CC. A new species of *Eimeria* Schneider (Apicomplexa: Eimeriidae) from *Carlia* spp. (Sauria: Scincidae) from Papua New Guinea. *Systematic Parasitology*. 2013c; 86:53–57. [PubMed: 23949649]
- McAllister CT, Duszynski DW, Austin CC, Fisher RN. Three new species of coccidia (Apicomplexa: Eimeriidae) from skinks, *Lipinia* spp. (Sauria: Scincidae), from Oceania. *Journal of Parasitology*. 2014; 100 in press.
- Modrý D, Jirk M. Three new species of coccidia (Apicomplexa: Eimeriidae) from the marble-throated skink, *Marmorophax tricolor* Bavay, 1869 (Reptilia: Scincidae), endemic to New Caledonia with a taxonomic revision of *Eimeria* spp. from scincid hosts. *Parasitology Research*. 2006; 99:416–428.
- Ovezmukhammadov O. New species of coccidia *Eimeria beyeri* sp. n. of the desert lizard (*Ablepharus deserti* Stauch, 1868) in Turkmenistan. *Izvestiya Akademii Nauk Turkmenskoi SSR Seriya Biologicheskikh Nauk*. 1977; 0(4):83–84.
- Paperna I, Landsberg JH. Description and taxonomic discussion of eimerian coccidians from African and Levantine geckoes. *South African Journal of Zoology*. 1989; 24:345–355.
- Perry G, Buden DW. Ecology, behavior and color variation of the green tree skink, *Lamprolepis smaragdina* (Lacertilia: Scincidae), in Micronesia. *Micronesica*. 1999; 31:263–273.
- UETZ, P. [Accessed 28 April 2013] The TIGR Reptile Database. 2013. World Wide Web electronic publication. <http://www.reptile-database.org/>

- Wilber PG, Duszynski DW, Upton SJ, Seville RS, Corliss JO. A revision of the taxonomy and nomenclature of the *Eimeria* spp. (Apicomplexa: Eimeriidae) from rodents in the Tribe Marmotini (Sciuridae). *Systematic Parasitology*. 1998; 39:113–135.
- Zug, GR. *Reptiles and amphibians of the Pacific Islands: a comprehensive guide*. University of California Press; Berkeley: 2013. p. 320



Figs 1–4.

DIC photomicrographs of oöcysts of *Eimeria nuiailan* n. sp. (1–2) and *Eimeria auffenbergi* n. sp. (3–4). Abbreviations: PG, polar granule; SR, sporocyst residuum; SP, sporocyst; sporozoite, SZ; VS, valve suture. Scale bar: 10 μ m.



Figs. 5–6.

Composite line drawings of oocysts of *Eimeria nuaiailan* sp. n. (5) and *Eimeria auffenbergi* sp. n. (6). Scale-bar: 10 μ m.