4-2003

Valuation of Continuous and Intermittent Phragmites Control

R. Eugene Turner

R. S. Warren

Follow this and additional works at: https://digitalcommons.lsu.edu/oceanography_coastal_pubs

Digital Commons
Part of the [Oceanography Commons](#)
Commons

Network

Logo

Valuation of Continuous and Intermittent *Phragmites* Control

R. E. TURNER^{1,*} and R. S. WARREN²

¹ Coastal Ecology Institute, Louisiana State University, Baton Rouge, Louisiana 70803

² 109 New London Hall, Box 5362, Connecticut College, 270 Mohegan Avenue, New London, Connecticut 06320

ABSTRACT: Quantifying the relative value of *Phragmites australis* and *Spartina alterniflora* habitat is important to evaluate the benefits and risks of different attempts to address *Phragmites* expansion on the U.S. eastern seaboard. Two contrasting approaches commonly used to restore tidal marsh habitats invaded by *Phragmites* communities involve spraying *Phragmites* with herbicide only when its coverage of a particular marsh area is near or close to 100%. Alternatively, after the first application, herbicide is annually applied on any surviving patches of *Phragmites* present in a mosaic of other marsh vegetation. A model is introduced to evaluate the relative habitat value of these control regimes, here termed the Intermittent and Continuous. Compared to the Intermittent approach, the area of herbicide application in the Continuous approach is higher in the first 6 yr, but lower thereafter. The cumulative gain in habitat quality after 20 yr in either approach is sensitive to the presumed relative values of *Phragmites* versus *Spartina* habitat, and may even be negative if they are nearly equal. Annual applications of herbicide to patches of *Phragmites* appears to generate more habitat value and with less herbicide than occasional applications when *Phragmites* cover is at its maximum.

Introduction

Phragmites australis (hereafter, *Phragmites*) expanded inland and seaward throughout East Coast (United States) tidal marshes in the last 50 years (Chambers et al. 1999; Bertness et al. 2002). This expansion, the result of both direct and indirect consequences of anthropogenic modifications to coastal environments and the introduction of new genetic strains, probably from Europe (Besitka 1996; Saltenstall 2002), has been accompanied by concerns that the resulting near monoculture of *Phragmites* compromise various natural attributes that society values (e.g., habitat for fish, fur, and fowl) as well as biodiversity and aesthetic attributes (Weinstein and Balletto 1999). Various programs have been instituted to control its spread, with particular reference to reestablishing displaced tidal marsh vegetation dominated by *Spartina alterniflora*, meadows of *Spartina patens*, *Juncus*, *Eliochoris*, *Scirpus*, and other low-growing graminoids, and lower salinity, tall reed marshes of *Typha* and *Scirpus*. It is interesting to compare this reaction to the loss of *Spartina* sp. habitat on the eastern U.S. coast to that to the increased *Spartina* sp. growth (invasion?) on the western U.S. coast. Both *Phragmites* and *Spartina* sp. habitat expansions are presumed to increase mudflat colonization and sediment accretion and to lower shorebird habitat (Evans 1986; Weinstein and Balletto 1999; Feist and Simenstad 2000; Stevenson et al. 2000). *Spartina* sp. invasion on the West Coast is thought to compro-

mise fish and invertebrate habitat because of the loss of mudflats (Feist and Simenstad 2000). *Phragmites* control on the East Coast is meant to increase the area of *Spartina* sp., not mudflats, with the goal of restoring fish and bird habitat and angiosperm biodiversity (Weinstein and Balletto 1999). The valuation of the same plant, *Spartina* sp., it seems, has a high habitat value on one coast but a low value on another coast.

It costs money and time to control *Phragmites*, and is most often accomplished with herbicide applications followed by burning or mowing to remove standing dead stems. Suitable alternative management strategies to this approach are not yet developed (e.g., rhizome cutting, mowing, sulfide treatments, etc.). We assume that some combination of herbicide application, mowing, and burning will continue to be the dominant approach to controlling *Phragmites* in the near future. This situation makes it very important to know how well spray programs control *Phragmites* expansion, stability or decline, the degree to which they restore more desirable habitats, and at what cost and benefit. To our knowledge there are no published efforts that have systematically weighed the results of various *Phragmites* control approaches or to compare these results with the value of the lost and replaced habitats. A reasonable management and science goal might be to establish these habitat values and to put them in a framework that managers can use to evaluate alternative control approaches. Below we propose a model to support these efforts.

Two different management approaches to control *Phragmites* are what might be called the Inter-

* Corresponding author: tele: 225/578-6454; fax: 225/578-6326; e-mail: euturne@lsu.edu.

TABLE 1. Examples used of *Phragmites* (Pa) and *Spartina* sp. (Sa) plant cover (as a percentage of the sum of plant cover by the two communities in year 0) under intermittent (every 6 yr) herbicide application on *Phragmites* at a hypothetical salt marsh.

Year	%Pa	%Sa	Treatment
0	90%	10%	Prior to spraying
1	0%	100%	After spraying
2	20%	80%	Post spray
3	40%	60%	Post spray
4	60%	40%	Post spray
5	80%	20%	Post spray
6	100%	0%	Post spray
7	0%	100%	Spraying in late summer
8	20%	80%	Post spray
9	40%	60%	Post spray
10	60%	40%	Post spray
11	80%	20%	Post spray
12	100%	0%	Post spray
13	0%	100%	Spraying in late summer
14	20%	80%	Post spray
15	40%	60%	Post spray
16	60%	40%	Post spray
17	80%	20%	Post spray
18	100%	0%	Post spray
19	0%	100%	Spraying in late summer
20	20%	80%	Post spray

mittent and the Continuous spraying approaches. The Intermittent approach involves a one-time spraying of all *Phragmites* stands, usually after its has come to dominate a system. This application is followed by years of inattention until the whole area is again covered by *Phragmites*, after which it is treated again. The Continuous approach also begins with spraying of all *Phragmites* in year one, but is followed by continuous monitoring to identify and treat remaining and new *Phragmites* patches in each of the subsequent years. We determined the habitat value of marshes managed under these two different approaches by constructing a model of a hypothetical marsh managed to control *Phragmites* and comparing this to two managed marshes.

Methods

The three examples discussed below do not represent the whole world of possibilities, and were chosen as polar opposites to illustrate the two management options routinely used in practice.

INTERMITTENTLY MANAGED MARSH COVER

The habitat value was calculated for two marshes under Intermittent management. We assumed that plant cover in a hypothetical marsh was 100% *Spartina* sp. for all of the sprayed *Phragmites* area after spraying (year 1), but that *Spartina* sp. was replaced by *Phragmites* in each of 5 subsequent years (without a herbicide application) at which time there was again 90% *Phragmites* cover on the area sprayed in year 1. After the fifth year, the vegetation control

TABLE 2. *Phragmites* (Pa) and *Spartina* sp. (Sa) plant cover (as a percentage of the sum of plant cover by the two communities in year 0) as a result of Intermittent herbicide application on *Phragmites* (every 9 yr) at the Connecticut River salt marsh restoration site (data are from Warren et al. 2001).

Year	%Pa	%Sa	Treatment
0	90%	10%	Prior to spraying
1	2%	20%	After spraying
2	5%	95%	Post spray
3	10%	90%	Post spray
4	20%	80%	Post spray
5	35%	65%	Post spray
6	55%	45%	Post spray
7	70%	30%	Post spray
8	85%	15%	Post spray-extrapolation
9	95%	5%	Post spray-extrapolation
10	2%	20%	After spraying
11	5%	95%	Post spray-extrapolation
12	10%	90%	Post spray-extrapolation
13	20%	80%	Post spray-extrapolation
14	35%	65%	Post spray-extrapolation
15	55%	45%	Post spray-extrapolation
16	70%	30%	Post spray-extrapolation
17	85%	15%	Post spray-extrapolation
18	95%	5%	Post spray-extrapolation
19	2%	20%	After spraying
20	5%	95%	Post spray-extrapolation

and reestablishment cycle started again. This approach assumes an optimistic management result because, in practice, *Spartina* spp. cover does not reach 90% in year 1 (post-spray) and significant amounts of *Phragmites* may survive the herbicide treatment. The return of both *Spartina* spp. and *Phragmites* following herbicide is spatially and temporally variable (PSE&G 2001; Warren et al. 2001).

The second data set is from a marsh at the Connecticut River salt marsh restoration site described by Warren et al. (2001) and is for a marsh managed on a 9-yr spray rotation. Data beyond year 6 do not exist, and so we estimated the cover in years 7, 8, and 9 by extrapolation. The data for both marshes are in Tables 1 and 2.

CONTINUOUSLY MANAGED MARSH COVER

The vegetative cover of *Phragmites* and *S. alterniflora* was obtained for 4 salt marsh restoration sites in the Delaware estuary that were treated with a herbicide (Rodeo and Li-700 surfactant) for at least 3 consecutive years. These 4 sites are part of a 4,000 ha salt marsh restoration program (Weinstein and Balletto 1999). The data used are from annual reports of the monitoring program of the Public Service Electric and Gas Co., Inc. (PSE&G) that are provided to the New Jersey Department of Environmental Protection as part of regulatory oversight for a nuclear power plant discharge permit. Vegetation on mudflats is excluded from these analyses, which are the result of interpretations of aerial photos taken in the spring and fall of several

Fig. 1. The ratio (log transformed) of the plant cover of *Phragmites* and *Spartina* sp. for four *Phragmites* control programs in Delaware Bay estuary that had continuous (annual) applications of Rodeo herbicide to the *Phragmites* plant cover.

consecutive years before spraying began and also for each year of spraying. The maximum area with both plant communities at these 4 sites is 147, 199, 269, and 598 ha. The *Phragmites* and *S. alterniflora* cover ratios were transformed into logarithmic values and plotted against the years of consecutive spraying (Fig. 1). A simple linear regression of this relationship was accomplished using Statview II version 1.04 software.

The Continuous approach assumes the same first year herbicide treatment as above, followed by additional spraying of surviving and new *Phragmites* patches in all following years. Table 2 presents data from the Cedar Swamp restoration site (New Castle County, Delaware) as an example of *Phragmites* control using annual herbicide applications targeted at surviving or new *Phragmites* stands (PSE&G 2001). Cedar swamp had 90% *Phragmites* coverage before restoration began, which suggests the presence of a more tenacious *Phragmites* management problem there than in the more saline marshes of Delaware Bay.

DETERMINATION OF HABITAT VALUE

A habitat value (HV) of 0, 0.25, 0.5, 0.75, and 1.0 was assigned to *Phragmites*-dominated marshes, where 1.0 is the equal to a *Spartina* spp. marsh in any given year and 0 = no value. We do not define the basis for the determination of HV, which might involve floral, fauna, aesthetic, or political interests, among other things. The average HV of a marsh was calculated by proportioning the HV of both species among their respective coverage in an individual year. The cumulative habitat value (Σ HV) for up to 20 yr is the cumulative sum of the HV in all years. The net marsh Σ HV is the gain

(loss) in Σ HV under all management scenarios less the Σ HV in the same marsh without herbicide application (i.e., habitat conditions in year 0 are perpetuated to year 20).

AREA SPRAYED

The cumulative area sprayed with herbicide under Intermittent and Continuous approaches was estimated for 20 yr, using the data for plant cover for the 3 marshes described above. We assumed for purposes of this exercise that the herbicide used poses no significant or immediate health or other habitat risks.

Results

The ratio of *Phragmites:Spartina* plant cover declines in a linear manner for the first 4 years of herbicide application using the continuous approach (Fig. 1). The R^2 values for a linear regression of the ratio (Y) versus year (X) ranges from 0.70 to 0.99. Spraying for more than 3 consecutive years results in less *Phragmites* cover than *Spartina* sp. coverage. Four years of continuous spraying results in a *Phragmites* cover of less than 10%. The changes in plant cover in these four marshes is such that it may take 5 consecutive years of spraying before 5% or less of the land is covered with *Phragmites* habitat, and the ultimate level of *Phragmites* cover after multiple years of spraying is not known. We wish to emphasize that the choice of these two different spray programs or approaches was for illustrative purposes and that the results should not be expected at every site. *Phragmites* cover will probably increase if the spray program is reduced or stops, and the recovery rates will vary depending upon a number of local variables (e.g., salinity, elevation).

The annual HV, Σ HV, and net Σ HV for the 3 marshes under 5 different HV for *Phragmites* are shown in Fig. 2. The marsh HV swings widely under the 5 HV for *Phragmites* cover, as expected. There is a gain in marsh HV with each year, although the continuous spraying example shows a lag in Σ HV for the first few years. The net gain in Σ HV varies from zero or less to positive in all 3 marsh types (Fig. 2, Table 3). The marsh under Continuous management has the highest valuation of all three marshes after 20 yr when the HV for *Phragmites* is <0.5.

Continuous management results in a net loss of habitat up to 10 yr if the HV of *Phragmites* is > 0.75, and up to 10 yr if the HV of *Phragmites* equals 1. Habitat losses become gains by year 3 when the *Phragmites* habitat value is >0.5, and within 5 yr if the *Phragmites* value is 0.75 or higher (Fig. 2).

The cumulative area sprayed in years 9 through 15 is about the same at all marshes, after which the

Fig. 2. The habitat value (HV) of three marshes managed under the different management scenarios described in the text. Five different valuations of *Phragmites* plant cover are shown, ranging from 1 (equal value to *Spartina* sp.) to 0 (no value). The arrow on the x axis indicate when the herbicide application occurs. Spraying occurs in all years in the Cedar Swamp example. A) Annual value. B) Cumulative value. C) Net value (the difference between the Σ HV under managed and no-management scenarios).

marsh under Continuous management show minor increases in comparison to the large gains in the other two marshes (Fig. 3). The area sprayed under the Intermittent approach continues to rise (Fig. 3) until the cumulative amount is 40% greater than in the Continuous management program by year 20 (and continues to rise thereafter; Table 3).

Discussion

Spartina cover increases through herbicide application (Fig. 1), but is replaced by *Phragmites* when herbicide applications cease. It would improve management choices to know the relationship between herbicide application and vegetation response for a variety of other sites. The goal of the scientific approach to this management problem should be, and can be, to establish the trajectory of the cost:benefit ratio under different man-

Fig. 3. The cumulative area sprayed with herbicide of a marsh in each year for 20 years under three different management scenarios described in the text and Table 2. The area is normalized to a 100 ha marsh.

agement practices. Does *Phragmites* self-restoration occur at a faster rate if sprayed every year or in alternate years? Can a small herbicide application every year, by hand, reduce the total herbicide application amount anticipated if done every third year? Do different marshes types (salinity, elevation) respond in predictable ways? Do sites have a maximum restoration level, beyond which additional spraying has no discernible impact? If this information is collected and analyzed in a carefully documented and systematic manner, then the arguments about herbicide control of *Phragmites* can involve not only qualitative arguments, but also quantitative appraisals.

The HV for the two approaches clearly demonstrates changes over 20 yr (Fig. 2). The net HV for a marsh under Intermittent spraying in the hypothetical example results in a net loss of habitat after 5 yr if the HV of *Phragmites* is > 0.75 compared to *Spartina* sp., and a net gain after 1 yr if the *Phragmites* HV is < 0.5 (Fig. 2). These are optimal results, because *Spartina* sp. does not recolonize sprayed *Phragmites* vegetative zones as quickly as *Phragmites* cover is diminished after one herbicide application. For example, *Spartina* sp. cover was essentially unchanged at the Cedar Creek marsh restoration site even after 3 consecutive years of spraying (Table 4).

The root paradigm about the way *Phragmites* is currently managed is largely that it is an either/or situation, where *Phragmites* has a zero value compared to *Spartina* sp. When sprayed the first year,

TABLE 3. Summary of results for 20 years of 3 different *Phragmites* control programs.

Example	Spray Program	Spray Interval (yr)	Cumulative Sprayed Area (ha)	% Net Habitat Gain when <i>Phragmites</i> HV =				
				0	0.25	0.5	0.75	1
Hypothetical	Intermittent	6	270	405	88	32	8	-5
Connecticut River	Intermittent	9	280	416	100	43	19	6
Cedar Swamp	Continuous	each year	195	703	154	55	13	-10

TABLE 4. *Phragmites* (Pa) and *Spartina* sp. (Sa) plant cover (as a percentage of the sum of plant cover by the two communities in year 0) as a result of Intermittent herbicide application on *Phragmites* (every 9 yr) at the Cedar Swamp salt marsh restoration site in the Delaware Bay estuary.

Year	%Pa	%Sa	Treatment
0	90%	10%	Prior to spraying
1	28%	11%	Post spray
2	8%	4%	Spraying remnant patches
3	11%	14%	Spraying remnant patches
4	14%	51%	Spraying remnant patches
5	15%	63%	Spraying remnant patches
6	7%	75%	Spraying remnant patches
7	5%	95%	Spraying remnant patches-extrapolation
8	4%	96%	Spraying remnant patches-extrapolation
9	3%	97%	Spraying remnant patches-extrapolation
10	1%	99%	Spraying remnant patches-extrapolation
11	1%	99%	Spraying remnant patches-extrapolation
12	1%	99%	Spraying remnant patches-extrapolation
13	1%	99%	Spraying remnant patches-extrapolation
14	1%	99%	Spraying remnant patches-extrapolation
15	1%	99%	Spraying remnant patches-extrapolation
16	1%	99%	Spraying remnant patches-extrapolation
17	1%	99%	Spraying remnant patches-extrapolation
18	1%	99%	Spraying remnant patches-extrapolation
19	1%	99%	Spraying remnant patches-extrapolation
20	1%	99%	Spraying remnant patches-extrapolation

regardless of the two approaches, both *Phragmites* and *Spartina* sp. are typically greatly reduced. The expectation is that, in the long term, there will be a net gain in HV as a result of subsequent *Spartina* sp. growth. If the HV of *Phragmites* is half that of *Spartina* sp., then it takes a longer time to realize a net gain in marsh HV than if the HV of *Phragmites* is zero. The result depends what the relative value of *Phragmites* habitat—on whose ox (amphipod) gets gored. Intermittent management (under these assumptions) will not result in a net habitat gain for 5 or more years under circumstances of relative high *Phragmites* valuation (Fig. 2). Under continuous spraying, it might take 7 or 8 yr before there is a net habitat gain under most HV for *Phragmites* of 0.5, and much longer if at all, if the HV of *Phragmites* approaches 1 (Fig. 2). The point to be made here is that the length of time to evaluate results is important; how *Phragmites* is valued is important, and that the HV of *Phragmites* is probably not zero or 100% (equivalent to *Spartina* sp.), but something in-between, which is presently difficult to evaluate; and spraying once a year every 5 or 6 yr may not result in a net gain in habitat value, but will use much more herbicide in comparison to that used in the Continuous approach (Fig. 2, Table 3). By spraying remnant patches every year, then there may be a point where spraying only a few percent of the marsh each year is necessary to control the areal cover of *Phragmites*.

The determination of the relative value of *Spartina* sp. and *Phragmites* is a crucial to the evaluation

of management choices. *Phragmites* habitat value is probably not going to be equal to *Spartina* sp. for many issues, but it is not to be zero, either. For example, if the perceived HV of *Phragmites* to fish is 0 (the wolf in sheep's clothing), then any habitat control through spraying is beneficial from a fisheries habitat viewpoint, but not if the value is > 0.75. Fell et al. (1998), Able and Hagan (2000), Weinstein and Litvin (2000), Wainwright et al. (2000), and Weis et al. (2002) have shown that the value of *Phragmites* for larval, small, and large juvenile fish, and for decapods is greater than zero and may sometimes be close to 1.0 (the sheep in wolf's clothing).

Whether the costs of spraying are worth the benefits may involve questions about the health threats of the herbicide itself. If herbicide use is to be minimized, then an annual spray program of small amounts will result in much less herbicide applied than if the spraying is done on a less frequent basis or erratically. Suppose that 5% of a marsh can be hand-treated every year to keep it at 98% *Spartina*, but another marsh requires aerial spraying of 40% every year to keep it at 60% *Spartina*. Given our present knowledge, these two alternative scenarios are reasonable outcomes, depending upon salinity and other environmental variables. But they involve vastly different amounts of herbicides, and at different exposure rates, expenses, and commitments by management and public.

Useful improvements to the approach discussed herein include refined estimates of invasion and reestablishment rates and accurate descriptions of the variability in plant cover before restoration starts and the decline in *Phragmites:Spartina* ratios after restoration begins. Additional information on the interactive effects of burning, mowing, and timing of spray operations are also desirable.

ACKNOWLEDGMENTS

The authors thank the organizers of the Technical Forum, *Phragmites australis*: A Sheep in Wolf's Clothing? for the invitation to attend the meeting. The scientists and managers with the PSE&G Estuary Enhancement Program, including J. Balletto, B. Evans, G. Bickle, R. Hinkle, J. Pantazes, K. Philipp, K. Strait, J. Teal, and L. Weishar, were part of many stimulating discussions concerning the salt marsh restoration program in Delaware Bay. The company of M. Bruno, J. Gallagher, W. Meredith, W. Mitsch, J. Shisler, and L. Widjeskog when visiting the *Phragmites* communities in these marshes was always a treat.

LITERATURE CITED

- ABLE, K. AND W. S. M. HAGAN. 2000. Effects of common reed (*Phragmites australis*) invasion on marsh surface macrofauna: Response of fishes and decapod crustaceans. *Estuaries* 23:633–646.
- BERTNESS, M. D., P. J. EWANCHUK, AND B. R. SILLIMAN. 2002. Anthropogenic modification of New England salt marsh landscapes. *Proceeding of the National Academy of Sciences (USA)* 99: 1395–1398.

- BESITKA, M. R. A. 1996. An ecological and historical study of *Phragmites australis* along the Atlantic coast. Masters Thesis, Drexel University, Philadelphia, Pennsylvania.
- CHAMBERS, R. M., L. A. MEYERSON, AND K. SALTONSTALL. 1999. Expansion of *Phragmites australis* into tidal wetlands of North America. *Aquatic Botany* 64:261–273.
- CHAMBERS, R. M., D. T. OSGOOD, AND N. KALAPASEV. 2002. Hydrologic and chemical control of *Phragmites* growth in tidal marshes of SW Connecticut, USA. *Marine Ecology Progress Series* 239:83–91.
- EVANS, P. R. 1986. Use of herbicide 'Dalapon' for control of *Spartina* encroaching on intertidal mudflats: Beneficial effects on shorebirds. *Colonial Waterbirds* 9:171–175.
- FEIST, B. E. AND C. A. SIMENSTAD. 2000. Expansion rates and recruitment frequency of exotic smooth cordgrass, *Spartina alterniflora* (Loisel), colonizing unvegetated littoral flats in Willapa Bay, Washington. *Estuaries* 23:267–274.
- FELL, P., S. P. WISSBACH, D. A. JONES, M. A. FALLON, J. A. ZEPPIERI, E. K. FAISON, K. A. LENNON, K. J. NEWBERRY, AND L. K. REDDINGTON. 1998. Does invasion of oligohaline tidal marshes by reed grass, *Phragmites australis* (Cac.) Trin. ex Steud., affect the availability of prey sources for the mummichog, *Fundulus heteroclitus* L. *Journal Experimental Marine Biology and Ecology* 222:59–77.
- PSE&G. 2001. Cedar Swamp and the rocks *Phragmites*-dominated wetland restoration sites, Blackbird and Appoquinimink Hundreds, New Castle County, Delaware. 2000 site status report dated 29 June 2001, to the New Jersey Department of Environmental Protection—Land Use Regulation Program. U.S. Army Corps of Engineers and Management Plan Advisory Committee by Public Service Electric and Gas Co., Inc., Salem, New Jersey.
- SALTONSTALL, K. 2002. Cryptic invasion by a non-native genotype of the common reed, *Phragmites australis*, into North America. *Proceeding of the National Academy of Science (USA)* 99: 2445–2449.
- STEVENSON, C., J. E. ROTH, M. S. KEARNEY, AND K. L. SUNDERG. 2000. The health and long term stability of natural and restored marshes in Chesapeake Bay, p. 709–735. In M. Weinstein and K. Kreeger (eds.), *Concepts and Controversies in Tidal Marsh Ecology*. Kluwer Academic Publishing, Dordrecht, The Netherlands.
- WAINWRIGHT, S. C., M. P. WEINSTEIN, K. W. ABLE, AND C. A. CURRIN. 2000. Relative importance of benthic microalgae, phytoplankton and the detritus of smooth cordgrass (*Spartina*) and the common reed (*Phragmites*) to brackish marsh food webs. *Marine Ecology Progress Series* 200:77–91.
- WARREN, R. S., P. E. FELL, J. L. GRIMSBY, E. L. BUCK, C. G. RILLING, AND R. A. FERTIK. 2001. Rates, patterns, and impacts of *Phragmites australis* expansion and effects of experimental *Phragmites* control on vegetation, macroinvertebrates, and fish within the tidelands of the lower Connecticut River. *Estuaries* 24:90–107.
- WEINSTEIN, M. P. AND J. H. BALLETO. 1999. Does the common reed, *Phragmites australis*, affect essential fish habitat? *Estuaries* 22:793–202.
- WEINSTEIN, M. P. AND S. Y. LITVIN. 2000. The role of tidal salt marsh as an energy source for marine transient and resident finfishes: A stable isotope approach. *Transactions American Fisheries Society* 129:797–2000.
- WEIS, J. S., L. WINDHAM, C. SANTIAGO-BASS, AND P. WEIS. 2002. Growth, survival, and metal content of marsh invertebrates fed diets of detritus from *Spartina alterniflora* Loisel. and *Phragmites australis* Cav. Trin. ex Steud. from metal-contaminated and clean sites. *Wetlands Ecology and Management* 10:71–84.

Received for consideration, May 7, 2002

Revised, September 23, 2002

Accepted for publication, October 15, 2002