6-1-2013

Partially Gaussian stationary stochastic processes in discrete time

K R Parthasarathy

Follow this and additional works at: <https://digitalcommons.lsu.edu/cosa>

Part of the [Analysis Commons](#), and the [Other Mathematics Commons](#)

Recommended Citation

Parthasarathy, K R (2013) "Partially Gaussian stationary stochastic processes in discrete time," *Communications on Stochastic Analysis*: Vol. 7 : No. 2 , Article 2.

DOI: 10.31390/cosa.7.2.02

Available at: <https://digitalcommons.lsu.edu/cosa/vol7/iss2/2>

PARTIALLY GAUSSIAN STATIONARY STOCHASTIC PROCESSES IN DISCRETE TIME

K. R. PARTHASARATHY

ABSTRACT. We present here an elementary example, for every fixed positive integer k , of a strictly stationary nongaussian stochastic process in discrete time, all of whose k -marginals are gaussian.

1. Introduction

It is well known that, for every positive integer $n > 1$, there exists a probability distribution in $\mathbb{R}^n$ which is not gaussian but has all its $(n-1)$ -dimensional marginal distributions gaussian. (See, for example, Section 10.3 in Stoyanov [2].) Using the finer theory of pathwise stochastic integrals and martingale methods, Föllmer, Wu and Yor [1] have shown that, for every positive integer k , there exists a variety of nongaussian stochastic processes with continuous trajectories in the interval $[0, \infty)$ which have the same k -dimensional marginals as the standard brownian motion process. Here we present an elementary example, for every fixed positive integer $k > 1$, a discrete time stationary stochastic process which is not gaussian but has all its $(k-1)$ -marginals gaussian. However, we do not know how to construct such processes in continuous time.

2. The Basic Construction

Let $k > 1$ be any fixed positive integer and let ν be a probability distribution in $\mathbb{R}^k$, which is not gaussian but has all its $(k-1)$ -marginals gaussian with mean $\mathbf{0}$ and covariance matrix identity. For example, we may choose ν to have the probability density function

$$\psi(\mathbf{x}) = (2\pi)^{-\frac{k}{2}} \left\{ 1 + x_1 x_2 \cdots x_k e^{-\frac{1}{2}|\mathbf{x}|^2} \right\} e^{-\frac{1}{2}|\mathbf{x}|^2}$$

where $\mathbf{x} = (x_1, x_2, \dots, x_k)$. If $(X_1, X_2, \dots, X_k)$ is an $\mathbb{R}^k$ -valued random variable with distribution ν , then the sequence $X_1, X_2, \dots, \widehat{X}_i, \dots, X_k$ with the i -th term omitted consists of i.i.d. $N(0, 1)$ random variables, for each i .

Now consider a bilateral sequence $\{(X_{n1}, X_{n2}, \dots, X_{nk}), -\infty < n < \infty\}$ of i.i.d. $\mathbb{R}^k$ -valued random variables with the common distribution ν as described in the preceding paragraph. Define

$$Y_n = X_{nk} + X_{n+1, k-1} + X_{n+2, k-2} + \cdots + X_{n+k-1, 1}, -\infty < n < \infty.$$

Received 2012-12-17; Communicated by Hui-Hsiung Kuo.

2000 *Mathematics Subject Classification.* Primary 60G15, 60G10, 60G09.

Key words and phrases. Gaussian distribution, strictly stationary process, partially gaussian process.

It is to be noted that the sum of the two suffixes in each summand on the right hand side is equal to $n + k$.

Theorem 2.1. *The sequence $\{Y_n, -\infty < n < \infty\}$ is strictly stationary, $(k-1)$ -step independent with every $(k-1)$ -dimensional marginal being gaussian with mean $\mathbf{0}$ and covariance matrix kI , I being the identity matrix of order $k-1$. In particular, $\{Y_n\}$ is ergodic.*

Proof. Fix an integer m and consider the two sets $\{Y_n, n \leq m\}$ and $\{Y_n, n \geq m+k\}$. Since $Y_m = X_{mk} + X_{m+1\ k-1} + \cdots + X_{m+k-1\ 1}$ and $Y_{m+k} = Y_{m+k\ k} + X_{m+k+1\ k-1} + \cdots + X_{m+2\ k-1\ 1}$ and the first suffix in the last summand in the definition of Y_m is less than the first suffix in the first summand in the definition of Y_{m+k} it follows that the two sets $\{Y_n, n \leq m\}$ and $\{Y_n, n \geq m+k\}$ are independent. In other words $\{Y_n\}$ is a $(k-1)$ -step independent process.

We now look at the column vector-valued random variable

$$\begin{bmatrix} Y_{n+1} \\ Y_{n+2} \\ \vdots \\ Y_{n+m} \end{bmatrix} = \begin{bmatrix} X_{n+1\ k} + X_{n+2\ k-1} + \cdots + X_{n+k\ 1} \\ X_{n+2\ k} + X_{n+3\ k-1} + \cdots + X_{n+k+1\ 1} \\ \vdots \\ X_{n+m\ k} + X_{n+m+1\ k-1} + \cdots + X_{n+m+k-1\ 1} \end{bmatrix}$$

and express it as $S_1 + S_2 + S_3$ where

$$\begin{aligned} S_1 &= \begin{bmatrix} X_{n+1\ k} \\ 0 \\ \vdots \\ 0 \end{bmatrix} + \begin{bmatrix} X_{n+2\ k-1} \\ X_{n+2\ k} \\ 0 \\ \vdots \\ 0 \end{bmatrix} + \cdots + \begin{bmatrix} X_{n+k-1\ 2} \\ X_{n+k-1\ 3} \\ \vdots \\ X_{n+k-1\ k} \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \\ S_2 &= \begin{bmatrix} X_{n+k\ 1} \\ X_{n+k\ 2} \\ \vdots \\ X_{n+k\ k} \\ 0 \\ \vdots \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ X_{n+k+1\ 1} \\ X_{n+k+1\ 2} \\ \vdots \\ X_{n+k+1\ k} \\ 0 \\ \vdots \\ 0 \end{bmatrix} + \cdots + \begin{bmatrix} 0 \\ \vdots \\ 0 \\ X_{n+m\ 1} \\ X_{n+m\ 2} \\ \vdots \\ X_{n+m\ k} \end{bmatrix}, \\ S_3 &= \begin{bmatrix} 0 \\ \vdots \\ 0 \\ X_{n+m+1\ 1} \\ X_{n+m+1\ 2} \\ \vdots \\ X_{n+m+1\ k-1} \end{bmatrix} + \begin{bmatrix} 0 \\ \vdots \\ 0 \\ X_{n+m+2\ 1} \\ X_{n+m+2\ 2} \\ \vdots \\ X_{n+m+2\ k-2} \end{bmatrix} + \cdots + \begin{bmatrix} 0 \\ \vdots \\ \vdots \\ 0 \\ X_{n+m+k-1\ 1} \end{bmatrix}. \end{aligned}$$

In each column on the right hand side of S_1 or S_3 there are at most $k-1$ nonzero entries whereas in each column on the right hand side of S_2 there are exactly k entries. All the column vectors appearing in S_1, S_2, S_3 together are mutually independent. By the choice of the measure ν , S_1 and S_3 are gaussian random vectors. Denote by $\mu([i, j])$ the $(j-i+1)$ -dimensional standard normal distribution imbedded in $\mathbb{R}^m$ so that the first $i-1$ and the last $m-j$ coordinates are 0 when $1 \leq i \leq j \leq m$. Similarly, denote by $\nu([j, k+j-1])$ the k -dimensional distribution ν imbedded in $\mathbb{R}^m$ with the first $j-1$ and the last $m-k-j+1$ coordinates 0 for $1 \leq j \leq m-k+1$, assuming $m \geq k$. Then it follows that the random variables $Y_{n+1}, Y_{n+2}, \dots, Y_{n+m}$ expressed as a single column vector has the m -dimensional distribution ν_m (in $\mathbb{R}^m$) given by

$$\begin{aligned} \nu_m &= \mu([1, 1]) * \mu([1, 2]) * \dots * \mu([1, k-1]) \\ &\quad * \nu([1, k]) * \nu([2, k+1]) * \dots * \nu([m-k+1, m]) \\ &\quad * \mu([m-k+2, m]) * \mu([m-k+3, m]) * \dots * \mu([m, m]), \end{aligned}$$

for every $m \geq k$. Since ν_m is independent of n it follows that $\{Y_n\}$ is a strictly stationary process. Since $\nu([1, k])$ is nongaussian it is clear that ν_m is not gaussian for every $m \geq k$.

We now observe that Y_n , being a sum of k independent $N(0, 1)$ random variables, is an $N(0, k)$ variable with mean 0 and variance k . Now consider the pair (Y_0, Y_m) . If $m \geq k$ we have already seen that Y_0 and Y_m are independent. If $m < k$, we write

$$\begin{aligned} \begin{bmatrix} Y_0 \\ Y_m \end{bmatrix} &= \begin{bmatrix} X_{0k} + X_{1 \ k-1} + \dots + X_{m-1 \ k-m+1} \\ 0 \end{bmatrix} + \begin{bmatrix} X_{m \ k-m} \\ X_{m \ k} \end{bmatrix} \\ &\quad + \begin{bmatrix} X_{m+1 \ k-m-1} \\ X_{m+1 \ k-1} \end{bmatrix} + \dots + \begin{bmatrix} X_{k-1 \ 1} \\ X_{k-1 \ m+1} \end{bmatrix} \\ &\quad + \begin{bmatrix} 0 \\ X_{km} + X_{k+1 \ m-2} + \dots + X_{k+m-1 \ 1} \end{bmatrix}. \end{aligned}$$

Now the special choice of ν implies that Y_0 and Y_m are independent $N(0, k)$ random variables. Stationarity of the process $\{Y_n\}$ implies that Y_{n_1} and Y_{n_2} are independent $N(0, k)$ random variables for any n_1, n_2 .

Now consider, for any $n_1 < n_2 < \dots < n_{k-1}$ the random vector

$$\tilde{\mathbf{Y}} = \begin{bmatrix} Y_{n_1} \\ Y_{n_2} \\ \vdots \\ Y_{n_{k-1}} \end{bmatrix} = \begin{bmatrix} X_{n_1 k} + X_{n_1+1 \ k-1} + \dots + X_{n_1+k-1 \ 1} \\ X_{n_2 k} + X_{n_2+1 \ k-1} + \dots + X_{n_2+k-1 \ 1} \\ \vdots \\ X_{n_{k-1} k} + X_{n_{k-1}+1 \ k-1} + \dots + X_{n_{k-1}+k-1 \ 1} \end{bmatrix}.$$

The right hand side can be expressed as a sum of column vectors in which the entries in each column are either 0 or an X_{rs} where the first suffix r is fixed and the second suffix takes at most $k-1$ values from the set $\{1, 2, \dots, k\}$. The different column vectors are independent and by the choice of ν each column has a multivariate gaussian distribution. Thus $\tilde{\mathbf{Y}}$ is gaussian. Since any two Y_i and Y_j are independent where $k > 2$, it follows that $Y_{n_1}, Y_{n_2}, \dots, Y_{n_{k-1}}$ are i.i.d $N(0, k)$ random variables. This completes the proof. $\square$

Remark 2.2. From the proof of Theorem 2.1 it is clear that any $k - 1$ of the random variables $\{Y_n\}$ are i.i.d $N(0, k)$. This motivates the introduction of the following notion of limited exchangeability. We say that a stationary random process $\{Z_n, -\infty < n < \infty\}$ is *k-exchangeable* if any $Z_{n_1}, Z_{n_2}, \dots, Z_{n_k}$ has the same distribution for any k -point set $\{n_1, n_2, \dots, n_k\} \subset \mathbb{Z}$. The probability measures of all such k -exchangeable stationary processes constitute a convex set. One wonders what are the extreme points of this convex set.

References

1. Föllmer, H., Wu, C. T., and Yor, M.: On weak brownian motions of arbitrary order, *Ann. Inst. Henri Poincaré* **36** (2000) 447–487.
2. Stoyanov, J.: *Counterexamples in Probability*. Wiley, 1987.

K. R. PARTHASARATHY: THEORETICAL STATISTICS & MATHEMATICS UNIT, INDIAN STATISTICAL INSTITUTE, DELHI CENTRE, 7, S. J. S. SANSANWAL MARG, NEW DELHI - 110016, INDIA
E-mail address: `krp@isid.ac.in`