


6-1-2015

Regular transformation groups based on Fourier-Gauss transforms

Maximilian Bock

Wolfgang Bock

Follow this and additional works at: <https://digitalcommons.lsu.edu/cosa>

 Part of the [Analysis Commons](#), and the [Other Mathematics Commons](#)

Recommended Citation

Bock, Maximilian and Bock, Wolfgang (2015) "Regular transformation groups based on Fourier-Gauss transforms," *Communications on Stochastic Analysis*: Vol. 9 : No. 2 , Article 3.

DOI: 10.31390/cosa.9.2.03

Available at: <https://digitalcommons.lsu.edu/cosa/vol9/iss2/3>

REGULAR TRANSFORMATION GROUPS BASED ON FOURIER-GAUSS TRANSFORMS

MAXIMILIAN BOCK AND WOLFGANG BOCK

ABSTRACT. We discuss different representations of the white noise test functions $(E)_\beta$, $0 \leq \beta < 1$ by introducing generalized Wick tensors. As an application we state the following generalization of the Mehler formula for the Ornstein-Uhlenbeck semigroup, see [7, p. 237]: Let $a, b \in \mathbb{C}$, $0 \leq \beta < 1$. Then $a \cdot \Delta_G + b \cdot N$ is the infinitesimal generator of the following regular differentiable one parameter transformation group $\{P_{a,b,t}\}_{t \in \mathbb{R}} \subset GL((E)_\beta)$:

(i) if $b \neq 0$ then for all $\varphi \in (E)_\beta$, $t \in \mathbb{R}$:

$$P_{a,b,t}(\varphi) = \int_{E^*} \varphi\left(\sqrt{\left(1 - \frac{a}{b}\right)(1 - e^{2bt})} \cdot x + e^{bt} \cdot y\right) d\mu(x)$$

(ii) if $b = 0$ then for all $\varphi \in (E)_\beta$, $t \in \mathbb{R}$:

$$P_{a,0,t}(\varphi) = \int_{E^*} \varphi(\sqrt{2at} \cdot x + y) d\mu(x)$$

On an informal level the second case of the above theorem may be looked upon as a special case of the first one. Note that by the rules of l'Hôpital we have $\lim_{b \rightarrow 0} (b - a) \frac{1 - e^{2bt}}{b} = 2at$.

1. Introduction

In finite dimensional analysis convolution and Fourier transform appear in many different applications. Since the Fourier transform is symmetric with respect to the dual pairing it extends naturally to generalized functions. The Fourier transform on $(E)_\beta^*$ resembles the finite dimensional Fourier transform in the sense that it is the adjoint of a continuous linear operator from the space of test functions into itself. This adjoint is a so called Fourier Gauss transform. In Gaussian Analysis, the Fourier-Gauss transform $\mathfrak{G}_{a,b}(\varphi)$ of $\varphi \in (E)_\beta$ is defined by

$$\mathfrak{G}_{a,b}(\varphi)(y) = \int_{E^*} \varphi(ax + by) d\mu(x), \quad a, b \in \mathbb{C}, \quad 0 \leq \beta < 1$$

Received 2015-6-12; Communicated by Hui-Hsiung Kuo.

2010 *Mathematics Subject Classification*. Primary 60H40; Secondary 46F25.

Key words and phrases. White noise theory, Gross Laplace operator, number operator, Fourier-Gauss transform, infinitesimal generator, convolution operator, generalized Wick tensors.

see e.g. [7], [11] and [10]. The Fourier-Gauss transform is in $L((E)_\beta, (E)_\beta)$ and the operator symbol is given by

$$\widehat{\mathfrak{G}}_{a,b}(\xi, \eta) = \exp \left[\frac{1}{2} (a^2 + b^2 - 1) \langle \xi, \xi \rangle + b \langle \xi, \eta \rangle \right], \quad \text{for all } \xi, \eta \in E_{\mathbb{C}},$$

see [10, Theorem 11.29, p. 168-169]. Hence

$$\mathfrak{G}_{a,b} = \Gamma(b \text{ Id}) \circ \exp \left(\frac{1}{2} (a^2 + b^2 - 1) \Delta_G \right), \quad (1.1)$$

where Id denotes the identity operator. Considering this formula, we show that we can find a suitable representation of the white noise test functions, such that equation (1.1) will be just consisting of second quantization operators and is thus comparable to the approach in [2]. This will be done by introducing generalized Wick tensors. For $m \in \mathbb{N}$, $\kappa_{0,m} \in (E_{\mathbb{C}}^{\otimes m})_{sym}^*$ we define generalized Wick tensors by

$$: x^{\otimes n} :_{\kappa_{0,m}} = \sum_{k=0}^{\lfloor \frac{n}{m} \rfloor} \frac{n!}{(n - mk)! k!} \left(-\frac{1}{2} \right)^k x^{\otimes (n - mk)} \widehat{\otimes} \kappa_{0,m}^{\widehat{\otimes} k}.$$

We show that for each test function φ in the space $(E)_\beta$, with $\frac{\max(0, m-2)}{m} \leq \beta < 1$, we have a unique representation

$$\varphi = \sum_{n=0}^{\infty} \langle : x^{\otimes n} :_{\kappa_{0,m}}, \varphi_n \rangle,$$

with $\varphi_n \in (E_{\mathbb{C}}^{\otimes n})_{sym}$ for all $n \in \mathbb{N}_0$. Indeed we can rewrite the Fourier-Gauss transform $\mathfrak{G}_{a,b}$ as a suitable second quantization operator $\{\Gamma_{\kappa_{0,r}}(bId)\}$. Using generalized Wick tensors it is very natural to find a larger class of one parameter transformation groups, see 6.15. As an application we deduce explicitly the regular one parameter groups corresponding to the infinitesimal generators $a\Delta_G + bN$.

In order to classify regularity we use and extend the result from [13, Theorem 5.6, p. 671]. There it is shown, that there exists a continuous homomorphism C from $(E)_\beta^*$ to $L((E)_\beta, (E)_\beta)$. We show furthermore that this homomorphism is even an isomorphism from $(E)_\beta^*$ onto $\text{Im}(C)$. We use this theorem to prove easily the differentiability of some one parameter transformation groups. Moreover we want to show that these transformation groups regular. In particular we show that every differentiable one-parameter subgroup of $GL(X)$, where X is a nuclear Fréchet space, has a regular generator, see 5.8.

The investigation of regular transformation groups in this manuscript can be compared with [3]. There the authors first construct a two-parameter transformation group G on the space of white noise test functions (E) in which the adjoints of Kuo's Fourier and Kuo's Fourier-Mehler transforms are included. They show that the group G is a two-dimensional complex Lie group whose infinitesimal generators are the Gross Laplacian Δ_G and the number operator N , and then find an explicit description of a differentiable one-parameter subgroup of G whose infinitesimal generator is $a\Delta_G + bN$, which is identical with the one in this article.

2. Preliminaries on White Noise Distribution Theory

Starting point of the white noise distribution theory is the Gel'fand triple

$$E \subset L^2(\mathbb{R}, dt) \subset E^*,$$

where E is a nuclear real countably Hilbert space which is densely embedded in the Hilbert space of square integrable functions with respect to the Lebesgue measure $L^2(\mathbb{R}, dt)$ and E^* its topological dual space. Via the Bochner-Minlos theorem, see e.g. [1], we obtain the Gaussian measure μ on E^* by its Fourier transform

$$\int_{E^*} \exp(i\langle x, \xi \rangle) d\mu(x) = \exp(-\frac{1}{2}|\xi|_0^2), \quad \xi \in E$$

where $|\cdot|_0$ denotes the Hilbertian norm on $L^2(\mathbb{R}, dt)$. The topology on E is induced by a positive self-adjoint operator A on the space of real-valued functions $H = L^2(\mathbb{R}, dt)$ with $\inf \sigma(A) > 1$ and Hilbert-Schmidt inverse A^{-1} . We set $\rho := \|A^{-1}\|_{OP}$ and $\delta := \|A^{-1}\|_{HS}$. Note that the complexification $E_{\mathbb{C}}$ are equipped with the norms $|\xi|_p := |A^p \xi|_0$ for $p \in \mathbb{R}$. We denote $H_{\mathbb{C}} := L^2(\mathbb{R}, \mathbb{C}, dt)$ furthermore $E_{\mathbb{C}, p} := \{\xi \in E_{\mathbb{C}}^* \mid |\xi|_p < \infty\}$ and $E_p^* := \{\xi \in E^* \mid |\xi|_p < \infty\}$, for $p \in \mathbb{R}$.

Now we consider the following Gel'fand triple of White Noise test and generalized functions.

$$(E)_{\beta} \subset (L^2) := L^2(E^*, \mu) \subset (E)_{\beta}^*, \quad 0 \leq \beta < 1$$

By the Wiener-Itô chaos decomposition theorem, see e.g. [7, 12, 10] we have the following unitary isomorphism between (L^2) and the Boson Fock space $\Gamma(H_{\mathbb{C}})$:

$$(L^2) \ni \Phi(x) = \sum_{n=0}^{\infty} \langle :x^{\otimes n} :, f_n \rangle \leftrightarrow (f_n) \sim \Phi \in \Gamma(H_{\mathbb{C}}), \quad f_n \in H_{\mathbb{C}}^{\hat{\otimes} n}, \quad (2.1)$$

where $:x^{\otimes n} :$ denotes the Wick ordering of $x^{\otimes n}$ and $H_{\mathbb{C}}^{\hat{\otimes} n}$ is the symmetric tensor product of order n of the complexification $H_{\mathbb{C}}$ of H . Moreover the (L^2) -norm of $\Phi \in (L^2)$ is given by

$$\|\Phi\|_0^2 = \sum_{n=0}^{\infty} n! |f_n|_0^2$$

The elements in $(E)_{\beta}$ are called white noise test functions, the elements in $(E)_{\beta}^*$ are called generalized white noise functions. We denote by $\langle\langle \cdot, \cdot \rangle\rangle$ the canonical \mathbb{C} -bilinear form on $(E)_{\beta}^* \times (E)_{\beta}$. For each $\Phi \in (E)_{\beta}^*$ there exists a unique sequence $(F_n)_{n=0}^{\infty}, F_n \in (E_{\mathbb{C}}^{\hat{\otimes} n})^*$ such that

$$\langle\langle \Phi, \varphi \rangle\rangle = \sum_{n=0}^{\infty} n! \langle F_n, f_n \rangle, \quad (f_n) \sim \varphi \in (E)_{\beta} \quad (2.2)$$

Thus we have, see e.g. [7, 12, 10]:

$$(E)_{\beta} \ni \Phi \sim (f_n),$$

if and only if for all $p \in \mathbb{R}$ we have $\|\Phi\|_{p,\beta} := \left(\sum_{n=0}^{\infty} (n!)^{1+\beta} |f_n|_p^2 \right)^{\frac{1}{2}} < \infty$. Moreover for its dual space we obtain

$$(E)_\beta^* \ni \Phi \sim (F_n), \quad (2.3)$$

if and only if there exists a $p \in \mathbb{R}$ such that $\|\Phi\|_{p,-\beta} := \left(\sum_{n=0}^{\infty} (n!)^{1-\beta} |F_n|_p^2 \right)^{\frac{1}{2}} < \infty$.

For $p \in \mathbb{R}$ we define

$$(E)_{p,\beta} := \left\{ \varphi \in (E)_\beta^* : \|\varphi\|_{p,\beta} < \infty \right\}$$

and

$$(E)_{p,-\beta} := \left\{ \varphi \in (E)_\beta^* : \|\varphi\|_{p,-\beta} < \infty \right\}.$$

It follows

$$(E)_\beta := \text{proj} \lim_{p \rightarrow \infty} (E)_{p,\beta}$$

and

$$(E)_\beta^* = \text{ind} \lim_{p \rightarrow -\infty} (E)_{p,-\beta}$$

Moreover $(E)_\beta$ is a nuclear (F)-space. In the following we use the abbreviation $(E) := (E)_0$. The exponential vector or Wick ordered exponential is defined by

$$\Phi_\xi(x) := \sum_{n=0}^{\infty} \frac{1}{n!} \langle : x^{\otimes n} :, \xi^{\otimes n} \rangle \quad \text{for } \xi \in E_{\mathbb{C}} \text{ and } x \in E^*. \quad (2.4)$$

For $y \in E_{\mathbb{C}}^*$ we use the same notation, which is only symbolic, and define $\Phi_y \in (E)_\beta^*$ by:

$$(E)_\beta \ni \psi \sim (f_n)_{n \in \mathbb{N}_0} : \langle \langle \Phi_y, \psi \rangle \rangle := \sum_{n=0}^{\infty} \langle y^{\otimes n}, f_n \rangle$$

Since $\Phi_\xi \in (E)_\beta$, for $\xi \in E_{\mathbb{C}}$ and $0 \leq \beta < 1$, we can define the so called S -transform of $\Psi \in (E)_\beta^*$ by

$$S(\Psi)(\xi) = \langle \langle \Phi_\xi, \Psi \rangle \rangle,$$

Moreover we call $S(\Psi)(0)$ the generalized expectation of $\Psi \in (E)_\beta^*$. The Wick product of $\Theta \in (E)_\beta^*$ and $\Psi \in (E)_\beta^*$ is defined by

$$\Psi \diamond \Theta := S^{-1}(S(\Psi) \cdot S(\Theta)) \in (E)_\beta^*,$$

see e.g. [7, 10, 12].

In the following we list some basic facts about integral kernel operators.

Definition 2.1. Let l, m be nonnegative integers. For each $\kappa_{l,m} \in (E_{\mathbb{C}}^{\otimes(l+m)})^*$ the integral kernel operator $\Xi_{l,m}(\kappa_{l,m}) \in L((E)_\beta, (E)_\beta^*)$ with kernel distribution $\kappa_{l,m}$ (see [12, Proposition 4.3.3, p. 82]) is defined by

$$\Xi_{l,m}(\kappa_{l,m})(\varphi) = \sum_{n=0}^{\infty} \frac{(n+m)!}{n!} \left\langle : x^{\otimes(l+n)} :, \kappa_{l,m} \otimes_m f_{n+m} \right\rangle, \quad \varphi \sim (f_n) \in (E)_\beta, \quad (2.5)$$

where $\kappa_{l,m} \otimes_m f_{n+m}$ is the right contraction (see [12, Section 3.4, p. 53 ff]).

Remark 2.2. Note that $\langle : x^{\otimes(l+n)} :, \kappa_{l,m} \otimes_m f_{n+m} \rangle = \langle : x^{\otimes(l+n)} :, \kappa_{l,m} \hat{\otimes}_m f_{n+m} \rangle$, because $: x^{\otimes(l+n)} :$ is symmetric and $\hat{\otimes}_m$ means the right symmetric contraction.

The correspondence

$$\kappa_{l,m} \leftrightarrow \Xi_{l,m}(\kappa_{l,m})$$

is one-to-one if

$$\kappa_{l,m} \in (E_{\mathbb{C}}^{\otimes(l+m)})_{sym(l,m)}^*$$

Furthermore it is known (see [10, Section 10.2, p. 123 ff]):

$$\Xi_{l,m}(\kappa_{l,m}) \in L((E)_{\beta}, (E)_{\beta}) \Leftrightarrow \kappa_{l,m} \in (E_{\mathbb{C}}^{\otimes l}) \otimes (E_{\mathbb{C}}^{\otimes m})^* \quad (2.6)$$

By a dual argument to [12, Theorem 4.3.9] it is known that an integral kernel operator $\Xi_{l,m}(\kappa_{l,m})$ is extendable to an operator in $L((E)_{\beta}^*, (E)_{\beta}^*)$ if and only if $\kappa_{l,m} \in (E_{\mathbb{C}}^{\otimes l})^* \otimes (E_{\mathbb{C}}^{\otimes m})$.

The operator symbol of $\Xi \in L((E)_{\beta}, (E)_{\beta}^*)$ is defined as $\hat{\Xi}(\xi, \eta) := \langle \Xi(\Phi_{\xi}), \Phi_{\eta} \rangle$ for $\xi, \eta \in E_{\mathbb{C}}$.

3. Convergence of Generalized Functions

Let X be a locally convex Hausdorff space and X^* its dual space with respect to the strong dual topology. Recall that $L(X, X)$ is equipped with the topology of bounded convergence, namely the locally convex topology is defined by the semi-norms:

$$\|T\|_{B,p} := \sup_{\xi \in B} |T\xi|_p, \quad T \in L(X, X) \quad (3.1)$$

where B runs over the bounded subsets of X and $|\cdot|_p$ is an arbitrary semi-norm on X . The topology on $L(X^*, X^*)$ is defined in the same way. We state the following proposition:

Proposition 3.1. *Let X be a reflexive (F)-space and X^* its dual space with the strong dual topology. Then the mapping $*$: $L(X, X) \rightarrow L(X^*, X^*)$, $T \mapsto T^*$ with $T^*(\xi) := \xi \circ T$ is a topological vector space isomorphism.*

Proof. For details see [6, 2.7 Satz, p. 84]. We present the main idea: Let M be a bounded subset of E^* and B be a bounded subset of X . Let $T \in L(X, X)$. Since X is reflexive, the topological isomorphism follows by

$$\sup_{\xi \in M} |T^*(\xi)|_{p_{B^{\circ}}} = \sup_{\xi \in M, b \in B} |\langle T^*\xi, b \rangle| = \sup_{b \in B} |Tb|_{p_M^{\circ}} \quad (3.2)$$

□

The following theorem can be found in e.g. [9],[7, Theorem 4.41, p. 127 f] and [10, Theorem 8.6, p. 86f].

Theorem 3.2. *Let $0 \leq \beta < 1$. For all $n \in \mathbb{N}$ let $\Psi_n \in (E)_{\beta}^*$ and $F_n = S(\Psi_n)$, where S denotes the S -transform. Then $(\Psi_n)_{n \in \mathbb{N}}$ converges strongly in $(E)_{\beta}^*$ if and only if the following conditions are satisfied:*

- (i) $\lim_{n \rightarrow \infty} F_n(\xi)$ exists for each $\xi \in E_{\mathbb{C}}$
- (ii) There exist nonnegative constants $C, K, p \geq 0$, independent of n , such that

$$|F_n(\xi)| \leq C \exp(K|\xi|_p^{\frac{2}{1-\beta}}), \quad \forall n \in \mathbb{N}, \xi \in E_{\mathbb{C}}. \quad (3.3)$$

As an example we consider the convergence of Wick exponentials.

Lemma 3.3. *Let $\phi \in (E)^*$ and $\phi \sim (\phi_0, \phi_1, \dots, \phi_m, 0, 0, 0, \dots)$ be its Wiener-Itô chaos decomposition. Then the expression*

$$\exp^\diamond(\phi) := \sum_{n=0}^{\infty} \frac{1}{n!} \phi^{\diamond n} \quad (3.4)$$

converges in $(E)_\beta^*$, where $1 > \beta \geq \frac{\max(0, m-2)}{m}$.

Proof. We show that the conditions from Theorem 3.2 are fulfilled. Without loss of let $m \geq 1$ and $\phi_m \neq 0$. Since $\phi \in (E)_\beta^*$ there exists $p > 0$ such that $\|\phi\|_{-p, \beta} < \infty$.

For $n \in \mathbb{N}_0$ and $\xi \in E_{\mathbb{C}}$ let $\Psi_n := \sum_{k=0}^n \frac{1}{k!} \phi^{\diamond k}$ and $F_n(\xi) := \sum_{k=0}^n \frac{1}{k!} \langle \langle \phi^{\diamond k}, \Phi_\xi \rangle \rangle$.

Condition (i) from Theorem 3.2 is fulfilled since

$$F_n(\xi) = \sum_{k=0}^n \frac{1}{k!} (S(\phi)(\xi))^k \rightarrow \exp(S(\phi)(\xi)).$$

It follows $|F_n(\xi)| \leq \exp(|S(\phi)(\xi)|)$. Since for $r \geq m$ we have

$$\begin{aligned} |S(\phi)(\xi)| &\leq |\phi_0| + \dots + |\langle \phi_m, \xi^{\otimes m} \rangle| \\ &\leq |\phi_0| + m \cdot \max_{1 \leq k \leq m} (|\phi_k|_{-p}) \cdot (1 + |\xi|_p^r) \end{aligned}$$

we obtain that condition (ii) from Theorem 3.2 is fulfilled for $\beta \geq 0$ and $r = \frac{2}{1-\beta}$. \square

The following is an immediate consequence of Lemma 3.3 and comparison of the S -transforms.

Corollary 3.4. *Let $\phi, \psi \in (E)^*$ with corresponding chaos decompositions $\phi = (\phi_0, \phi_1, \dots, \phi_m, 0, 0, 0, \dots)$ and $\psi = (\psi_0, \psi_1, \dots, \psi_m, 0, 0, 0, \dots)$. Then the formula*

$$\exp^\diamond(\phi + \psi) = \exp^\diamond(\phi) \diamond \exp^\diamond(\psi)$$

is valid in $(E)_\beta^*$, for all $1 > \beta \geq \frac{1}{m} \max(0, m-2)$.

4. Wick Multiplication and Convolution Operator

The Wick multiplication operator and its dual, often denoted as convolution operator are well known objects in white noise theory, see e.g. [13]. We give a proof, that $(E)_\beta^*$ with the Wick product may be considered as a commutative sub-algebra of $L((E)_\beta, (E)_\beta)$. For a similar approach see also [13].

Proposition 4.1.

(i) *Let $\varphi \in (E)_\beta^*$. The Wick multiplication operator M_φ , defined by*

$$M_\varphi(\psi) := \varphi \diamond \psi$$

is a well defined operator in $L((E)_\beta, (E)_\beta^)$.*

(ii) *Let $\varphi \in (E)_\beta^*$. The Wick multiplication operator \widetilde{M}_φ , defined by*

$$\widetilde{M}_\varphi(\psi) := \varphi \diamond \psi$$

is a well defined continuous extension of M_φ to an operator in $L((E)_\beta^, (E)_\beta^*)$.*

Proof. For (ii), see [10, Theorem 8.12., p. 92]. To prove (i) we know by [10, Theorem 8.12., see Remark, p.92], that for any $p \geq 0$ there exist suitable $\alpha > 0$, $c > 0$, such that

$$\begin{aligned} \|\varphi \diamond \psi\|_{-p-\alpha, -\beta} &\leq c \cdot \|\varphi\|_{-p, -\beta} \cdot \|\psi\|_{-p, -\beta} \\ &\leq c \cdot \|\varphi\|_{-p, -\beta} \cdot \|\psi\|_{p, \beta} \end{aligned}$$

Thus $\varphi \diamond (\cdot) \in L((E)_\beta, (E)_\beta^*)$. \square

Definition 4.2. The dual operator of the Wick operator \widetilde{M}_φ is called *convolution operator* and denoted by C_φ .

The next proposition gives the Fock expansion of the Wick multiplication operator.

Proposition 4.3. *Let $\varphi \in (E)_\beta^*$ and $\varphi \sim (\varphi_0, \varphi_1, \dots)$. Then M_φ , considered as $M_\varphi \in L((E)_\beta, (E)_\beta^*)$, has the following Fock expansion:*

$$M_\varphi = \sum_{n \in \mathbb{N}_0} \Xi_{n,0}(\varphi_n).$$

Proof. Let $\xi, \eta \in E_{\mathbb{C}}$. Then

$$\begin{aligned} \langle \langle M_\varphi \Phi_\xi, \Phi_\eta \rangle \rangle &= S(\varphi \diamond \Phi_\xi)(\eta) = S(\varphi)(\eta) \cdot S(\Phi_\xi)(\eta) \\ &= S(\varphi)(\eta) \cdot e^{\langle \xi, \eta \rangle} \end{aligned}$$

By [10, (10.21) p. 145] the Taylor expansion of M_φ is given by

$$S(\varphi)(\eta) = \sum_{n \in \mathbb{N}_0} \langle \varphi_n, \eta^{\otimes n} \otimes \xi^0 \rangle$$

completes the proof, compare also [12, Proposition 4.5.3, p.98-99]. \square

Proposition 4.4. *Let $\varphi \in (E)_\beta^*$ and $\varphi \sim (\varphi_0, \varphi_1, \dots)$. The convolution operator C_φ is in $L((E)_\beta, (E)_\beta)$ and has the Fock expansion*

$$C_\varphi = \sum_{n \in \mathbb{N}_0} \Xi_{0,n}(\varphi_n).$$

Proof. By definition we have $C_\varphi = \widetilde{M}_\varphi^* \in L((E)_\beta, (E)_\beta)$. Moreover $M_\varphi^*|(E)_\beta = C_\varphi$, since \widetilde{M}_φ is an extension of M_φ . Then, by Proposition 4.3 we have

$$C_\varphi = M_\varphi^* = \left(\sum_{n \in \mathbb{N}_0} \Xi_{n,0}(\varphi_n) \right)^* = \sum_{n \in \mathbb{N}_0} \Xi_{0,n}(\varphi_n).$$

\square

The following statement is an immediate consequence from the definition of integral kernel operators.

Corollary 4.5. *Let $\varphi \in (E)_\beta^*$ and $\varphi \sim (\varphi_0, \varphi_1, \dots)$. Then*

$$C_\varphi(\Phi_\xi) = [S(\varphi)(\xi)] \Phi_\xi, \quad \xi \in E_{\mathbb{C}} :$$

Lemma 4.6. *Let $\Xi \in L((E)_\beta, (E)_\beta^*)$ and $\Xi = \sum_{l,m=0}^{\infty} \Xi_{l,m}(\kappa_{l,m})$ be the corresponding unique representation as sum of integral kernel operators. Then:*

$$\Xi(\Phi_0) = \sum_{l=0}^{\infty} \langle : x^{\otimes l} :, \kappa_{l,0} \rangle \quad (4.1)$$

Proof. Consider the equation

$$\Xi_{l,m}(\kappa_{l,m})(\varphi) = \sum_{n=0}^{\infty} \frac{(n+m)!}{n!} \langle : x^{\otimes(l+n)} :, \kappa_{l,m} \hat{\otimes}_m f_{n+m} \rangle, \quad \varphi \sim (f_n) \in (E)_\beta, \quad (4.2)$$

then for $\Phi_0 \sim (f_n)$ with $f_0 = 1$ and $f_n = 0$ for all $n \geq 1$:

$$\Xi(\Phi_0) = \sum_{l=0}^{\infty} \langle : x^{\otimes l} :, \kappa_{l,0} \hat{\otimes}_0 1 \rangle$$

□

Theorem 4.7. *Let $0 \leq \beta < 1$. The mapping*

$$C : ((E)_\beta^*, +, \cdot, \diamond) \text{ to } (L((E)_\beta, (E)_\beta), +, \cdot, \diamond) \quad \varphi \mapsto \mathcal{C}_\varphi$$

is an injective continuous vector algebra homomorphism. In particular C is a topological isomorphism from $(E)_\beta^$ to $\text{Im}(C)$.*

Proof. Linearity is given by the definition of integral kernel operators. In order to prove homomorphy, let $\xi \in E_C$, $\varphi_1, \varphi_2 \in (E)_\beta^*$. Then by Corollary 4.5

$$\begin{aligned} C_{\varphi_1} \circ C_{\varphi_2}(\Phi_\xi) &= S(\varphi_1)(\xi) \cdot S(\varphi_2)(\xi) \cdot \Phi_\xi \\ &= S(\varphi_1 \diamond \varphi_2)(\xi) \cdot \Phi_\xi = C_{\varphi_1 \diamond \varphi_2}(\Phi_\xi) \end{aligned}$$

For injectivity let $\varphi \in (E)_\beta^*$ and $\mathcal{C}_\varphi = 0$. Then $\mathcal{C}_\varphi^* = 0$ and by 4.4 and 4.6 we obtain $\varphi = \mathcal{C}_\varphi^*(\Phi_0) = 0$. To prove continuity let $\varphi \in (E)_\beta^*$ and $\varphi := \sum_{n \in \mathbb{N}_0} \langle : x^{\otimes n} :, \varphi_n \rangle$ be the corresponding chaos decomposition. Now let $p \geq 0$. Moreover let $r > 0$, such that

$$(2^{\frac{1-\beta}{2}} \rho^r < 1),$$

where $\rho := \|A^{-1}\|_{OP}$. Choose $q > 0$ such that $\rho^{-q} \geq 2$. Then by [10, Theorem 10.5, p. 128] we have for all $\phi \in (E)_\beta$ and $m \in \mathbb{N}_0$

$$\begin{aligned} \|\Xi_{0,m}(\varphi_m)(\phi)\|_{p,\beta} &\leq (m!2^m)^{\frac{1-\beta}{2}} \cdot \|\varphi_m\|_{-(p+q+r)} \cdot \|\phi\|_{(p+q+r),\beta} \\ &\leq m!^{\frac{1-\beta}{2}} \|\varphi_m\|_{-(p+q)} (2^{\frac{1-\beta}{2}} \rho^r)^m \cdot \|\phi\|_{(p+q+r),\beta}. \end{aligned}$$

Now let $K := (\sum_{m \in \mathbb{N}_0} (2^{\frac{1-\beta}{2}} \rho^r)^{2m})^{\frac{1}{2}}$. Then by Cauchy-Schwartz we obtain

$$\begin{aligned} \sum_{m \in \mathbb{N}_0} \|\Xi_{0,m}(\varphi_m)(\phi)\|_{p,\beta} &\leq \sum_{m \in \mathbb{N}_0} m!^{\frac{1-\beta}{2}} \|\varphi_m\|_{-(p+q)} (2^{\frac{1-\beta}{2}} \rho^r)^m \cdot \|\phi\|_{(p+q+r),\beta} \\ &\leq K \|\varphi\|_{-(p+q),-\beta} \cdot \|\phi\|_{(p+q+r),\beta} \end{aligned}$$

Now let $M \subset (E)_\beta$ be bounded. Then $\sup_{\phi \in M} |\phi|_{p+q+r,\beta} \leq \text{const}(M, p, q, r) < \infty$ and

finally $\|\mathcal{C}_\varphi\|_{M, \{p,\beta\}} \leq K \text{const}(M, p, q, r) \cdot \|\varphi\|_{-(p+q), -\beta}$

To prove the last claim, we use Proposition 3.1 and and represent C^{-1} as composition of the two continuous mappings:

$$C^{-1} : \mathcal{C}_\varphi \xrightarrow{(\cdot)^*} \mathcal{C}_\varphi^* \xrightarrow[\text{evaluation}]{\text{point}} \mathcal{C}_\varphi^*(\Phi_0) = \varphi.$$

□

The following statement is a consequence from Lemma 3.3 and Theorem 4.7.

Corollary 4.8. *Let $\varphi \in (E)^*$ and $\varphi \sim (\varphi_0, \varphi_1, \dots, \varphi_m, 0, 0, 0, \dots)$ be it's chaos decomposition. Then the expression*

$$\exp(\mathcal{C}_\varphi) := \sum_{n=0}^{\infty} \frac{1}{n!} (\mathcal{C}_\varphi)^n \quad (4.3)$$

converges in $L((E)_\beta, (E)_\beta)$, where $1 > \beta \geq \max(0, \frac{m-2}{m})$.

5. Regular One Parameter Groups

Let X a nuclear (F)-space over \mathbb{C} and $(|\cdot|_n)_{n \in \mathbb{N}}$ be a family of Hilbertian semi-norms, with $|\cdot|_n \leq |\cdot|_{n+1}$ for all $n \in \mathbb{N}$, topologizing X , see e.g. [12, Proposition 1.2.2 and proposition 1.3.2]. For $x^* \in (X, |\cdot|_n)^*$ we define

$$|x^*|_{-n} := \sup_{|x|_n \leq 1} |\langle x^*, x \rangle|$$

By a Hahn-Banach argument, we have for all $x \in X$:

$$|x|_n = \sup \{ |\langle x^*, x \rangle| : x^* \in (X, |\cdot|_n)^* \wedge |x^*|_{-n} \leq 1 \}$$

for all semi-norms $|\cdot|_n$, $n \in \mathbb{N}$.

Definition 5.1. A family $\{\Omega_\theta\}_{\theta \in \mathbb{R}} \subset L(X)$ is called *differentiable* in $\theta_0 \in \mathbb{R}$, if

$$\lim_{\theta \rightarrow \theta_0} \frac{\Omega_\theta \phi - \Omega_{\theta_0} \phi}{\theta - \theta_0}$$

converges in X for any $\phi \in X$. In that case a linear operator Ω'_{θ_0} from X into itself is defined by

$$\Omega'_{\theta_0} \phi := \lim_{\theta \rightarrow \theta_0} \frac{\Omega_\theta \phi - \Omega_{\theta_0} \phi}{\theta - \theta_0}$$

$\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ is called *differentiable* if it is differentiable at each $\theta \in \mathbb{R}$. In the following we use the abbreviation:

$$\Omega' := \Omega'_0$$

Proposition 5.2. *Let $\theta_0 \in \mathbb{R}$ and $\{\Omega_\theta\}_{\theta \in \mathbb{R}} \subset L(X)$ be a family of operators which is differentiable in θ_0 . Then Ω'_{θ_0} is continuous, i.e. $\Omega'_{\theta_0} \in L(X)$. Moreover we have uniformly convergence on every compact (or equivalently, bounded) subset of X , i.e.:*

$$\lim_{\theta \rightarrow \theta_0} \sup_{\phi \in K} \left| \frac{\Omega_\theta \phi - \Omega_{\theta_0} \phi}{\theta - \theta_0} - \Omega'_{\theta_0} \phi \right|_n = 0$$

for any $n \in \mathbb{N}$ and any compact (or bounded) subset $K \subset X$.

Proof. First note that a subset of the nuclear space X is compact if and only if it is closed and bounded. The assertion follows by an application of the Banach-Steinhaus theorem. (For the uniform convergence on compact subsets see e.g. [14, 4.6 Theorem, p. 86].) \square

Definition 5.3. A family $\{\Omega_\theta\}_{\theta \in \mathbb{R}} \subset L(X)$ is called *regularly differentiable* in $\theta_0 \in \mathbb{R}$, if there exists $\Omega'_{\theta_0} \in L(X)$ such that for any $n \in \mathbb{N}$ there exists $m \in \mathbb{N}$ such that

$$\lim_{\theta \rightarrow \theta_0} \sup_{|\phi|_m \leq 1} \left| \frac{\Omega_\theta \phi - \Omega_{\theta_0} \phi}{\theta - \theta_0} - \Omega'_{\theta_0} \phi \right|_n = 0.$$

$\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ is called *regular* if it is regularly differentiable at each $\theta \in \mathbb{R}$.

Definition 5.4. Let $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ be a family of operators in $L(X)$ with

$$\forall \theta_1, \theta_2 \in \mathbb{R} : \Omega_{\theta_1 + \theta_2} = \Omega_{\theta_1} \circ \Omega_{\theta_2}, \quad \Omega_0 = \text{Id}$$

Then, as easily seen, $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ is a subgroup of $GL(X)$ and is called a *one-parameter subgroup* of $GL(X)$.

In the following we collect facts about one-parameter subgroups, for details see e.g. [12, Section 5.2].

Lemma 5.5. Let $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ be a one-parameter subgroup of $GL(X)$.

(i) Then $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ is differentiable at each $\theta \in \mathbb{R}$ if and only if $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ is differentiable at 0, i.e. there exists $\Omega' \in L(X)$ such that

$$\lim_{\theta \rightarrow 0} \left| \frac{\Omega_\theta \phi - \phi}{\theta} - \Omega' \phi \right|_n = 0.$$

for all $\phi \in X$ and $n \in \mathbb{N}$.

(ii) Let $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ be a differentiable subgroup of $GL(X)$. Then we have

a) Ω' is an element of $L(X)$, further unique and is called the *infinitesimal generator* of the differentiable one parameter subgroup $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ of $GL(X)$. Conversely a differentiable one parameter subgroup $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ of $GL(X)$ is uniquely defined by its infinitesimal generator, see [12, Proposition 5.2.2, p. 119]. If $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ is regular the infinitesimal generator Ω' is called a *regular generator*.

b) $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ is infinitely many differentiable at each $\theta \in \mathbb{R}$ and

$$\forall n \in \mathbb{N} : \frac{d^n}{d\theta^n} \Omega_\theta = (\Omega')^n \circ \Omega_\theta = \Omega_\theta \circ (\Omega')^n$$

c) $\mathbb{R} \rightarrow L(X)$, $\theta \mapsto \Omega_\theta$ is continuous. Note that $L(X)$ is equipped with the topology of bounded convergence, compare [12, Section 5.2, Eq. (5.27), p. 119].

Lemma 5.6. Let $T : \mathbb{R} \rightarrow L(X)$ be a continuous mapping. If K is a compact subset of \mathbb{R} , then $T(K)$ is equicontinuous.

Proof. $T(K)$ is compact, hence bounded, especially pointwisely bounded, by the definition of the topology of bounded convergence. The theorem of Banach and Steinhaus then completes the proof. \square

Proposition 5.7. *Let $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ be a differentiable one parameter subgroup of $GL(X)$.*

(i) $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ is regular if and only if for any $n \in \mathbb{N}$ there exists $m \in \mathbb{N}$ such that

$$\lim_{\theta \rightarrow 0} \sup_{|\phi|_m \leq 1} \left| \frac{\Omega_\theta \phi - \phi}{\theta} - \Omega' \phi \right|_n = 0.$$

(ii) For any compact (or bounded) subset $K \subset X$ we have

$$\lim_{\theta \rightarrow 0} \sup_{\phi \in K} \left| \frac{\Omega_\theta \phi - \phi}{\theta} - \Omega' \phi \right|_n = 0.$$

(iii) Let $\theta_0 > 0$. Then for any $n \in \mathbb{N}$ there exists $m \in \mathbb{N}$ and $K(n, m) > 0$, such that for all $\phi \in X$

$$\sup_{|\theta| \leq \theta_0, \theta \neq 0} \left| \frac{\Omega_\theta \phi - \phi}{\theta} \right|_n \leq K(n, m) |\phi|_m$$

Proof. The statement (i) is easily verified, moreover (ii) is an immediate consequence of Proposition 5.2. To prove statement (iii) we have by (ii) and Proposition 5.5 (ii) a), that the mapping

$$\mathbb{R} \longrightarrow L(X), \quad \theta \longmapsto \begin{cases} \frac{\Omega_\theta - \text{Id}}{\theta} & , \theta \neq 0 \\ \Omega' & , \theta = 0 \end{cases}$$

is continuous. Then the claim is obtained by Lemma 5.6. \square

Theorem 5.8 (Regularity). *Let $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ be a differentiable one-parameter subgroup of $GL(X)$. Then $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ is regular.*

Proof. Let $n \in \mathbb{N}$ and $\xi \in X$, $\eta \in (X, |\cdot|_n)^*$. We define for all $t \in \mathbb{R}$:

$$f(t) := \langle \eta, \Omega_t \xi \rangle$$

Then we have by Proposition 5.5

$$\begin{aligned} f'(t) &= \langle \eta, \Omega'_t \xi \rangle \\ f''(t) &= \langle \eta, (\Omega')^2 \Omega_t \xi \rangle \end{aligned}$$

Now let $\theta_0 > 0$ be fixed. Then by Proposition 5.5 (ii) c) and Lemma 5.6, $\{(\Omega')^2 \Omega_t\}_{|t| \leq \theta_0}$ is equicontinuous. Thus there exists $K = K(\theta_0, n, m, \Omega') > 0$ with $m \in \mathbb{N}_0$ and such that

$$\max_{|t| \leq \theta_0} |f''(t)| \leq K |\xi|_{n+m} |\eta|_{-n}$$

Let $\theta \in \mathbb{R}$ with $|\theta| \leq \theta_0$. By Taylor expansion we have

$$\begin{aligned} |f(\theta) - f(0) - \theta \cdot f'(0)| &\leq \frac{|\theta|^2}{2} \max_{|t| \leq \theta_0} |f''(t)| \\ &\leq \frac{|\theta|^2}{2} K |\xi|_{n+m} |\eta|_{-n} \end{aligned}$$

and for $\theta \neq 0$

$$\sup_{|\xi|_{n+m} \leq 1} \sup_{|\eta|_{-n} \leq 1} \left| \frac{\langle \eta, \Omega_\theta \xi \rangle - \langle \eta, \xi \rangle}{\theta} - \langle \eta, \Omega' \xi \rangle \right| \leq \frac{|\theta|}{2} K.$$

Hence

$$\sup_{|\xi|_{n+m} \leq 1} \left| \frac{\Omega_\theta \xi - \xi}{\theta} - \Omega' \xi \right|_n \leq \frac{|\theta|}{2} K$$

such that

$$\lim_{\theta \rightarrow 0} \sup_{|\xi|_{n+m} \leq 1} \left| \frac{\Omega_\theta \xi - \xi}{\theta} - \Omega' \xi \right|_n = 0.$$

□

Proposition 5.9. *Let $\{S_\theta\}_{\theta \in \mathbb{R}}$ be a differentiable one parameter subgroup of $GL(X)$ and $\{T_\theta\}_{\theta \in \mathbb{R}}$ be a family of operators in $L(X)$ which is differentiable in zero with $\lim_{\theta \rightarrow 0} T_\theta \phi = \phi$ for all $\phi \in X$. Then*

$$\frac{d}{d\theta} \Big|_{\theta=0} S_\theta \circ T_\theta = S' + T'$$

If $\{T_\theta\}_{\theta \in \mathbb{R}}$ is regularly differentiable in zero, then

$$\{S_\theta \circ T_\theta\}_{\theta \in \mathbb{R}}$$

is also regularly differentiable in zero.

Proof. Let $n \in \mathbb{N}$, $\phi \in X$. Then $\{S_\theta \mid |\theta| \leq 1\}$ is equicontinuous by Proposition 5.5, (ii) c). Then there exist a semi-norm $|\cdot|_m$, with $m \in \mathbb{N}$ and $m \geq n$, and a constant $K > 0$ such that $|S_\theta(\phi)|_n \leq K \cdot |\phi|_m$ for all $\phi \in X$ and $\theta \in \mathbb{R}$ with $|\theta| \leq 1$. Now let $\phi \in X$ be arbitrarily chosen. For $\theta \in \mathbb{R}$ with $|\theta| \leq 1$ we have:

$$\begin{aligned} & \left| \frac{S_\theta \circ T_\theta(\phi) - \phi}{\theta} - S' \phi - T' \phi \right|_n = \left| \frac{S_\theta \circ (T_\theta(\phi) - \phi)}{\theta} + \frac{S_\theta(\phi) - \phi}{\theta} - S' \phi - T' \phi \right|_n \\ & \leq \left| \frac{S_\theta \circ (T_\theta(\phi) - \phi)}{\theta} - T' \phi \right|_n + \left| \frac{S_\theta(\phi) - \phi}{\theta} - S' \phi \right|_n \\ & \leq K \cdot \left| \frac{(T_\theta(\phi) - \phi)}{\theta} - S_{-\theta} T' \phi \right|_m + \left| \frac{S_\theta(\phi) - \phi}{\theta} - S' \phi \right|_n \\ & \leq K \cdot \left| \frac{(T_\theta(\phi) - \phi)}{\theta} - T' \phi \right|_m + K \cdot |S_{-\theta}(T' \phi) - (T' \phi)|_m + \left| \frac{S_\theta(\phi) - \phi}{\theta} - S' \phi \right|_n, \end{aligned}$$

where last inequality follows by Theorem 5.8 and the continuity of T' . □

We use the following notation, due to [12, Eq. (4.66), p. 106]

Definition 5.10.

$$\begin{cases} \gamma_n(T) & := \sum_{k=0}^{n-1} \text{Id}^{\otimes k} \otimes T \otimes \text{Id}^{\otimes (n-1-k)}, \quad n \geq 1 \\ \gamma_0(T) & := 0 \end{cases}$$

Now let $T \in L(E_{\mathbb{C}})$. We recall the definition of the second quantization operator of T , denoted by $\Gamma(T)$ and $d\Gamma(T)$, the differential second quantization operator. Suppose $\phi \in (E)_{\beta}$ is given as

$$\phi(x) = \sum_{n=0}^{\infty} \langle : x^{\otimes n} :, f_n \rangle, \quad x \in E^*, \quad f_n \in (E_{\mathbb{C}}^{\otimes n})_{sym}$$

as usual. We put

$$\Gamma(T)\phi(x) := \sum_{n=0}^{\infty} \langle : x^{\otimes n} :, T^{\otimes n} f_n \rangle$$

and

$$d\Gamma(T)\phi(x) := \sum_{n=0}^{\infty} \langle : x^{\otimes n} :, \gamma_n(T) f_n \rangle$$

We have $\Gamma(T), d\Gamma(T) \in L((E)_\beta)$, for all $0 \leq \beta < 1$.

Lemma 5.11. *Let $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ be a differentiable one parameter subgroup of $GL(X)$ with infinitesimal generator Ω' . Then, for all $n \in \mathbb{N}$, $\{\Omega_\theta^{\otimes n}\}_{\theta \in \mathbb{R}}$ is a differentiable one parameter subgroup of $GL(X^{\otimes n})$ with*

$$\frac{d}{d\theta} \Big|_{\theta=0} \Omega_\theta^{\otimes n} = \gamma_n(\Omega')$$

Proof. We show the case $n = 2$. The general case is similar. It holds

$$\Omega_\theta \otimes \Omega_\theta = (\Omega_\theta \otimes \text{Id}) \circ (\text{Id} \otimes \Omega_\theta)$$

Now apply Proposition 5.9 □

Proposition 5.12. *Let $0 \leq \beta < 1$. Further let $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ be a differentiable one parameter subgroup of $GL(E_{\mathbb{C}})$ with infinitesimal generator Ω' . Then $\Gamma(\Omega_\theta)_{\theta \in \mathbb{R}}$ is a differentiable one-parameter subgroup of $GL((E)_\beta)$ with infinitesimal generator $d\Gamma(\Omega')$.*

Proof. Note that for $0 \leq \beta < 1$, the sequence $(\alpha(n))_{n \in \mathbb{N}_0}$ with $\alpha(n) := n!^\beta$ fullfills the conditions of [5, Theorem 4.2, p.696], where a detailed proof, based on the characterization theorem, is given. On the other hand the expected result is easily seen by Lemma 5.11. □

For a similar statement like Proposition 5.12, see [12, 5.4.5, p. 130-131].

Proposition 5.13. *Let $\{\Omega_\theta\}_{\theta \in \mathbb{R}}$ be a differentiable one parameter subgroup of $GL(E_{\mathbb{C}})$. Furthermore let $r \in \mathbb{N}$, $1 > \beta \geq \frac{\max(0, r-2)}{r}$ and $\kappa_{0,r} \in (E_{\mathbb{C}}^{\otimes r})_{sym}^*$.*

Then $\{\exp(\Xi_{0,r}((\Omega_\theta^{\otimes r})^ \kappa_{0,r}))\}_{\theta \in \mathbb{R}} \subset GL((E)_\beta)$ is an in zero differentiable family of operators with*

$$\frac{d}{d\theta} \Big|_{\theta=0} \exp(\Xi_{0,r}((\Omega_\theta^{\otimes r})^* \kappa_{0,r})) = \Xi_{0,r}((\gamma_r(\Omega'))^* \kappa_{0,r}) \circ \exp(\Xi_{0,r}((\kappa_{0,r})).$$

Proof. First, because \mathbb{R} is a metric space, it is enough to consider sequential convergence, i.e. the limit process for any arbitrary sequence $(\theta_n)_{n \in \mathbb{N}}$ in \mathbb{R} with $\lim_{n \rightarrow \infty} \theta_n = 0$. Consider the sequence $\{\exp^\diamond \langle : x^{\otimes r} :, (\Omega_{\theta_n}^{\otimes r})^* \kappa_{0,r} \rangle\}_{n \in \mathbb{N}}$. Then the limit process will be transferred to $L((E)_\beta, ((E)_\beta))$ by continuity, using Theorem 4.7. Now let $(\theta_n)_{n \in \mathbb{N}}$ be an arbitrary sequence in \mathbb{R} with $\lim_{n \rightarrow \infty} \theta_n = 0$ and $\theta_n \neq 0$ for all $n \in \mathbb{N}$. Further define

$$\varphi_n := \frac{\exp^\diamond \langle : x^{\otimes r} :, (\Omega_{\theta_n}^{\otimes r})^* \kappa_{0,r} \rangle - \exp^\diamond \langle : x^{\otimes r} :, \kappa_{0,r} \rangle}{\theta_n}$$

for all $n \in \mathbb{N}$. Then, by Lemma 3.3, we have $\varphi_n \in (E)_\beta^*$ for all $n \in \mathbb{N}$. We verify the conditions of Theorem 3.2.

Let $\xi \in E_{\mathbb{C}}$. Then for all $n \in \mathbb{N}$

$$S(\varphi_n)(\xi) = \frac{\exp(\langle \kappa_{0,r}, (\Omega_{\theta_n}^{\otimes r}) \xi^{\otimes r} \rangle) - \exp(\langle \kappa_{0,r}, \xi^{\otimes r} \rangle)}{\theta_n}$$

First note that for all $n \in \mathbb{N}$ the S -transform $S(\varphi_n)$ is entire holomorphic. By Lemma 5.11 and Proposition 5.5 we obtain $\{\Omega_\theta^{\otimes r}\}_{\theta \in \mathbb{R}}$ as a regular one-parameter subgroup of $(E_{\mathbb{C}}^{\otimes r})_{sym}$ with

$$\frac{d}{d\theta} \Omega_\theta^{\otimes r} = \gamma_r(\Omega') \Omega_\theta^{\otimes r}.$$

Hence the function $\theta \mapsto \langle \kappa_{0,r}, (\Omega_\theta^{\otimes r}) \xi^{\otimes r} \rangle$ is infinitely often differentiable on \mathbb{R} and the same holds for $\theta \mapsto \exp(\langle \kappa_{0,r}, (\Omega_\theta^{\otimes r}) \xi^{\otimes r} \rangle)$ as composition of two infinitely many differentiable functions with

$$\frac{d}{d\theta} \exp(\langle \kappa_{0,r}, (\Omega_\theta^{\otimes r}) \xi^{\otimes r} \rangle) = \langle \kappa_{0,r}, \gamma_r(\Omega') \Omega_\theta^{\otimes r} \xi^{\otimes r} \rangle \cdot \exp(\langle \kappa_{0,r}, (\Omega_\theta^{\otimes r}) \xi^{\otimes r} \rangle)$$

Hence $\lim_{n \rightarrow \infty} (S(\varphi_n)(\xi))$ exists and we have:

$$\lim_{n \rightarrow \infty} S(\varphi_n)(\xi) = \langle \kappa_{0,r}, \gamma_r(\Omega') \xi^{\otimes r} \rangle \cdot \exp(\langle \kappa_{0,r}, \xi^{\otimes r} \rangle)$$

by the chain rule and Proposition 5.12.

For the growth estimate let $p \geq 0$. Without loss of generality let $|\theta_n| < 1$ for all $n \in \mathbb{N}$. Then, since $\{\Omega_\theta\}_{|\theta| \leq 1}$ and $\{\Omega' \Omega_\theta\}_{|\theta| \leq 1}$ are compact by 5.6, there exists a $q \geq 0$ such that, $\forall \theta \in \mathbb{R}$, $|\theta| \leq 1$ we have $|\Omega_\theta(\xi)|_p \leq |\xi|_{p+q}$ and $|\Omega' \Omega_\theta(\xi)|_p \leq |\xi|_{p+q}$. Then, by the mean value theorem and by Proposition 5.5 (ii) b), it follows for each $n \in \mathbb{N}$:

$$\begin{aligned} |S(\varphi_n)(\xi)| &\leq \sup_{|\theta| \leq 1} \left| \langle \kappa_{0,r}, \gamma_r(\Omega') \Omega_\theta^{\otimes r} \xi^{\otimes r} \rangle \cdot \exp(\langle \kappa_{0,r}, (\Omega_\theta^{\otimes r}) \xi^{\otimes r} \rangle) \right| \\ &\leq |\kappa_{0,r}|_{-p} \cdot r \cdot |\xi|_{p+q}^r \cdot \exp(|\kappa_{0,r}|_{-p} \cdot |\xi|_{p+q}^r) \\ &\leq |\kappa_{0,r}|_{-p} \cdot r(r!) \cdot \exp((1 + |\kappa_{0,r}|_{-p}) \cdot |\xi|_{p+q}^r) \\ &\leq |\kappa_{0,r}|_{-p} \cdot r(r!) \cdot \exp((1 + |\kappa_{0,r}|_{-p}) \cdot (1 + |\xi|_{p+q}^{\frac{2}{1-\beta}})), \end{aligned}$$

for all $1 > \beta \geq \frac{\max(0, r-2)}{r}$. The claim is a consequence of Theorem 4.7. \square

The same idea as in the proof of Proposition 5.13 combined with Theorem 5.8 leads to the following result.

Proposition 5.14. *Let $r \in \mathbb{N}$, $1 > \beta \geq \frac{1}{r} \max(0, r-2)$. Further let $\varphi \in (E)_\beta^*$ with $\varphi \sim (\varphi_0, \dots, \varphi_r, 0, 0, 0, \dots)$. Define $\Xi_\theta := \exp(\theta \mathcal{C}_\varphi)$ for $\theta \in \mathbb{R}$. Then $\{\Xi_\theta : \theta \in \mathbb{R}\}$ is a regular one parameter subgroup of $GL((E)_\beta)$ with infinitesimal generator \mathcal{C}_φ .*

Example 5.15. Let $y \in E^*$. As simple example consider the operator $\Xi_{0,1}(y)$. Recall that $D_y = \Xi_{0,1}(y)$ and the translation operator $T_y = \exp(D_y)$.

Let $z \in \mathbb{C}$. From $z\Delta_G = \Xi_{0,2}(z\tau)$, where τ is the trace operator, we conclude that $\exp(z\Delta_G) \in L((E)_\beta, (E)_\beta)$, for all $0 \leq \beta < 1$.

6. Generalized Wick Tensors and an Application

Our goal in this section is to rewrite Fourier-Gauss transforms as second quantization operators. This will be accomplished by suitable basis-transformations which will be explicitly calculated. For this purpose we introduce generalized Wick tensors. In this context Fourier-Gauss transforms appear as second quantization operators of the form $\{\Gamma_{\kappa_0, r}(\Omega_\theta)\}$. As an application we deduce explicitly the regular one parameter groups corresponding to the infinitesimal generators $a\Delta_G + bN$.

First we repeat some definitions. Let $a, b \in \mathbb{C}$, $0 \leq \beta < 1$. The Fourier-Gauss transform $\mathfrak{G}_{a,b}(\varphi)$ of $\varphi \in (E)_\beta$ is defined to be the function

$$\mathfrak{G}_{a,b}(\varphi)(y) = \int_{E^*} \varphi(ax + by) d\mu(x)$$

The Fourier-Gauss transform is in $L((E)_\beta, (E)_\beta)$ and the operator symbol is given by

$$\widehat{\mathfrak{G}}_{a,b}(\xi, \eta) = \exp \left[\frac{1}{2} (a^2 + b^2 - 1) \langle \xi, \xi \rangle + b \langle \xi, \eta \rangle \right], \quad \text{for all } \xi, \eta \in E_{\mathbb{C}}$$

see e.g. [10, Theorem 11.29, p. 168-169]. Hence

$$\mathfrak{G}_{a,b} = \Gamma(b \text{ Id}) \circ \exp \left(\frac{1}{2} (a^2 + b^2 - 1) \Delta_G \right)$$

By [10, Lemma 11.22, p. 163] the operator symbol of the Fourier transform is given by

$$\widehat{\mathfrak{F}}(\xi, \eta) = \exp(-i \langle \xi, \eta \rangle - \frac{1}{2} \langle \eta, \eta \rangle), \quad \text{for all } \xi, \eta \in E_{\mathbb{C}}.$$

Consequently

$$\mathfrak{F} = \exp \left(-\frac{1}{2} \Delta_G \right)^* \circ \Gamma(-i \text{ Id})$$

Moreover the operator symbol of the Fourier-Mehler transform is given by

$$\widehat{\mathfrak{F}}_\theta(\xi, \eta) = \exp(e^{i\theta} \langle \xi, \eta \rangle + \frac{i}{2} e^{i\theta} \sin \theta \langle \eta, \eta \rangle), \quad \text{for all } \xi, \eta \in E_{\mathbb{C}}, \theta \in \mathbb{R},$$

see e.g. [10, 11., p. 180]. Thus the Fourier-Mehler transform is given by the formula

$$\mathfrak{F}_\theta = \left[\exp \left(\frac{i}{2} e^{i\theta} \sin \theta \Delta_G \right) \right]^* \circ \Gamma(e^{i\theta} \text{ Id}).$$

In the following let \mathcal{G}_θ denote the adjoint of the Fourier-Mehler transform \mathfrak{F}_θ .

Finally by [12, Proposition 4.6.9, p. 105] the operator symbol of the scaling operator is given by

$$\widehat{S}_\lambda(\xi, \eta) = \exp \left((\lambda^2 - 1) \langle \xi, \xi \rangle / 2 + \lambda \langle \xi, \eta \rangle \right), \quad \text{for all } \xi, \eta \in E_{\mathbb{C}}, \lambda \in \mathbb{C}.$$

and consequently

$$S_\lambda = \Gamma(\lambda \text{ Id}) \circ \exp \left(\frac{\lambda^2 - 1}{2} \Delta_G \right)$$

Definition 6.1. Let $m \in \mathbb{N}$ and $\kappa_{0,m} \in ((E_{\mathbb{C}})^{\otimes m})_{sym}^*$. For $x \in E^*$ we define the *renormalized tensor power* $:x^{\otimes n} :_{\kappa_{0,m}}$ as follows:

$$:x^{\otimes n} :_{\kappa_{0,m}} = \sum_{k=0}^{\lfloor \frac{n}{m} \rfloor} \frac{n!}{(n-mk)!k!} \cdot \left(-\frac{1}{2}\right)^k \cdot x^{\otimes(n-mk)} \hat{\otimes} (\kappa_{0,m})^{\hat{\otimes} k}$$

For the usual tensor power $x^{\otimes n}$, $x \in E^*$ it holds the following relation:, see [12, Corollary 2.2.4, p. 25]

$$x^{\otimes n} = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n!}{(n-2k)!k!} \cdot \left(\frac{1}{2}\right)^k \cdot :x^{\otimes(n-2k)} :_{\tau} \hat{\otimes} (\tau)^{\hat{\otimes} k}$$

As a simple example we have for $x \in E$ the relation $:x^{\otimes n} : = :x^{\otimes n} :_{\tau}$. More usually is the abbreviation $:x^{\otimes n} :_{\sigma^2} \stackrel{def}{=} :x^{\otimes n} :_{\sigma^2 \tau}$.

Note that $\kappa_{0,m} = 0$ is permitted, e.g. $:x^{\otimes n} :_0 = x^{\otimes n}$. But we don't permit $m = 0$.

There exists Θ in $L((E), (E))$ defined by $\Theta(\exp(\langle \cdot, \xi \rangle)) := \Phi_{\xi}$, see e.g. [10, Theorem 6.2]. Since $\int_{E^*} \Phi_{\xi} d\mu = 1$ we call Θ the renormalization operator.

Proposition 6.2. (i) $\Theta = \exp(-\frac{1}{2}\Delta_G)$

(ii) For all $f_m \in (E_{\mathbb{C}}^{\otimes m})_{sym}$ we have:

$$\Theta(\langle x^{\otimes m}, f_m \rangle) = \langle :x^{\otimes m} :, f_m \rangle$$

Proof. By Corollary 4.5 we have for all $\xi \in E_{\mathbb{C}}$:

$$\begin{aligned} \exp(\frac{1}{2}\Delta_G)\Phi_{\xi} &= \exp(\frac{1}{2}\langle \xi, \xi \rangle)\Phi_{\xi} \\ &= \exp(\frac{1}{2}\langle \xi, \xi \rangle) \exp(-\frac{1}{2}\langle \xi, \xi \rangle) e^{\langle \cdot, \xi \rangle} \\ &= e^{\langle \cdot, \xi \rangle} \end{aligned}$$

By Proposition 5.14 Θ is invertible and the first statement is proved. To proof the second, let $m \in \mathbb{N}_0$. Note that for $m < 2n$ we have

$$\Xi_{0,2n}(\tau^{\otimes n})(\langle :x^{\otimes m} :, f_m \rangle) = 0,$$

further $\Theta^{-1} = \exp(\frac{1}{2}\Delta_G) = \sum_{n=0}^{\infty} \frac{1}{2^n n!} \Xi_{0,2n}(\tau^{\otimes n})$. Let $m \geq 2n$ and $\delta_{i,j}$ be the Kronecker symbol. Then by [12, Proposition 4.3.3, Eq. (4.23), p. 82] it holds:

$$\begin{aligned} \Xi_{0,2n}(\tau^{\otimes n})(\langle :x^{\otimes m} :, f_m \rangle) &= \sum_{k=0}^{\infty} \frac{(k+2n)!}{k!} (\langle :x^{\otimes k} :, \tau^{\otimes n} \otimes_{2n} \delta_{k+2n,m} \cdot f_m \rangle) \\ &= \frac{m!}{(m-2n)!} \langle :x^{\otimes(m-2n)} :, \tau^{\otimes n} \otimes_{2n} f_m \rangle \\ &= \frac{m!}{(m-2n)!} \langle :x^{\otimes(m-2n)} : \otimes \tau^{\otimes n}, f_m \rangle \\ &= \frac{m!}{(m-2n)!} \langle :x^{\otimes(m-2n)} : \hat{\otimes} \tau^{\hat{\otimes} n}, f_m \rangle \end{aligned}$$

where the last equation is due to the symmetricity of f_m . Then

$$\begin{aligned} \exp\left(\frac{1}{2}\Delta_G\right)(\langle : x^{\otimes m} :, f_m \rangle) &= \left\langle \sum_{n=0}^{\lfloor \frac{m}{2} \rfloor} \frac{m!}{(m-2n)!n!2^n} : x^{\otimes m-2n} : \hat{\otimes} \tau^{\otimes n}, f_m \right\rangle \\ &= \langle x^{\otimes m}, f_m \rangle, \end{aligned}$$

compare also [12, Corollary 2.2.4]. \square

Note that $e^{(\cdot, \xi)} \in (E)$ since $\Phi_\xi \in (E)$.

Corollary 6.3. *Let $\xi \in E_{\mathbb{C}}$. Then the series $e^{(\cdot, \xi)} = \sum_{n=0}^{\infty} \frac{1}{n!} \langle \cdot, \xi \rangle^n$ converges in (E) .*

Proof. Let $\xi \in E_{\mathbb{C}}$. Since

$$\Phi_\xi = \sum_{n=0}^{\infty} \frac{1}{n!} \langle : x^{\otimes n} :, \xi^{\otimes n} \rangle$$

converges in (E) also

$$\exp\left(\frac{1}{2}\Delta_G\right)\Phi_\xi = \sum_{n=0}^{\infty} \frac{1}{n!} \langle x^{\otimes n}, \xi^{\otimes n} \rangle$$

converges in (E) . \square

The following proposition generalizes Proposition 6.2.

Proposition 6.4. *Let $r \in \mathbb{N}$ and $\kappa_{0,r} \in ((E_{\mathbb{C}})^{\otimes r})_{sym}^*$. For all $f_m \in (E_{\mathbb{C}}^{\otimes m})_{sym}$ we have:*

$$\exp\left(-\frac{1}{2}\Xi_{0,r}(\kappa_{0,r})\right)(\langle x^{\otimes m}, f_m \rangle) = \langle : x^{\otimes m} :_{\kappa_{0,r}}, f_m \rangle$$

Proof. Let $m \in \mathbb{N}_0$. Note that for $m < rn$ we have

$$\Xi_{0,rn}(\kappa_{0,r}^{\otimes n})(\langle : x^{\otimes m} :, f_m \rangle) = 0,$$

further $\exp\left(-\frac{1}{2}\Xi_{0,r}(\kappa_{0,r})\right) = \sum_{n=0}^{\infty} \frac{1}{n!} \left(-\frac{1}{2}\right)^n \Xi_{0,rn}(\kappa_{0,r}^{\otimes n})$. Let $m \geq rn$ and $\delta_{i,j}$ be the Kronecker symbol. Then by Theorem 4.7 and Proposition 6.2, using [12, Proposition 4.3.3, Eq. (4.23), p. 82], we have:

$$\begin{aligned} \Xi_{0,rn}(\kappa_{0,r}^{\otimes n})(\langle x^{\otimes m}, f_m \rangle) &= \exp\left(\frac{1}{2}\Delta_G\right) \circ \Xi_{0,rn}(\kappa_{0,r}^{\otimes n})(\langle : x^{\otimes m} :, f_m \rangle) \\ &= \exp\left(\frac{1}{2}\Delta_G\right) \sum_{k=0}^{\infty} \frac{(k+rn)!}{k!} (\langle : x^{\otimes k} :, \kappa_{0,r}^{\otimes n} \otimes_{rn} \delta_{k+rn,m} \cdot f_m \rangle) \\ &= \frac{m!}{(m-rn)!} \langle x^{\otimes m-rn}, \kappa_{0,r}^{\otimes n} \otimes_{rn} f_m \rangle \\ &= \frac{m!}{(m-rn)!} \langle x^{\otimes m-rn} \otimes \kappa_{0,r}^{\otimes n}, f_m \rangle = \frac{m!}{(m-rn)!} \langle x^{\otimes m-rn} \hat{\otimes} \kappa_{0,r}^{\otimes n}, f_m \rangle \end{aligned}$$

where the last equation follows because f_m is symmetric.

Then by the above definition we have:

$$\begin{aligned} \exp\left(-\frac{1}{2}\Xi_{0,r}(\kappa_{0,r})\right)\langle x^{\otimes m}, f_m \rangle &= \left\langle \sum_{n=0}^{\lfloor \frac{m}{r} \rfloor} \frac{m!}{(m-rn)!n!} \left(-\frac{1}{2}\right)^n x^{\otimes m-rn} \hat{\otimes} \kappa_{0,r}^{\hat{\otimes} n}, f_m \right\rangle \\ &= \langle :x^{\otimes m} :_{\kappa_{0,r}}, f_m \rangle \end{aligned}$$

□

Notation 6.5. In the following we use $0 \cdot \tau$ in order to express that we consider $0 \in (E^{\otimes 2})^*$.

Theorem 6.6 (Representation theorem). *Let $r \in \mathbb{N}$, $1 > \beta \geq \frac{\max(0, r-2)}{r}$ and $\kappa_{0,r} \in ((E_{\mathbb{C}})^{\otimes r})_{sym}^*$. Then each $(\varphi_n)_{n \in \mathbb{N}_0} \sim \varphi \in (E)_{\beta}$ has a unique decomposition*

$$\varphi(x) = \sum_{n=0}^{\infty} \langle :x^{\otimes n} :_{\kappa_{0,r}}, (\psi_n)_{\kappa_{0,r}} \rangle, \quad \text{for all } x \in E^*$$

with $(\psi_n)_{\kappa_{0,r}} \in (E_{\mathbb{C}}^{\otimes n})_{sym}$, which we denote as $\kappa_{0,r}$ -representation of φ .

Proof. First note that φ has a unique Wiener-Itô chaos decomposition $(\varphi_n)_{n \in \mathbb{N}_0}$, see e.g. [12, Theorem 3.1.5]. Then the existence and uniqueness of the above representation follows by the bijectivity of $\exp(-\frac{1}{2}\Delta_G) \circ \exp(\frac{1}{2}\Xi_{0,r}(\kappa_{0,r}))$. Recall Corollary 4.8 and the following chain of mappings:

$$\langle :x^{\otimes n} :_{\kappa_{0,r}}, \varphi_n \rangle \xrightarrow{\exp(\frac{1}{2}\Xi_{0,r}(\kappa_{0,r}))} \langle x^{\otimes n}, \varphi_n \rangle \xrightarrow{\exp(-\frac{1}{2}\Delta_G)} \langle :x^{\otimes n} :, \varphi_n \rangle$$

□

Remark 6.7. The S -transform of $\varphi \in (E)$ is a restriction of the 0τ -representation of $\exp(\frac{1}{2}\Delta_G)\varphi$ from E^* to E .

Note that we do not claim that the above decomposition is an orthogonal decomposition with respect to the measure μ , like the chaos decomposition. We claim only the uniqueness.

Definition 6.8. Let $r \in \mathbb{N}$, $1 > \beta \geq \frac{\max(0, r-2)}{r}$, $\kappa_{0,r} \in (E_{\mathbb{C}}^{\otimes r})_{sym}^*$. For $T \in L((E)_{\beta}, (E)_{\beta})$, we define:

$$T_{\kappa_{0,r}} := \exp\left(\frac{1}{2}(\Delta_G - \Xi_{0,r}(\kappa_{0,r}))\right) \circ T \circ \exp\left(-\frac{1}{2}(\Delta_G - \Xi_{0,r}(\kappa_{0,r}))\right)$$

$T_{\kappa_{0,r}}$ is called the *renormalization* of T corresponding to $\kappa_{0,r}$. Obviously $T_{\kappa_{0,r}} \in L((E)_{\beta}, (E)_{\beta})$. For $\Omega \in L((E)_{\beta})$ we abbreviate

$$\begin{aligned} \Gamma_{\kappa_{0,r}}(\Omega) &:= (\Gamma(\Omega))_{\kappa_{0,r}}, \\ d\Gamma_{\kappa_{0,r}}(\Omega) &:= (d\Gamma(\Omega))_{\kappa_{0,r}}. \end{aligned}$$

Remark 6.9. It is clear, that $T_{\kappa_{0,r}}$ acts formally on white noise test functions in $\kappa_{0,r}$ -representation like T on white noise test functions in the standard representation as Boson Fock space. We precise this statement by the following proposition.

Proposition 6.10. *Let $r \in \mathbb{N}$, $1 > \beta \geq \frac{\max(0, r-2)}{r}$, $\kappa_{0,r} \in (E_{\mathbb{C}}^{\otimes r})_{sym}^*$. For $T \in L((E)_{\beta}, (E)_{\beta})$ and $\varphi \in (E)_{\beta}$ with $\varphi = \sum_{n=0}^{\infty} \langle : x^{\otimes n} :, f_n \rangle$, we use the notation $T(\varphi) = \sum_{n=0}^{\infty} \langle : x^{\otimes n} :, (f_n)_T \rangle$. Then it follows:*

$$T_{\kappa_{0,r}} \left(\sum_{n=0}^{\infty} \langle : x^{\otimes n} :_{\kappa_{0,r}}, f_n \rangle \right) = \sum_{n=0}^{\infty} \langle : x^{\otimes n} :_{\kappa_{0,r}}, (f_n)_T \rangle$$

Proof. The expression $\sum_{n=0}^{\infty} \langle : x^{\otimes n} :_{\kappa_{0,r}}, f_n \rangle$ is well defined because

$$\sum_{n=0}^{\infty} \langle : x^{\otimes n} :_{\kappa_{0,r}}, f_n \rangle = \exp\left(-\frac{1}{2}\Xi_{0,r}(\kappa_{0,r})\right) \circ \exp\left(\frac{1}{2}\Delta_G\right) \left(\sum_{n=0}^{\infty} \langle : x^{\otimes n} :, f_n \rangle \right).$$

By Corollary 3.4 and Theorem 4.7 we have

$$\exp\left(-\frac{1}{2}(\Delta_G - \Xi_{0,r}(\kappa_{0,r}))\right) = \exp\left(-\frac{1}{2}\Delta_G\right) \circ \exp\left(\frac{1}{2}\Xi_{0,r}(\kappa_{0,r})\right).$$

Then using Proposition 6.2 and Proposition 6.4, the claim follows with the same idea as in the proof of Theorem 6.6. \square

The following formula is suitable for the calculation of renormalized second quantization operators. Recall that for all $T \in L(E_{\mathbb{C}}, E_{\mathbb{C}})$, we have $\Gamma(T) \in L((E)_{\beta}, (E)_{\beta})$, where $0 \leq \beta < 1$.

Corollary 6.11. *Let $r \in \mathbb{N}$, $1 > \beta \geq \frac{\max(0, r-2)}{r}$, $\kappa_{0,r} \in (E_{\mathbb{C}}^{\otimes r})_{sym}^*$. Then for all $T \in L(E_{\mathbb{C}}, E_{\mathbb{C}})$ we have*

$$\begin{aligned} \Gamma_{\kappa_{0,r}}(T) &= \Gamma(T) \circ \exp\left(\frac{1}{2}\Xi_{0,2}((T^{\otimes 2} - \text{Id}^{\otimes 2})^*(\tau))\right) \\ &\quad \circ \exp\left(-\frac{1}{2}\Xi_{0,r}((T^{\otimes r} - \text{Id}^{\otimes r})^*(\kappa_{0,r}))\right). \end{aligned}$$

Proof. On the one hand we calculate

$$\begin{aligned} &\Gamma(T) \circ \exp\left(\frac{1}{2}\Xi_{0,2}((T^{\otimes 2} - \text{Id}^{\otimes 2})^*(\tau))\right) \circ \exp\left(-\frac{1}{2}\Xi_{0,r}((T^{\otimes r} - \text{Id}^{\otimes r})^*(\kappa_{0,r}))\right) \Phi_{\xi} \\ &= \left(\exp\left(\frac{1}{2}\langle \tau, (T^{\otimes 2} - 1)\xi^{\otimes 2} \rangle\right)\right) \cdot \left(\exp\left(-\frac{1}{2}\langle (\kappa_{0,r}), (T^{\otimes r} - 1)\xi^{\otimes r} \rangle\right)\right) \Phi_{T\xi} \end{aligned}$$

On the other hand by Definition 6.8

$$\begin{aligned} &\Gamma_{\kappa_{0,r}}(T)(\Phi_{\xi}) \\ &= \exp\left(\frac{1}{2}(\Delta_G - \Xi_{0,r}(\kappa_{0,r}))\right) \circ \Gamma(T) \circ \exp\left(-\frac{1}{2}(\Delta_G - \Xi_{0,r}(\kappa_{0,r}))\right)(\Phi_{\xi}) \\ &= \exp\left(-\frac{1}{2}\langle \xi, \xi \rangle - \langle \kappa_{0,r}, \xi^{\otimes r} \rangle\right) \cdot \exp\left(\frac{1}{2}(\Delta_G - \Xi_{0,r}(\kappa_{0,r}))\right)(\Phi_{T\xi}) \\ &= \exp\left(-\frac{1}{2}\langle \xi, \xi \rangle - \langle \kappa_{0,r}, \xi^{\otimes r} \rangle\right) \cdot \exp\left(\frac{1}{2}\langle T\xi, T\xi \rangle - \langle \kappa_{0,r}, (T\xi)^{\otimes r} \rangle\right) \Phi_{T\xi} \\ &= \left(\exp\left(\frac{1}{2}\langle \tau, (T^{\otimes 2} - 1)\xi^{\otimes 2} \rangle\right)\right) \cdot \left(\exp\left(-\frac{1}{2}\langle (\kappa_{0,r}), (T^{\otimes r} - 1)\xi^{\otimes r} \rangle\right)\right) \Phi_{T\xi} \end{aligned}$$

\square

We consider some concrete examples to Corollary 6.11.

Example 6.12. Let $b \in \mathbb{C}$, $\theta \in \mathbb{R}$ and $\kappa_{0,2} \in (E_{\mathbb{C}}^{\otimes 2})_{sym}$. Then

(i)

$$\Gamma_{\kappa_{0,2}}(b \text{ Id}) = \Gamma(b \text{ Id}) \exp\left(\frac{1}{2}(b^2 - 1)\Xi_{0,2}(\tau - \kappa_{0,2})\right)$$

(ii)

$$\Gamma_{\kappa_{0,2}}(e^{i\theta} \text{ Id}) = \Gamma(e^{i\theta} \text{ Id}) \circ \exp\left(i \cdot e^{i\theta} \sin \theta \Xi_{0,2}(\tau - \kappa_{0,2})\right)$$

(iii)

$$\Gamma_{\frac{1}{2}\tau}(e^{i\theta} \text{ Id}) = \Gamma(e^{i\theta} \text{ Id}) \circ \exp\left(\frac{i}{2} e^{i\theta} \sin \theta \Delta_G\right)$$

By [12, Lemma 5.6.1, p.140] we conclude that $\Gamma_{\frac{1}{2}\tau}(e^{i\theta} \text{ Id})$ is the adjoint operator of the Fourier-Mehler transform \mathcal{F}_{θ} .

The next example gives conditions under which the Fourier-Gauss transforms are renormalized second quantization operators.

Example 6.13. Let $a, b \in \mathbb{C}$ and $b \notin \{-1, 1\}$. Choose σ with $\sigma^2 = \frac{a^2}{1-b^2}$. Then

$$\Gamma_{\sigma^2\tau}(b \text{ Id}) = \mathfrak{G}_{a,b}$$

Which follows immediately by Example 6.12(i) with $\kappa_{0,2} = \sigma^2\tau$.

Remark 6.14. In the special case $a = 0$, we have for the scaling operator $S_b := \mathfrak{G}_{0,b} = \Gamma_{0\tau}(b \text{ Id})$, i.e. $S_b(\langle x^{\otimes n}, f_n \rangle) = \langle x^{\otimes n}, b^n \cdot f_n \rangle$ for all $n \in \mathbb{N}_0$, $f_n \in (E_{\mathbb{C}}^{\otimes n})_{sym}$. With $a, b \in \mathbb{C}$, $b \notin \{-1, 1\}$ and $\sigma^2 = \frac{a^2}{1-b^2}$, we have

$$\Gamma_{\sigma^2\kappa_{0,2}}(b \text{ Id}) = \Gamma(b \text{ Id}) \circ \exp\left(\frac{1}{2} [a^2\Xi_{0,2}(\kappa_{0,2}) + (b^2 - 1) \Delta_G]\right)$$

Theorem 6.15. Let $\{T_{\theta}\}_{\theta \in \mathbb{R}}$ be a differentiable one parameter subgroup of $GL(E_{\mathbb{C}})$ with infinitesimal generator T' . Further let $r \in \mathbb{N}$, $1 > \beta \geq \frac{\max(0, r-2)}{r}$ and $\kappa_{0,r} \in (E_{\mathbb{C}}^{\otimes r})_{sym}^*$. Then $\{\Gamma_{\kappa_{0,r}}(T_{\theta})\}_{\theta \in \mathbb{R}}$ is a regular one parameter subgroup of $GL((E)_{\beta})$ with

$$d\Gamma_{\kappa_{0,r}}(T') = \frac{d}{d\theta} \Big|_{\theta=0} \Gamma_{\kappa_{0,r}}(T_{\theta}) = d\Gamma(T') + \frac{1}{2}\Xi_{0,2}(\gamma_2(T')^* \tau) - \frac{1}{2}\Xi_{0,r}(\gamma_r(T')^* \kappa_{0,r})$$

Proof. By Corollary 6.11 we have

$$\begin{aligned} \Gamma_{\kappa_{0,r}}(T_{\theta}) &= \Gamma(T_{\theta}) \circ \exp\left(\frac{1}{2}\Xi_{0,2}((T_{\theta}^{\otimes 2} - \text{Id}^{\otimes 2})^*(\tau))\right) \\ &\quad \circ \exp\left(-\frac{1}{2}\Xi_{0,r}((T_{\theta}^{\otimes r} - \text{Id}^{\otimes r})^*(\kappa_{0,r}))\right). \end{aligned}$$

First note that $(G \circ \Gamma(T_{\theta}) \circ G^{-1})_{\theta \in \mathbb{R}}$ is obviously a regular one-parameter subgroup of $GL((E)_{\beta})$ for all $G \in GL((E)_{\beta})$.

Consequently $(\Gamma_{0\tau}(T_{\theta}))_{\theta \in \mathbb{R}}$, with

$$\Gamma_{0\tau}(T_{\theta}) = \Gamma(T_{\theta}) \circ \exp\left(\frac{1}{2}\Xi_{0,2}((T_{\theta}^{\otimes 2} - \text{Id}^{\otimes 2})^*(\tau))\right),$$

is a regular one-parameter subgroup of $GL((E)_{\beta})$. Because

$$\begin{aligned} \Gamma_{\kappa_{0,r}}(T_\theta) = & \left[\Gamma(T_\theta) \circ \exp\left(\frac{1}{2}\Xi_{0,2}((T_\theta^{\otimes 2} - \text{Id}^{\otimes 2})^*(\tau))\right) \right] \\ & \circ \exp\left(-\frac{1}{2}\Xi_{0,r}((T_\theta^{\otimes r} - \text{Id}^{\otimes r})^*(\kappa_{0,r}))\right), \end{aligned}$$

the claim follows by Proposition 5.9, then Proposition 5.12 and Proposition 5.13. \square

Corollary 6.16. *Let $a, b \in \mathbb{C}$ and $b \neq 0$. Then*

$$bN_{(1-\frac{a}{b})\tau} = \frac{d}{d\theta}\Big|_{\theta=0} \Gamma_{(1-\frac{a}{b})\tau}(e^{b\theta} \text{Id}) = a\Delta_G + bN.$$

Proof. By [12, Proposition 4.6.13., p.107] we have $N := \Xi_{1,1}(\tau) = d\Gamma(\text{Id})$. Because $b\text{Id}$ is the infinitesimal generator of $\{e^{b\theta}\text{Id}\}_{\theta \in \mathbb{R}}$, it follows by Proposition 5.12 that

$$\frac{d}{d\theta}\Big|_{\theta=0} (\Gamma(e^{b\theta} \text{Id})) = d\Gamma(b \text{Id}) = \Xi_{1,1}(b\tau) = bN,$$

for $b \in \mathbb{C}$.

On the other hand, in the general case $r \in \mathbb{N}$, $1 > \beta \geq \frac{\max(0, r-2)}{r}$ and $\kappa_{0,r} \in (E_{\mathbb{C}}^{\otimes r})_{sym}^*$, it holds from Theorem 6.15, with $T' = \text{Id}$, that

$$N_{\kappa_{0,r}} = \frac{d}{d\theta}\Big|_{\theta=0} \Gamma_{\kappa_{0,r}}(e^\theta \text{Id}) = N + \Delta_G - \frac{r}{2}\Xi_{0,r}(\kappa_{0,r}). \quad (6.1)$$

Multiplicating both sides with b , the claim is immediate with $\kappa_{0,r} = (1 - \frac{a}{b})\tau$. \square

Remark 6.17. Let $b \neq 0$. The regular one-parameter subgroup

$$\left\{ \Gamma_{(1-\frac{a}{b})\tau}(e^{b\theta} \text{Id}) \right\}_{\theta \in \mathbb{R}} \subset GL((E)_\beta), \quad 0 \leq \beta < 1$$

can, Example 6.13, be identified as the Fourier-Gauss transforms $\{\mathfrak{G}_{x, e^{b\theta}}\}$, where $x^2 = (1 - \frac{a}{b})(1 - e^{2b\theta})$.

The case $b = 0$ is solved by 5.14. We get $\{\mathfrak{G}_{\sqrt{2a\theta}, 1}\}$ as solution.

Note, that the special choice of x from x^2 has no influence, because $\{\mathfrak{G}_{x, e^{b\theta}}\}$ only depends on x^2 and $e^{b\theta}$.

We summarize this discussion using the definition of the Fourier-Gauss transform in [10, Definition 11.24, p. 164]. The following theorem is a generalization of the Mehler formula for the Ornstein-Uhlenbeck semigroup, see e.g. [7, p. 237]

Theorem 6.18. *Let $a, b \in \mathbb{C}$, $0 \leq \beta < 1$. Then $a \cdot \Delta_G + b \cdot N$ is the infinitesimal generator of the following regular transformation group $\{P_{a,b,t}\}_{t \in \mathbb{R}} \subset GL((E)_\beta)$:*

(i) if $b \neq 0$ then for all $\varphi \in (E)_\beta$, $t \in \mathbb{R}$:

$$P_{a,b,t}(\varphi) = \int_{E^*} \varphi\left(\sqrt{\left(1 - \frac{a}{b}\right)(1 - e^{2bt})} \cdot x + e^{bt} \cdot y\right) d\mu(x)$$

(ii) if $b = 0$ then $\varphi \in (E)_\beta$, $t \in \mathbb{R}$:

$$P_{a,0,t}(\varphi) = \int_{E^*} \varphi(\sqrt{2at} \cdot x + y) d\mu(x)$$

On an informal level the second case of the above theorem may be considered as a special case of the first one. Note that by the rules of l'Hôpital we have $\lim_{b \rightarrow 0} (b - a) \frac{1 - e^{2bt}}{b} = 2at$.

7. Summary and Bibliographical Notes

The investigation of regular transformation groups in this manuscript can be compared with [3], where a two-parameter transformation group G on the space of white noise test functions (E) was constructed, which includes the adjoints of Kuo's Fourier and Kuo's Fourier-Mehler transforms. Their description of a differentiable one-parameter subgroup of G whose infinitesimal generator is $a\Delta_G + bN$ is identical with our findings. However using the unique representation of white noise test functions via generalized Wick tensors and corresponding renormalized operators we have presented a different way which leads to a more general result for the spaces $(E)_\beta$. Using convolution operators and the fact that differentiable one parameter transformation groups on nuclear Fréchet spaces are always regular, convergence problems could be solved.

Acknowledgment. The authors would like to thank to Prof. H.-H. Kuo for his encouragement to publish this manuscript. W. Bock would like to thank for the financial support from the FCT project PTDC/MAT-STA/1284/2012. We thank the referees for their helpful suggestions to improve the presentation of this article.

References

1. Berezansky, Y. M. and Kondratiev, Y. G. (1995): *Spectral Methods in Infinite-dimensional Analysis. Vol. 2*. Dordrecht: Kluwer Academic Publishers. Translated from the 1988 Russian original by Malyshev P. V. and Malyshev D. V. and revised by the authors.
2. Bock, W.: *Generalized Scaling Operators in White Noise Analysis and Applications to Hamiltonian Path Integrals with Quadratic Action*, in *Stochastic and Infinite Dimensional Analysis*, Springer, 2015.
3. Chung, D. M. and Ji, U. C.: *Transformation Groups on White Noise Functionals and Their Applications*, *Applied Mathematics & Optimization*, 1998, Volume 37, Number 2, Seiten 205–223.
4. Chung, D. M., Ji, U. C., and Obata, N.: Transformations on White Noise functions associated with second order differential operators of diagonal type, *Nagoya Math. J.*, **149** (1998), 173–192.
5. Chung, C.-H., Chung, D. M., and Ji, U. C.: One-parameter groups and cosine families of operators on white noise functions, *Bull. Korean. Math. Soc.* **37** (2000), No. 5, 687–705.
6. Floret, K. and Wloka, J.: *Einführung in die Theorie der Lokalkonvexen Räume*, Lecture Notes in Mathematics 1968, (56) Walter Springer-Verlag, Berlin Heidelberg New York.
7. Hida, T., Kuo, H.-H., Potthoff, J., and Streit L.: *White Noise: An Infinite-dimensional Calculus*, Kluwer Academic Publishers, Dordrecht, 1993.
8. Hida, T., Obata, N., and Saitô, K.: Infinite dimensional rotations and Laplacians in terms of white noise calculus, *Nagoya Math. J.* **128** (1992), 65–93.
9. Kondratiev, Yu. G. Leukert, P., Potthoff, J., Streit, L., and Westerkamp, W.: Generalized functionals in Gaussian spaces: The characterization theorem revisited. *J. Funct. Anal.* **141** (1996) Nr.2. 301–318.
10. Kuo, H.-H.: *White Noise Distribution Theory*, CRC Press, Boca Raton, 1996.
11. Lee, Y.-J.: Integral transforms of analytic functions on abstract Wiener spaces, *J. Funct. Anal.* **47** (1983) 153–164.

12. Obata, N.: *White Noise Analysis and Fock Space*, Lecture Notes in Mathematics, Vol. 1577, Springer-Verlag, Berlin, 1994.
13. Obata, N. and Ouerdiane, H.: A note on convolution operators in white noise calculus, *IDAQP*, **14**, No. 4 (2011) 661–674, World Scientific Publishing Company, DOI: 10.1142/S0219025711004535.
14. Schaefer, H. H. and Wolff, M. P.: *Topological Vector Spaces*, Springer-Verlag New York Berlin Heidelberg, second edition 1999, ISBN 0-387-98726-6 SPIN 10707604.

MAXIMILIAN BOCK: SAARSTAHL AG, INFORMATIK, BISMARCKSTR. 57-59, 66333 VÖLKLINGEN, GERMANY

E-mail address: `maximilianbock61@gmail.com`

WOLFGANG BOCK: TECHNOMATHEMATICS GROUP, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KAISERSLAUTERN, 67653 KAISERSLAUTERN, GERMANY

E-mail address: `bock@mathematik.uni-kl.de`