

1-1-2012

The extrinsic proteins of Photosystem II

Terry M. Bricker
Louisiana State University

Johnna L. Roose
Louisiana State University

Robert D. Fagerlund
University of Otago

Laurie K. Frankel
Louisiana State University

Julian J. Eaton-Rye
University of Otago

Follow this and additional works at: https://digitalcommons.lsu.edu/biosci_pubs

Recommended Citation

Bricker, T., Roose, J., Fagerlund, R., Frankel, L., & Eaton-Rye, J. (2012). The extrinsic proteins of Photosystem II. *Biochimica et Biophysica Acta - Bioenergetics*, 1817 (1), 121-142. <https://doi.org/10.1016/j.bbabo.2011.07.006>

This Article is brought to you for free and open access by the Department of Biological Sciences at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.



Review

The extrinsic proteins of Photosystem II[☆]

Terry M. Bricker^{a,*}, Johnna L. Roose^a, Robert D. Fagerlund^b, Laurie K. Frankel^a, Julian J. Eaton-Rye^b
^a Department of Biological Sciences, Biochemistry and Molecular Biology Section, Louisiana State University, Baton Rouge, LA 70803, USA

^b Department of Biochemistry, University of Otago, P.O. Box 56, Dunedin, New Zealand

ARTICLE INFO

Article history:

Received 11 May 2011

Received in revised form 11 July 2011

Accepted 12 July 2011

Available online 27 July 2011

Keywords:

PsbO

PsbP

PsbQ

PsbU

PsbV

ABSTRACT

In this review we examine the structure and function of the extrinsic proteins of Photosystem II. These proteins include PsbO, present in all oxygenic organisms, the PsbP and PsbQ proteins, which are found in higher plants and eukaryotic algae, and the PsbU, PsbV, CyanoQ, and CyanoP proteins, which are found in the cyanobacteria. These proteins serve to optimize oxygen evolution at physiological calcium and chloride concentrations. They also shield the Mn₄CaO₅ cluster from exogenous reductants. Numerous biochemical, genetic and structural studies have been used to probe the structure and function of these proteins within the photosystem. We will discuss the most recent proposed functional roles for these components, their structures (as deduced from biochemical and X-ray crystallographic studies) and the locations of their proposed binding domains within the Photosystem II complex. This article is part of a Special Issue entitled: Photosystem II.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Photosystem II (PSII) functions as a light energy-driven water-plastoquinone oxidoreductase. In higher plants, eukaryotic algae and cyanobacteria, light is trapped by and funneled through light-harvesting pigment-protein complexes (light-harvesting complex II or phycobilisomes) to the reaction center of PS II, which contains a specialized pigment complex, P₆₈₀. After excitation, P₆₈₀ becomes photooxidized and donates an electron to the primary acceptor of PS II, a protein-bound pheophytin. This charge separation is then further stabilized by the transfer of this electron, first to Q_A and then to Q_B, protein-bound plastoquinones. The accumulation of two reducing equivalents on Q_B leads to its protonation and the formation of plastoquinol, which is then released from PS II. Concomitant to these electron transfers, P₆₈₀⁺ is reduced by Y_Z, a tyrosyl residue (¹⁶¹Y) located on the D1 protein. The subsequent reduction of Y_Z leads to the accumulation of an oxidizing equivalent in the oxygen-evolving complex. The accumulation of four such oxidizing equivalents, in a manner consistent with the observed S-state transitions, leads to the release of oxygen from the complex (for reviews, see [1–4]). The oxygen-evolving complex contains a metal cluster consisting of four mixed valence manganese ions, a calcium ion, and five oxo ligands (Mn₄CaO₅ cluster). Two chloride ions are also present flanking the cluster: Cl[−]1 is 6.7 Å from Mn4 and Cl[−]2 is 7.4 Å from Mn2 [5] in the S₁ oxidation state. All of the protein ligands to this metal cluster are

provided by the D1 protein, with the singular exception that one manganese ligand is provided by the ³⁵⁴E of CP43 [5–8].

In higher plants and cyanobacteria at least seven major intrinsic proteins appear to be absolutely required for oxygen evolution [9–11]. These are CP47, CP43, the D1 protein, the D2 protein, the α and β subunits of cytochrome *b*₅₅₉, and the 4.8 kDa *psbI* gene product. Insertional inactivation of the genes for any of these components leads to a loss of normal PS II assembly and a loss of the ability to evolve oxygen. Additionally, a large number of low molecular mass intrinsic components are associated with PS II, although the functions of the majority of these proteins remain obscure, as their deletion only modestly affects PS II function. While PS II complexes containing only these components can evolve oxygen, they do so at very low rates (about 25% of control) and require high, non-physiological levels of calcium and chloride [11]. In higher plants, three extrinsic proteins, PsbO, PsbP and PsbQ, are required for maximal rates of oxygen evolution under physiological conditions. A fourth, possibly extrinsic protein, PsbR, is also present in higher plants and green algae. In cyanobacteria, the PsbO and PsbQ homologue (CyanoQ) proteins are present along with the PsbU and PsbV extrinsic components [12]. These latter two proteins are not present in higher plants. The cyanobacteria also contain a PsbP homologue (CyanoP); the stoichiometry of this component remains controversial [13,14]. Interestingly, while the green algae contain the same extrinsic components present in higher plants (PsbO, PsbP, PsbQ and PsbR), the red algae contain PsbO, PsbP, PsbQ' (a 20 kDa homologue of CyanoQ), PsbU and PsbV [15,16]; PsbR is not present in the red algae.

In this review, we will examine the structural and functional properties of these extrinsic protein components which function in support of oxygen evolution. We will principally focus on findings

[☆] This article is part of a Special Issue entitled: Photosystem II.

* Corresponding author at: Department of Biological Sciences Louisiana State University Baton Rouge, LA 70803, USA. Tel.: +1 225 578 1555; fax: +1 225 578 2596. E-mail address: tbric@lsu.edu (T.M. Bricker).




Fig. 1. *Thermosynechococcus vulcanus* Photosystem II dimer highlighting the extrinsic subunits of the photosystem. The crystal structure of Umena et al. (PDB ID: 3ARC) is shown [5]. A., is a side view of the photosystem from within the plane of the thylakoid membrane, while B., illustrates the photosystem as viewed from the luminal space. PsbO is shown in orange, PsbU is shown in blue and PsbV is shown in green. Surface representations are provided for these subunits. All other PS II subunits are shown in gray. All figures were prepared in PyMol [259].

obtained during the last decade, which has witnessed an explosive growth in our understanding of the roles played by these proteins in the photosystem. Organizationally, we will first examine the PsbO, PsbP and PsbQ proteins. These components (or their homologues) are present in virtually all oxygenic organisms. The enigmatic PsbR protein, which is not present in cyanobacteria or red algae, will also be briefly examined. The cyanobacterial PsbU and PsbV proteins along with the cyanobacterial homologues of PsbQ (CyanoQ) and PsbP (CyanoP) will then be discussed. It should be noted here that we will not examine the extended PsbP and PsbQ families of proteins which are found in higher plants. The numerous members of these protein families may play other important roles within PS II, the Ndh dehydrogenase complex, and possibly other systems (for reviews, see [17–19]).

1.1. PsbO structure

Sequence alignments for the PsbO protein have been presented elsewhere [20,21] and indicate that PsbO is conserved across the higher plants, algal and cyanobacterial clades. Our understanding of the structural association of the PsbO protein within PS II comes principally from the crystal structures of the photosystem which have been obtained from the thermophilic cyanobacteria. These studies have, in large measure, confirmed earlier investigations which indicated that the PsbO component, in solution, exhibited a high degree of β -sheet secondary structure [22,23] and that the protein was significantly elongated [24,25]. It should be noted that no crystal structures for the higher plant protein either as an isolated subunit or associated with the photosystem are currently available.

PsbO, in association with PS II from *Thermosynechococcus vulcanus* [5] (henceforth *T. vulcanus*), is shown in Fig. 1 in orange. This protein is located on the luminal face of the complex and interacts with a variety of other PS II subunits including CP43, CP47, D1, D2 and PsbU. Since the PsbO protein can be removed from PS II membranes by treatment with 1 M CaCl_2 , it is hypothesized that charge-pair interactions provide the principal binding energy of PsbO to the photosystem. Within this context, it should be noted that PsbO remains firmly attached to PS II membranes even in the presence of 2 M NaCl. The higher charge density of calcium (or other divalent cations) appears to be required to disrupt the interaction of PsbO with the PS II core components. The PsbO protein can also be removed from PS II membranes by treatment with 2.6 M urea + 200 mM NaCl [9]. In this instance, the urea is hypothesized to weaken inter- or intramolecular hydrogen bond interactions sufficiently to allow release of the PsbO component by the 200 mM NaCl.

To identify possible salt-bridge interactions of surfaces within the PS II crystal structure, we utilized the program KFC [26], which uses shape and residue properties to identify contact domains on multimeric proteins predicted to contribute the highest binding energy. We then examined the residues within the contact domains for possible charge-pair interactions. Additionally, we used the program CAPTURE [27] to identify possibly relevant cation- π interactions. The results from this analysis are shown in Table 1.¹ CP43, CP47, D1 and D2 all appear to possess residues which could contribute to the binding of PsbO.

Only a small subset of these residues, principally on the CP47 subunit, has been examined in more detail. Within the *T. vulcanus* crystal structure [5], a charge-pair “complex” appears to be formed involving the residues ³⁸⁵R:CP47–¹⁶⁹D:PsbO–⁴²²R:CP47–¹⁷⁹E:PsbO–⁴²³K:CP47 (Fig. 2). Please note that ³⁸⁴R:CP47 appears to interact with the C-terminal carboxyl group of the D2 protein (³⁵²L:D2). These residues are partially buried, which would increase the relative strength of the charge-pair interactions. The basic residue pairs of CP47, ³⁸⁴R³⁸⁵R and ⁴²²R⁴²³R, have been examined by site-directed mutagenesis in *Synechocystis* sp. PCC 6803 (henceforth *Synechocystis* 6803). Positive charges at these positions are absolutely conserved across all CP47 proteins. While alteration of the ⁴²²R⁴²³R pair to neutral residues had a relatively small effect on oxygen evolution [28], modification of residues ³⁸⁴R³⁸⁵R to either neutral (³⁸⁴G³⁸⁵G) [29] or negatively charged (³⁸⁴E³⁸⁵E) residues [28] had a dramatic effect. Individual site-directed modification at either residue ³⁸⁴R or ³⁸⁵R also yielded mutants with significantly decreased oxygen evolution capability [28]. The double mutants evolved oxygen at rates comparable to that reported for the Δ psbO mutant. Later studies, indicating that permeabilized thylakoids isolated from the CP47 ³⁸⁴E³⁸⁵E mutant bound the PsbO protein very weakly,² and that the Δ psbO:³⁸⁴E³⁸⁵E triple mutant exhibited a phenotype very similar to the Δ psbO mutant, provided strong evidence that ³⁸⁴R³⁸⁵R of CP47 provided an important binding site for PsbO to PS II [30]. It should be noted that detergent-permeabilized thylakoids isolated from the CP47 ³⁸⁴G³⁸⁵G mutant bound the PsbO protein significantly more strongly than the CP47 ³⁸⁴E³⁸⁵E mutant [30], indicating that other, yet unidentified residues also contribute to the association of PsbO with PS II. Neither of the PsbO residues (¹⁶⁹D and ¹⁷⁹E) predicted from the KFC analysis described above to participate

¹ The numbering of the specific amino acid residues in the various PS II proteins is problematic as various authors have used numbering derived from the immature translated gene sequence (with or without removal of the N-terminal methionyl residue) or from the fully processed, mature protein, from a variety of different organisms. Except where indicated, we have used the numbering system from the *T. vulcanus* crystal structure of Umena et al. [5].

² His-tagged PS II particles containing mutations at residues ³⁸⁴R³⁸⁵R of CP47 also lacked the PsbO subunit (C. Putnam-Evans and T. M. Bricker, unpublished).

Table 1

Possible interacting residues of *Thermosynechococcus vulcanus* PsbO with other subunits of PS II. Binding domains were identified with KFC [26]. A putative cation– π interaction was identified by the program CAPTURE. A possible inter-monomer interaction is shown in parentheses. The numbering scheme is for *T. vulcanus* as derived from the Umena et al. [5] crystal structure. It should be noted that these distances are slightly different for the two monomers. The closest distances are shown below.

PsbO:CP 43	PsbO:CP 47	PsbO:D1	PsbO:D2
<i>Charge-pair interactions</i>			
⁸ D:362R, 3.05 Å	¹⁶⁹ D:385R, 4.88 Å	⁶⁹ K:103D, 3.2 Å	¹⁶⁰ K:302E, 3.41 Å
⁹⁹ D:381K, 2.45 Å	¹⁶⁹ D:422R, 2.70 Å	⁷³ R:104E, 3.39 Å	²²⁸ H:310E, 3.19 Å
	¹⁷⁹ E:422R, 4.27 Å		
	¹⁷⁹ E:423K, 5.20 Å		
	(⁵⁹ K:307E, 4.21 Å)		
<i>Cation–π interaction</i>			
	⁷ Y:370R, 4.67 Å		

in these charge-pair interactions with CP47 has been examined by site-directed mutagenesis.

PsbO also may participate in an inter-monomer interaction with CP47, possibly stabilizing the PS II dimer [20]. KFC analysis indicates that ⁵⁹K of PsbO could form a putative charge-pair interaction with ³⁰⁷E of CP47. Such an interaction might explain the loss of PS II dimers in $\Delta psbO$ mutants [31]. An alternative hypothesis, however, is that increased photodamage occurring in the absence of PsbO in the $\Delta psbO$ mutant accelerates the rate of monomer formation as a prelude to the repair of the photosystem.

Two residues in CP43 (³⁶²R and ³⁸¹K) are also predicted to participate in charge-pair interactions with PsbO (Table 1). The residues ³⁶²R and ³⁸¹K lie within the *Synechocystis* 6803 deletion mutants³ $\Delta F345/E354$ and $\Delta L364/D370$, respectively [32]. These mutants did not assemble functional PS II reaction centers. Since the observed phenotype was much more extreme than that observed in the $\Delta psbO$ mutant, the loss of assembly of the photosystem cannot be ascribed to the possible loss of PsbO binding alone. The severe phenotype may be the result of the removal of ten and seven residues, respectively in these two deletion strains. The effects of alteration of the individual CP43 residues ³⁶²R and ³⁸¹K have not been directly probed by site-directed mutagenesis. The putative ⁸D:PsbO–³⁶²R:CP43 interaction is particularly interesting, however. A variety of N-terminal modifications of higher plant PsbO have been shown to affect PsbO binding to the photosystem. Eaton-Rye and Murata [33], using spinach PS II membranes, demonstrated that proteolytic removal of the N-terminal 16–18 amino acid residues from PsbO abolished its ability to bind to PS II. Popelkova et al. [34] identified two N-terminal domains⁴ (⁷TYD¹⁰E and ¹⁵TYL¹⁸E) in the spinach PsbO protein which the authors demonstrated were required to bind the two copies of the PsbO protein per PS II monomer (see below) in higher plants. One of these domains is missing at the N-terminus of the cyanobacterial protein. Sequence alignment of higher plant and cyanobacterial PsbO sequences [21] indicates that the cyanobacteria uniformly exhibit a nine–ten amino acid deletion at the N-terminus (the ⁷TYD¹⁰E domain lies within this deletion). These results may explain a major discrepancy between site-directed mutagenesis studies performed in higher plants and cyanobacteria. In spinach, site-directed modification of ⁹D to either ⁹N or ⁹K had neither an effect on the binding of the modified PsbO protein to PsbO-depleted PS II membranes nor the restoration of oxygen-evolving activity [35]. In *Synechocystis* 6803, however, modification of ⁹D to ⁹K weakened the binding of PsbO to thylakoid membranes and resulted in a moderate loss of oxygen-evolving activity [30]. The findings of Popelkova et al. [34] demonstrate that the ⁹D residue is not equivalent in the higher plant and cyanobacterial systems; in spinach, this residue lies in the first of two

binding domains, while in cyanobacteria it lies in the sole binding domain present in these organisms, which is equivalent to the second binding domain found in higher plants (please note that the ⁹D residue in *Synechocystis* 6803 is equivalent to the ⁸D residue in the *T. vulcanus* crystal structure [5]). The putative interacting residue, ³⁶²R:CP43, has not been examined by site-directed mutagenesis.

Finally, a number of residues in the D1 and D2 proteins are predicted to interact via charge-pair interactions with PsbO (Table 1). No site-directed mutagenesis experiments examining the residues in D1 and D2 with respect to PsbO binding have been reported. ³⁰²E of the D2 protein is of particular interest, however. Its putative binding partner is ¹⁶⁰K of PsbO, the modification of which dramatically altered the binding properties of the mutated PsbO protein in *Thermosynechococcus elongatus*⁵ (henceforth, *T. elongatus*) [36]. Other residues in this vicinity have also been implicated in the functional association of PsbO with the photosystem. Examination of the crystal structure indicates that the backbone carbonyl of ¹⁵⁸D of PsbO may be hydrogen-bound to ³³⁴R of D1 (3.0 Å). This latter residue was modified in *Synechocystis* 6803 in an attempt to determine its involvement in extrinsic protein binding [37]. While no evidence was found indicating this role, the resultant mutant did exhibit a variety of PS II defects including altered S₂ state properties and altered oxygen release kinetics. Modification of ¹⁵⁸D of PsbO yielded a mutant⁶ (D159N) which exhibited a 30% loss of steady-state oxygen evolution and loss of photoautotrophy in a $\Delta psbV$ background, neither of which could be directly attributed to loss of PsbO from the photosystem [38]. These authors hypothesized that ¹⁵⁸D formed an intramolecular charge-pair with ¹⁶²R. Examination of the *T. vulcanus* crystal structure [5] indicates that this is plausible, as these residues are separated by 2.7 Å.

Many of the results discussed above apply directly only to cyanobacterial PS II. It must be noted that while the PS II crystal structures have provided important insights into the structure of the photosystem, some questions remain. None of the current structures contain the CyanoQ or the CyanoP subunits. The former is almost certainly a stoichiometric component of the photosystem [12,13], while the latter also may be present in stoichiometric quantities [14], although this is controversial [13,39]. The observation that loss of CyanoQ destabilizes the binding of PsbV [40] also raises questions concerning the overall integrity of the observed subunit composition in the cyanobacterial structures currently available.

A number of studies, obtained principally by biochemical examination of higher plant PS II, appear inconsistent with current cyanobacterial crystal structures. Given the absence of a higher plant crystal structure it is unclear if these differences are due to real differences between the cyanobacterial and higher plant photosystem, possible limitations in the current cyanobacterial structures (particularly with regard to subunit composition), or investigator error. Numerous crosslinking studies have indicated that in higher plants [35,41,42], green algae [43] and cyanobacteria [44] the PsbO protein can be crosslinked to CP47 using the 0-length crosslinker 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC). EDC crosslinks amino groups which are interacting with carboxylates via charge-pair interactions [45] and which are exposed to the bulk solvent. Odom et al. [46] mapped the crosslinked domain between these two components and reported that residues within the N-terminal tryptic peptide ¹E–⁷⁶K of PsbO were crosslinked to residues within the domain ³⁶⁴E–⁴⁴⁰D of CP47. Examination of the *T. vulcanus* [5] crystal structure indicates that the only possible crosslinking candidates for this interaction are ⁴⁰⁵E of CP47 and the residue ⁶⁶K of PsbO in spinach. The latter residue corresponds to ⁵⁹K in *T. vulcanus*. Interestingly, this would be the result of inter-monomer crosslinking. However, these

⁵ In the original paper, Motoki et al. [36] examined "...a thermophilic cyanobacteria, *Synechococcus elongatus*". This organism is now named *Thermosynechococcus elongatus*. *Synechococcus elongatus* is now the genus and species name used for *Synechococcus* sp. PCC 7942.

⁶ *Synechocystis* 6803 numbering.

³ *Synechocystis* 6803 numbering.

⁴ Spinach numbering.




Fig. 2. Putative binding domain on CP47 for the PsbO protein. Detail from the *T. vulcanus* crystal structure is illustrated (PDB ID: 3ARC[5]). PsbO is shown in blue, while CP47 is shown in gray and the C-terminal residue of D2 (³⁵²L) is shown in yellow. Putative charge-pair interactions are indicated by dotted lines and apparent distances are shown in angstroms (Å).

residues are separated by nearly 10 Å, thus precluding these residues from forming the charge-pair interaction required for EDC crosslinking.

Frankel and Bricker [47] reported that upon removal of the PsbO component from spinach PS II membranes, the domains ³⁰⁴K–³²¹K and ³⁸⁹K–⁴¹⁹K of CP47 became susceptible to modification with the amino group-reactive reagent NHS-biotin. Possible biotinylated residues include ³⁰⁴K, ³⁰⁸K and ³²¹K in the first domain and ³⁸⁹K, ⁴¹⁸K and ⁴¹⁹K in the second. Four of these CP47 residues are conserved in *T. vulcanus* (³⁰⁸K, ³²¹K, ³⁸⁹K, and ⁴¹⁸K). None of these residues appears closely associated with the PsbO protein in the *T. vulcanus* structure. Interestingly, residues ³⁰⁸K, ³⁸⁹K, and possibly ⁴¹⁸K are surface-exposed residues. If the structure of higher plant PS II is similar to that observed for cyanobacteria, these residues should have been labeled with NHS-biotin. No biotinylation, however, of any residues on CP47 was observed in the presence of PsbO in the higher plant system [47].

Enami et al. [48] used the crosslinker hexamethylene-diisocyanate (HMDI) to crosslink PS II components in spinach NaCl-washed n-heptylthioglucoside-solubilized spinach PS II particles. HMDI can react with a variety of functional groups and can crosslink proteins which are within 11 Å [49]. Individual crosslinked products were observed which contained both the PsbO protein and either PsbI or PsbE (α-subunit of cytochrome *b*₅₅₉). Examination of this result within the context of the *T. vulcanus* crystal structure yields interesting observations. The closest approach of PsbO with PsbI is 8–10 Å, a distance which is consistent with the crosslinking result [48]. The nearest approach of PsbE with PsbO, however, is over 30 Å, a distance which cannot be spanned with HMDI. One possible explanation for this result could be inter-dimer crosslinking. However, this is unlikely since the experiment was performed on solubilized PS II particles and, consequently, the effective local PS II concentration would be much lower than would be observed in PS II membranes.

2. Stoichiometry: cyanobacterial vs. higher plant

In all cyanobacterial PS II crystal structures published to date, one copy of the PsbO protein is present per monomer. This has led many workers to assume that one copy of PsbO is present in all oxygenic organisms [50,51]. In higher plants, however, persistent biochemical observations indicate that two copies of PsbO are present per monomer of the photosystem. These studies have been reviewed extensively elsewhere [21,52,53]. Briefly, this evidence falls into three main categories: protein quantification experiments, rebinding studies and mutant characterization. Direct immuno-quantification using internal PsbO protein standards indicates that two copies of PsbO are present per monomer in both PS II membranes and β-D-dodecyl maltoside-solubilized PS II particles [54]. The number of PsbO copies bound is independent of the presence or absence of an intact

Mn₄CaO₅ cluster, although the *K*_d for both PsbO copies is modulated by the intactness of the metal cluster [55]. Earlier, protein purification efforts had indicated that it was possible to isolate 1.7 mol of the PsbO component per mole PS II from PS II membranes; these membranes contain only strongly bound PsbO [56]. Rebinding studies from a number of laboratories examining higher plant PsbO binding to PS II membranes after extraction by CaCl₂ or NaCl-urea treatment demonstrated that binding of PsbO saturates at two copies of PsbO per PS II monomer [55,57–59]. The binding of native PsbO to PS II exhibits cooperativity with a Hill coefficient of 2.0 [55] and in some cases sigmoidal binding was evident [60]. Certain mutated⁷[57,61] and chemically modified PsbO proteins [62] exhibit markedly sigmoidal binding isotherms. These are critical observations. The observed sigmoidicity necessitates cooperative binding which, by definition, requires at least two binding components. Finally, analysis of N-terminal truncation mutants has identified two binding domains, both of which must be present to observe normal PsbO binding stoichiometry [34,63]. One of these domains is not conserved in cyanobacterial sequences, perhaps explaining the apparent lack of a second copy of PsbO in these organisms. Functional studies indicate that the two copies of PsbO perform different functions within the photosystem. The first bound PsbO stabilizes the Mn₄CaO₅ and lowers the chloride requirement for oxygen evolution while the second PsbO further lowers the chloride requirement and optimizes oxygen evolution capacity [64,65].

If higher plants do contain two copies of the PsbO component, where is the second copy bound? At least two models are possible. First, the second copy could be bound symmetrically with respect to the first copy, on the luminal surface of the PS II complex. This would necessitate the second copy associating principally with the CP47 protein in a similar manner as the first copy associates principally with CP43 in the cyanobacterial crystal structure. In this model, the second copy of PsbO would be quite distant from the Mn₄CaO₅ cluster. A second possibility is that the two PsbO copies form a homodimer. Three lines of circumstantial evidence support this hypothesis. First, during crystallization experiments of the spinach protein, only PsbO dimers are observed in the microcrystals which formed [66]. Second, single particle analysis of PS II supercomplexes has interpreted the density arising from the PsbO subunit as being aglobular and/or being present as a dimer [67]. Finally, re-examination of the chemical modification data presented by Frankel and Bricker [68] indicated that the majority of the PsbO residues which were labeled when PsbO was in solution but which were not labeled when PsbO was associated with NaCl-washed PS II membranes are solvent-exposed in the

⁷ It should be noted that in the only reconstitution study performed with cyanobacterial mutated PsbO proteins, sigmoidal binding was also observed for some mutants [36].

T. vulcanus crystal structure. These residues may define the interaction domain of the second copy of the PsbO protein with higher plant PS II. It should be noted that these models are not necessarily exclusive. A homodimer of PsbO would almost certainly also interact with the CP47 present in the other PS II monomer. If this were the case, it could explain a number of the discrepant results described above [46,47]. Clearly, resolution of this question of PsbO stoichiometry in higher plants ultimately will require crystal structures of intact higher plant PS II.

While the cyanobacterial and higher plant PsbO proteins are quite similar, some differences have been noted. In addition to the N-terminal truncation which is present in all cyanobacteria (see above), several loop domains are also different. These features have been examined in detail elsewhere [20,53]. The most striking of these differences is the presence of a fourteen amino acid residue “cyano-loop” [20] which is present in all cyanobacteria. In *T. vulcanus* this loop extends from ¹²⁹T–¹⁴²F while the analogous loop in spinach is quite small (¹³⁶G–¹⁴¹F). The functional significance of this difference is unclear; however, it has been noted that the cyano-loop effectively compensates for the ubiquitous cyanobacterial N-terminal truncation, so that all PsbO proteins are approximately the same length. It has been suggested that this may have functional implications [53,69].

3. PsbO function

Functionally, what parameters of the oxygen-evolving process are affected by the biochemical or genetic removal of PsbO? Numerous investigators have examined this question using higher plant PS II membranes which were depleted of the PsbO component by either CaCl₂ [70] or NaCl-urea [9] treatment. These preparations, while completely lacking the PsbO protein, retain the Mn₄CaO₅ cluster and the ability to evolve oxygen, albeit at very low rates [11]. It should be noted that these studies were typically performed at high chloride and calcium concentrations to maintain the intactness of the Mn₄CaO₅ cluster (see below) and to compensate for the absence of the PsbP and PsbQ subunits which are also removed by these treatments. Ono and Inoue [71], using thermoluminescence as a probe for S-state transitions, observed that the S₃ → [S₄] → S₀ transition was significantly slowed. Miyao et al. [72] performed flash oxygen yield measurements on PsbO-depleted PS II membranes. These studies indicated that the S₂ lifetime of the PsbO-depleted samples increased five-fold and that the lifetime of the S₃ state also increased. Additionally, the S₂ → S₃ transition was slowed and the S₃ → [S₄] → S₀ transition was retarded two- to three-fold. Similar results have been obtained *in vivo*. An *Arabidopsis thaliana* (hereafter *Arabidopsis*) mutant which lacks a functional PsbO-1 protein [59,73], the isoform responsible for normal PS II activity, exhibited long-lived S₂ and S₃ state lifetimes [74]. Thermoluminescence experiments performed with an *Arabidopsis* T-DNA mutant which lacked the PsbO-1 protein also indicated an increased stability of the S₂ state, and EPR measurements indicated a lower yield of the S₂ multiline signal [75]. The analysis of cyanobacterial *ΔpsbO* mutants has yielded very similar results. Flash oxygen yield measurements on the *ΔpsbO* mutant of *Synechocystis* 6803 showed an increased stabilization of the S₂ and S₃ states and that the S₃ → [S₄] → S₀ transition was slowed at least five-fold [76]. Thermoluminescence measurements on this strain also indicated a higher stability of the S₂ and S₃ states [76,77] in this mutant. A parallel retardation in the rate of Y₂• reduction was also observed [78] which indicated slower electron transfer from the Mn₄CaO₅ cluster to Y₂•.

Considerable circumstantial evidence is available which indicates that the PsbO protein functions to modulate the chloride and calcium⁸ requirements for oxygen evolution. The oxygen evolution requirements for both of these ions are significantly lowered in the presence of the PsbO component [11]. The effects of chloride and/or calcium

depletion on the oxygen evolution characteristics of higher plant PS II membranes are very similar qualitatively to the effects observed upon removal of the PsbO protein, as described above. This correlation is particularly strong with respect to chloride. Oxygen-evolving activity is markedly inhibited in chloride-depleted PS II membranes [79], with residual activities less than 20% of the control reported. Chloride depletion leads to a decrease in intensity of the S₂ multiline signal with a concomitant increase of the g = 4.0 signal [79,80]. A similar decrease in the multiline signal was observed upon removal of the PsbO component [81]. Flash-induced UV absorbance changes indicated a twenty-fold stabilization of the S₂ state lifetime, a retardation of the S₂ → S₃ transition and the absence of any detectable S₃ → [S₄] → S₀ transition upon chloride depletion [82]. No effect on S₃ lifetime, however, was observed. Proton release experiments indicated that S-state transitions were blocked after S₂ [83]. Ono et al. [84] reported that the XANES edge shifts associated with S-state advancement were blocked at the S₂ state. Finally, chloride was demonstrated to be required for the S₂ → S₃ and S₃ → [S₄] → S₀, but not for the S₀ → S₁ or the S₁ → S₂ transitions [82]. Depletion of calcium also prevents the S₂ → S₃ transition [85]; however, the S₁ → S₂ transition was also slowed [86,87]. These latter results may correlate to the observation that removal of the PsbO protein leads to a lower yield of the formation of the S₂-multiline signal [81]. Finally, thylakoids isolated from *Arabidopsis* mutants lacking the PsbO-1 protein exhibit long-lived S₂ and S₃ states [88]; this phenotype can be chemically complemented by the addition of calcium and a calcium ionophore but not by the addition of chloride, alone. The mechanism(s) by which the PsbO protein could modulate the chloride and calcium requirements for oxygen evolution, however, remain unclear.

The PsbO component was earlier termed the ‘manganese-stabilizing protein’ based on the observation that removal of this component from higher plant PS II membranes leads to a destabilization of the Mn₄CaO₅ cluster at low chloride concentrations. In the absence of added chloride, two of the four manganese become paramagnetically uncoupled [89] and are eventually lost to the bulk media [90,91]. In the presence of high chloride concentrations (>100 mM) the Mn₄CaO₅ cluster remains intact and oxygen can be evolved even in the complete absence of the PsbO component [11,90], although at significantly lower rates (20–40% of control). It has recently been reported that intermediate chloride concentrations (10–20 mM) preserve the integrity of the Mn₄CaO₅ cluster to varying degrees, and this is modulated in the higher plant system by the number of the PsbO subunits associated with each monomer [64]. While the PsbO protein stabilizes the active site Mn₄CaO₅ cluster, EXAFS experiments performed on PS II membranes depleted of PsbO indicated that essentially no change occurred in the structure of the metal cluster upon removal of PsbO [92]. This result strongly indicates that the PsbO protein does not provide any ligands directly to the Mn₄CaO₅ cluster. Early investigations also suggested that PsbO was directly involved in calcium binding at the oxygen-evolving site of PS II [93–95]; later studies indicated that PsbO was not involved in this process (see Seidler, [96], for an in-depth discussion). Seidler and Rutherford [97], for instance, demonstrated that in the absence of the PsbO protein, the high affinity binding site for calcium associated with the oxygen-evolving site was still present. These latter findings have been confirmed by the various cyanobacterial PS II crystal structures which indicate that the PsbO protein does not provide any ligands to either the calcium or manganese ions in the Mn₄CaO₅ cluster [5,7,8,98].

Putative low affinity calcium-binding sites, distinct from the active site metal cluster may, however, be present on the PsbO protein, although the functional significance of these is unclear. Murray and Barber [99] identified a calcium-binding site with the calcium ligands being ⁵⁴E, ¹¹⁴E and ²³¹H, which was also observed by Guskov et al. [8]. It has been speculated that this low affinity site may be associated with a putative proton exit pathway leading from the Mn₄CaO₅ cluster to the surface of the complex [7,20]. The authors note, however, that with the exception of ¹¹⁴E, these residues are rather poorly conserved

⁸ For excellent reviews on the role of chloride and calcium in PS II the reader should consult references [143,144].

[99]. This calcium-binding site was not observed in the latest crystal structure [5]; while the positions of ^{114}E and ^{231}H are similar to those observed in earlier structures [7,8,99], the side chain of ^{54}E is rotated away from the putative calcium-binding site and appears to associate with ^{57}K via a charge-pair interaction. It is unclear if calcium was lost from this position in the Umena et al. structure [5] or if this putative calcium-binding site was a possible crystallization artifact in the earlier models [7,8,99]. A different calcium-binding site on PsbO is located in the *T. vulcanus* structure [5]. The putative calcium ligands are ^{200}N , the backbone carbonyls of ^{138}T and ^{201}V and two water molecules (^{253}Wat and ^{1023}Wat). The possible physiological significance of this site is unknown. In any event, none of the amino acid residues in either of the putative calcium-binding sites have been examined by site-directed mutagenesis in either cyanobacterial or higher plant systems.

While in *Synechocystis* 6803 the absence of the PsbO protein does not prevent photoautotrophy or assembly of PS II, the ΔpsbO mutant grown on complete medium [10] grew slower than wild type, evolved oxygen at about 40% wild-type rates, and was more susceptible to photo-inactivation. This strain cannot grow under low calcium [100] or chloride [101] conditions. These observations indicate a role for the PsbO protein in the regulation of these ions at the oxygen-evolving site. Interestingly, in *Chlamydomonas reinhardtii* (hereafter *C. reinhardtii*) [102] and *Arabidopsis* [103], a profoundly different phenotype is observed. In these organisms, the absence of PsbO leads to a loss of photoautotrophic growth and the ability to assemble functional PS II reaction centers. The reason for the differences observed in the prokaryotic and eukaryotic systems is unclear at this time, although it should be noted that in the green algae and higher plants the loss of PsbO also leads to the concomitant loss of PsbP and PsbQ. In the cyanobacteria, however, the PsbU and PsbV proteins are not released upon the loss of the PsbO component. The retention of PsbU and PsbV may partially stabilize the manganese cluster in the cyanobacterial system. Another possibility is that the luminal chloride concentration is significantly higher in cyanobacterial thylakoids than in higher plants. If this were the case, then the integrity of the Mn_4CaO_5 cluster might be maintained in the cyanobacterial system even in the absence of the PsbO protein, while a putative lower chloride concentration in the higher plant thylakoids would lead to disassembly of the Mn_4CaO_5 cluster and destabilization of PS II.

While the evidence for a chloride and/or calcium regulatory function for the PsbO component is rather strong (if circumstantial), it is also possible that the PsbO protein may participate more directly in the water-oxidizing process. In an isotope editing experiment, PsbO-depleted PS II membranes were reconstituted with uniformly ^{13}C -labeled PsbO protein [104]. Fourier transform infrared spectroscopy (FTIR) was then used to monitor changes in the protonation state of the PsbO component during S state turnover. During the $\text{S}_1 \rightarrow \text{S}_2$ transition, spectra were obtained which were consistent with the deprotonation of the PsbO protein; the proton acceptor did not appear to reside on the protein. The authors hypothesized that PsbO may serve to stabilize the charged S_2 state. Interestingly, the oxidation state of the Mn_4CaO_5 cluster appears to modify the structure of the PsbO protein [105]. Treatment of PS II membranes with 100 μM hydroxylamine leads to the reduction of the Mn_4CaO_5 cluster. In the presence of the PsbO protein, the manganese and presumably calcium ions are not lost to the bulk solvent. This treatment leads to the exposure of tryptic sites on the PsbO protein. Trypsin treatment of hydroxylamine-reduced samples leads to the rapid digestion of the PsbO protein, which is normally quite resistant to trypsin attack when associated with an intact Mn_4CaO_5 cluster. This result indicates that reduction of the metal cluster leads to conformational changes in either the PsbO protein or other PS II subunits which interact with PsbO, leading to the exposure of normally trypsin inaccessible lysyl and/or arginyl residues on the PsbO protein. The locations of these residues have not been determined.

The mechanism by which the PsbO protein modulates the chloride (and possibly calcium) requirements for the photosystem is unclear. Several conserved residues on the PsbO protein have been modified by site-directed mutagenesis in spinach in an attempt to clarify this process. Modification of residues ^{151}R and ^{161}R in spinach, yielding the R151G, R151D, and R161G variants [106], lower the binding affinity of PsbO for PS II. When PsbO-depleted PS II membranes were saturated with two copies of the mutated PsbO proteins per monomer, normal sequestration of chloride was not observed and oxygen was evolved at only 20–40% of control rates. The K_{M} for chloride was 1.5–2.5 mM vs. 0.4 mM for NaCl-washed PS II membranes, which retain PsbO. No alteration for the affinity for calcium was observed. Mutations at position ^{157}D in spinach PsbO (D157N, D157K and D157E) are particularly interesting [61]. PsbO proteins with these mutations exhibit essentially normal binding characteristics when reconstituted onto PsbO-depleted PS II membranes. Nevertheless, all of these altered proteins exhibit lower affinity for chloride (K_{M} of 1.5–1.6 mM), no alteration in the affinity for calcium, and an inability to fully reconstitute steady-state oxygen evolution, with only about 50% of the control activity being observed. Interestingly, the D157E protein performs as poorly as the D157N and D157K variants. Additionally, fluorescence experiments indicated that the presence of these PsbO variants resulted in slow charge recombination in the presence of DCMU ($\text{Q}_{\text{A}}^- \rightarrow \text{Q}_{\text{B}}$ electron transfer was not affected), indicative of a more stable S_2 state. Flash oxygen yield experiments indicated that after dark incubation a higher proportion of the PS II reaction centers were in the S_3 state [107]. Consequently, these results are similar to, but not as extreme as, results obtained from PS II membranes which lack the PsbO component or have been depleted of chloride. Popelkova and Yocum have subsequently hypothesized [53] that these residues (^{151}D , ^{157}D and ^{161}R) along with the additional conserved PsbO residues ^{190}K , ^{224}D and ^{226}D , interact to properly fold a domain important for chloride retention. Examination of the interactions among the analogous residues (^{152}R , ^{158}D , ^{162}R , ^{189}R , ^{222}D and ^{224}D) in the *T. vulcanus* crystal structure [5] in large measure appears to confirm their hypothesis. PsbO residues ^{158}D and ^{162}R appear to form a charge-pair, while ^{152}R appears to hydrogen bond with the backbone carbonyl of ^{64}R of D1. ^{224}D of PsbO may participate in three relevant interactions. First, its carboxylate may hydrogen bond with the backbone amide of ^{310}E of D2; its backbone amide appears to hydrogen bond with the carboxylate of PsbO residue ^{222}D , while its backbone carbonyl forms a hydrogen bond with the backbone amide of PsbO residue ^{226}G . The backbone carbonyl of this latter residue appears to hydrogen bond with both ^{189}R and ^{184}R of PsbO. It should be noted that the domain ^{152}R – ^{162}R of PsbO closely approaches both Cl^{-1} (10.1 Å) and Cl^{-2} (12.2 Å) in the *T. vulcanus* crystal structure [5]. One can speculate that biochemical or genetic removal of the PsbO protein or site-directed modification of residues in the ^{152}R – ^{162}R domain could lead to conformational changes in intervening residues, such as residues ^{317}K of D2 and ^{334}R of D1 (including residues in the vicinity of ^{64}R which is hydrogen-bonded to ^{152}R of PsbO) at the Cl^{-1} site and/or ^{336}A – ^{338}N of D1 at the Cl^{-2} site, which would destabilize the bound chloride and result in the multitude of chloride-related defects described above.

Within PS II, access channel(s) must exist which allow the substrate water to reach the Mn_4CaO_5 cluster and exit channels must exist which allow the products, protons and molecular oxygen, to leave the active site. A number of *in silico* studies have sought to identify such channels and to determine the roles played by the various PS II subunits in their formation [20,108–110]. Recently, Gabdulkhakov et al. [111] have combined *in silico* studies using noble gases derivatization and dimethyl sulfoxide co-crystallization to identify nine possible channels in *T. elongatus*; the channels designated A1 and A2 are putative water entrance channels, channels B1 and B2 are putative oxygen egress channels and channels C–G are putative proton exit channels. A number of PsbO residues were predicted to be associated with water channel A2 and the proton exit channels C–E and G. The locations of these channels were calculated

using the 2.9 Å structure [8]. It will be most interesting to see similar analysis performed on the higher resolution *T. vulcanus* structure [5]. Determination of the actual functions performed by these putative channels will require additional experimentation.

4. PsbO as a GTPase

Evidence has been presented that the PsbO protein binds GTP with high affinity [112] and functions as a GTPase [113]. In this role, it has been hypothesized to act to control the phosphorylation state of the D1 protein [114]. Since the phosphorylation level of the D1 component appears to be coupled with its efficient PS II turnover [115,116], it was proposed that the PsbO protein may be an important regulatory component of this process [117]. In spinach, this proposed function is apparently performed by the single PsbO gene product found in this organism which also facilitates the oxygen-evolving process, as described above. The mechanisms regulating these two functions of the PsbO component are unclear. In *Arabidopsis*, two genes which encode the PsbO component (*psbO-1*, At5g66570 and *psbO-2*, At3g50820) are normally expressed, yielding two different PsbO proteins (PsbO-1 and PsbO-2, respectively). There are 11 amino acid differences between these two isoforms [73]. It has been hypothesized that the two functions ascribed to PsbO, support of oxygen evolution and regulation of PS II turnover, have differentiated between the two PsbO isoforms, with the primary function of the PsbO-1 protein being support of normal oxygen evolution by PS II and with the PsbO-2 protein principally acting to regulate the phosphorylation state and turnover of the D1 protein [114]. This suggestion was based largely on the observation that the PsbO-2 component exhibited substantially higher GTPase activity than the PsbO-1 protein while functioning poorly in support of oxygen evolution [117]. The hypothesis that the PsbO protein plays an important role in PS II turnover has been evaluated critically elsewhere [118].

5. PsbP and PsbQ

In higher plants, the PsbP and PsbQ proteins modulate the calcium and chloride requirement for oxygen evolution. In spinach, these proteins exhibit mature molecular masses of 20.2 kDa and 16.5 kDa, respectively [119]. Sequence alignments of these proteins from a variety of different organisms have been previously published [21]. Other members of the extended PsbP and PsbQ families have been found to function in PS II repair or as components of the chloroplastic NDH dehydrogenase and will not be examined in this review (for an in-depth examination of this topic, see [17–19]).

6. Structures

A number of groups have recently published crystal structures of PsbP from higher plants [120–122]. Fig. 3A illustrates the three-dimensional structure of the PsbP protein from spinach at 1.98 Å resolution (PDB ID: 2VU4, [121,122]). The structures of the spinach and tobacco (PDB ID: 1V2B, [120]) PsbP proteins and the cyanobacterial homologue CyanoP (PDB ID: 2XB3, [123]), which is shown in Fig. 3B, are quite similar although the cyanobacterial protein has very limited sequence homology with its higher plant counterparts [21]. The protein exhibits a two-strand anti-parallel β -sheet near its N-terminus, and a central anti-parallel six-strand β -sheet which is flanked by two α -helical domains. The N-terminus of the protein is disordered and is not resolved in any of the available structures. The protein contains a number of bound zinc ions, the biological consequences of which are unclear (see below). As originally pointed out by Ifuku et al. [120], the protein shares its fold with Mog1p (PDB ID: 1EQ6 [124]), a regulatory protein which interacts with the *ran* GTPase in yeast [125]. This is quite interesting as the PsbP protein is




Fig. 3. Crystal Structures of PsbP and CyanoP. A., The spinach 1.98 Å PsbP crystal structure (PDB ID: 2VU4 [122]) is shown. The lysyl residues identified by Tohri et al. [133] as being involved in electrostatic interactions with PS II are shown in blue. Please note that residue 33K is not visible in this rendering and is located behind and to the lower left of the molecule. B., The *T. elongatus* 2.8 Å CyanoP crystal structure (PDB ID: 2XB3 [123]) is shown.

known to interact with the PsbO component which, as described above, has been suggested to exhibit GTPase activity [112,113].

Several groups have published crystal structures of PsbQ from higher plants [126,127]. Fig. 4A illustrates the three-dimensional structure of the PsbQ protein from spinach at 1.49 Å resolution (PDB ID: 1VYK, [127]). The protein exhibits a four α -helix bundle in its C-terminal domain which is very similar to the fold exhibited by the cyanobacterial homologue CyanoQ (PDB ID: 3LSO, [128]), as shown in Fig. 4B. A two-strand parallel β -sheet is located in the N-terminal domain along with a polyproline type II (PPII) left-handed helix formed by four sequential prolyl residues (⁹Pro–¹²Pro) in the spinach protein; these features are absent in CyanoQ. The spinach protein contains a number of bound Zn²⁺ ions, again the function, if any, of these is unknown (see below). The loop which connects the two β -strands (¹³L–³⁴T) is not resolved in the current structure (probably because it exhibits high mobility) but has been modeled using a molecular dynamic simulation [129].

A variety of studies have examined the association of these components to the photosystem and to each other. Early studies indicated that the PsbO protein was required for PsbP binding and that the PsbP component was required for the association of PsbQ to PS II [130,131]. Relatively few studies have addressed the mechanism of PsbP and PsbQ binding to the photosystem. Ifuku and Sato [132] reported that the binding affinity of a recombinant mutant of the PsbP protein, of which




Fig. 4. Crystal Structures of PsbQ and CyanoQ. A., The spinach 1.49 Å PsbQ crystal structure (PDB ID: 1VYK [127]) is shown. The lysyl residues identified by Meades et al. [136] as being involved in electrostatic interaction with PS II are shown in blue. B. The *Synechocystis* 6803 1.8 Å CyanoQ crystal structure (PDB ID: 3LSO [128]) is shown.

19 N-terminal residues were truncated ($\Delta 19$ -PsbP), was weaker than that of native PsbP and did not exhibit saturation binding characteristics. Additionally, this truncated mutant exhibited functional defects (see below). The bound $\Delta 19$ -PsbP protein could support the binding of the PsbQ subunit, indicating that while PsbP's binding was weaker, at least a proportion of the protein was bound normally. Unfortunately, this N-terminal domain is not resolved in the current PsbP crystal structures. Using chemical modification with N-succinimidyl propionate (NSP), which modifies primary amino groups, Tohri et al. [133] identified lysyl residues on the PsbP protein, the modification of which dramatically decreased the binding of PsbP to PS II. These residues (^{33}K , ^{38}K , ^{143}K , ^{166}K , ^{170}K , and ^{174}K) were clustered at the N- and C-termini of the protein and were all located on one face of the PsbP protein (Fig. 3A). The residues ^{11}K and ^{13}K were also modified but are not resolved in the currently available structures. The authors hypothesized that a subset of these lysyl residues interact with the carboxylate residues on the PsbO protein which were identified to be required for the efficient binding of PsbP to the photosystem (see above, [134]). Interestingly, several of these residues appear to be conserved at similar positions in CyanoP (^{42}R , ^{168}K and ^{165}K).

Other investigators have attempted to identify the residues on PsbQ which are required for its association with the photosystem. Earlier work indicated that proteolytic removal of the N-terminal 12 amino acid residues of the protein abolished binding to the photosystem [135]. It was unclear, however, if this domain was sufficient to account for the interaction of PsbQ with the photosystem, and the amino acid residues responsible for the interaction were not determined. Meades et al. [136] used EDC coupled with glycine

methyl ester to modify lysyl residues on the PsbQ protein. They identified four lysyl residues (^{90}K , ^{96}K , ^{101}K , and ^{102}K) whose modification also abolished high affinity binding of PsbQ to PS II. These residues all clustered on one face of the protein and are in the vicinity of the N-terminus (Fig. 4A).

In *C. reinhardtii* PS II, evidence is available for the direct interaction of PsbP with PsbQ. Enami and coworkers, using the 0-length crosslinker EDC, identified a crosslinked product formed between PsbP and PsbQ [43]. ^{176}K of the PsbP protein was identified as being crosslinked to ^{28}D of PsbQ. It should be noted that no EDC-crosslinked products have been identified between PsbP and PsbQ in higher plants. This may be due to the fact that ^{28}D is not a conserved residue in higher plant PsbQ. ^{176}K of *C. reinhardtii* is analogous to residue ^{174}K in spinach, which was identified in the NSP-labeling experiments described above [133]. Additionally, Nagao et al. [43] showed immunologically that PsbP and the α subunit of cytochrome b_{559} (PsbE) were crosslinked with EDC, although the interacting residues were not identified. If this assignment is correct, then PsbP is bound to the opposite side of the PS II monomer from the PsbO protein. A direct interaction of PsbP with both PsbE and PsbO would require that PsbP occupy a position roughly similar to that occupied by PsbU and PsbV in the cyanobacterial crystal structure. This, however, would appear unlikely since the red algae contain stoichiometric amounts of PsbP, PsbQ', PsbU and PsbV [15,16] and, presumably, the PsbU and PsbV proteins in the red algae are bound to PS II in analogous locations to the PsbU and PsbV proteins in the cyanobacteria.

While most investigators believe that the standard model is correct (PsbO binding to the PS II core, PsbP binding to PsbO and PsbQ binding to PsbP), it should be noted that some results indicate that the association of PsbP and PsbQ with the photosystem may be more complicated and that the accepted model may need modification. Evidence has been presented indicating that green algal PsbP and PsbQ can bind independently of the PsbO component to the photosystem [137,138].

Additionally, three groups working in higher plant systems have reported that the PsbO protein can be removed from PS II while PsbP and PsbQ remain bound to the photosystem. Differential extraction experiments using either a butanol–water phase separation system [139] or HgCl_2 [140,141] indicated that PsbO could be removed from PS II preparations without the concomitant extraction of the PsbP and PsbQ proteins. In the latter studies, a moderate chloride concentration inhibited the release of the PsbO component. Unfortunately, none of these investigators demonstrated that the PsbP and PsbQ proteins were bound normally after HgCl_2 treatment and could subsequently be extracted by standard procedures (*i.e.* 1 M NaCl treatment). It is possible that PsbO is required for the initial binding of the PsbP component but that upon binding, conformational changes occur within the intrinsic components of PS II which allow the selective removal of PsbO while PsbP and PsbQ are retained [21]. In other studies, Caffarri et al. [142] used image analysis of detergent-solubilized (α -D-dodecyl maltoside) PS II particles (from several *Arabidopsis* mutants) which were resolved by sucrose density gradient ultracentrifugation, to examine the association of the light-harvesting antennae proteins with the PS II core complex. They reported that, while PsbO is present in all PS II core-containing particles, PsbQ was present only in supercomplexes which contained Lhcb and that PsbP binding required CP26/LHCII. Unfortunately, the majority of these experiments were performed at pH 7.5, a pH known to destabilize the oxygen-evolving complex. Additionally, the functional integrity of their preparations is difficult to determine since their oxygen evolution experiments also were performed at pH 7.5 and at sub-saturating light intensities. Nevertheless, it is possible that removal of subsets of the light-harvesting apparatus might induce conformational changes in the PS II core proteins which could alter the binding characteristics of either PsbP or PsbQ to the photosystem.

7. PsbP and PsbQ functions

Many early studies have demonstrated that the PsbP and PsbQ proteins function to optimize oxygen evolution at physiological concentrations of calcium and chloride. These findings have been reviewed in detail elsewhere [21,96,143,144]. Briefly, removal of the PsbP and PsbQ proteins by NaCl treatment of PS II membranes results in a marked loss of oxygen evolution activity (to about 25% of control). Much of the lost activity can be restored by reconstitution of the extracted protein(s) to protein-depleted PS II [90,145–147], but only if the calcium and chloride cofactors are present during the reconstitution [90,148,149]. It should be noted that high, non-physiological concentrations of calcium and chloride can support reasonably high rates of oxygen evolution even in the absence of the PsbP and PsbQ components [90,148,150]. Inhibition of oxygen-evolving activity after extraction of the PsbP and PsbQ proteins is accelerated in the light. This appears to be due to differential binding affinities for both calcium [151,152] and chloride [153,154], which are modulated by the S-state of the oxygen-evolving complex. Depletion of the PsbP and PsbQ proteins also markedly increases the accessibility of the Mn_4CaO_5 cluster to exogenous reductants [155].

While phenomenologically the PsbP and PsbQ proteins optimize oxygen evolution under physiological calcium and chloride concentrations, the mechanism(s) of such optimization remain largely unknown. At least two possibilities exist. First, the proteins could serve to stabilize a conformation of PS II required for the normal integration of these ions into the oxygen-evolving complex. This would imply that other components of PS II exist in different conformational states which are dependent on the presence or absence of the PsbP and/or PsbQ proteins. Direct evidence for such conformational changes exists. Removal of these proteins appears to introduce alterations in the function and structure of the PS II complex. Functionally, genetic or biochemical removal of PsbP (with the concomitant removal of PsbQ) leads not only to alterations in oxygen evolution capability but also to alterations in electron transport from Q_A^- to Q_B [156,157]. No reducing-side defects are associated with the removal of PsbQ alone [158], at least in plants grown under normal growth light conditions. Since both the Q_A and Q_B sites reside on the stromal side of the membrane and PsbP is bound to the luminal side, these results indicate that transmembrane conformational changes occur upon the removal of PsbP. Image analysis of PS II supercomplexes from higher plants indicates that removal of PsbP and PsbQ induces CP29 to move approximately 12 Å towards the central core of the PS II complex [67]. This result indicates that the structural organization of the core proteins of PS II is altered by the removal of PsbP and PsbQ. The most likely candidates for such conformational changes are the D1 and/or D2 proteins, which bear the Q_A and Q_B -binding sites. Conformational changes have also been observed directly upon PsbP binding to PS II. In an FTIR study using ^{13}C -labeled PsbP, Tomita and coworkers [159] specifically probed for conformational changes in PS II arising from PsbP binding [160]. Conformational changes occurring in the vicinity of the oxygen-evolving complex were identified with no identifiable changes being observed in the PsbP protein itself. When reconstitution was carried out with an N-terminally truncated PsbP ($\Delta 15$ -PsbP), the conformational changes were not observed. Previously, it had been shown that the $\Delta 15$ -PsbP truncated protein, while able to bind to the photosystem, could not restore normal calcium and chloride binding upon reconstitution onto PsbP-depleted PS II membranes [161]. These results indicate that the N-terminal 15 amino acid residues of PsbP are critical in supporting normal calcium and chloride function in the photosystem. Consequently, there is a correlation between the absence of induced conformational changes near the oxygen-evolving complex, the inability to restore oxygen evolution at physiological calcium and/or chloride concentrations and the absence of the N-terminal 15 amino acid residues of PsbP. Unfortunately, these

residues are not resolved in the current crystal structures [120–122]. It should be noted that this view is complicated by the observation that PsbQ may act in a coordinated manner with PsbP with respect to calcium sequestration. This hypothesis had been suggested previously, although without direct evidence [152,162]. The N-terminal truncation of 19 amino acid residues from PsbP yields a protein ($\Delta 19$ -PsbP) which exhibits non-saturating binding characteristics and the complete loss of the ability to lower the calcium requirement for oxygen evolution [132]. If, however, native PsbQ protein is subsequently reconstituted onto $\Delta 19$ -PsbP protein-containing PS II membranes, much of the oxygen evolution rate is recovered, even in the absence of added calcium. So, calcium retention can be restored in the presence of a truncated PsbP protein and, presumably, the absence of conformational alterations in the vicinity of the oxygen-evolving site. If the conformational changes observed are directly related to the ability to retain calcium, then similar conformational changes should be observed after reconstitution of PS II with the truncated PsbP protein and PsbQ. The possibility of such conformational changes has not been examined experimentally.

A second possibility is that the PsbP and/or the PsbQ components actively sequester the calcium and chloride cofactors required for active and stable oxygen evolution. While no direct evidence for this possibility exists, some circumstantial evidence is available, at least with respect to the possibility that the PsbP protein can bind divalent cations like calcium. The high resolution crystal structures of both PsbP and CyanoP contain bound zinc ion(s). In tobacco and spinach PsbP [121], the zinc appears to be ligated to ^{144}His and ^{165}Asp , which also corresponds to one of the four zinc-binding sites observed in CyanoP [123]. It is unclear at this time if this association is physiologically relevant or is an artifact. Three possibilities exist. First, the bound zinc could be physiologically relevant. Trace metal analysis of PS II membrane preparations from higher plants indicated that PS II membranes which contained 4 atoms of manganese per PS II also contained 1–2 atoms of zinc per PS II [163,164]. The physiological consequences of the removal of this zinc from PS II have not been thoroughly investigated, although it should be noted that zinc is an inhibitor of PS II activity in both higher plants and cyanobacteria [165,166]. A second possibility is the PsbP protein contains a physiologically relevant divalent cation binding site which, during crystallization [123], becomes occupied by a zinc atom. The most likely ions for such an interaction would be either calcium or manganese. As noted earlier, PsbP appears to function in regulating the calcium and chloride requirements for optimal PS II activity. If, *in vivo*, a calcium ion was bound to this site it could conceivably donate this ion during formation of the Mn_4CaO_5 cluster or alter the calcium equilibrium in favor of bound calcium within the active site. Calcium was not present during the crystallization process in any of these studies. In this regard, it should be pointed out that it has been hypothesized that PsbP is a manganese-binding protein which delivers manganese to the oxygen-evolving site during photoactivation. This hypothesis has been critically evaluated elsewhere [118]. Finally, given the high zinc concentrations (5 mM in the case of the spinach protein [121], 200 mM in the case of CyanoP [123]) present during the crystallization of PsbP, it is possible that the zinc binding is purely an artifact of the crystallization process. Analogous arguments can be made concerning the observed bound zinc in the PsbQ and Cyano Q structures (see below).

Both PsbP and PsbQ may be phosphorylated in *Arabidopsis*. While no phosphorylation has been reported using standard radiolabeling procedures in any system, two recent proteomic studies have identified putative phosphorylated peptides from these two components after IMAC or TiO_2 enrichment for phosphopeptides [167,168]. PsbP-1 and PsbP-2 were reported to contain multiple phosphorylation sites (^{56}T , ^{66}T , ^{70}S and ^{82}S of the mature protein). While phosphorylated PsbQ-1 peptides were not identified, phosphorylation of the mature PsbQ-2 protein at ^{91}S of the mature protein was detected. Both PsbP and PsbQ

phosphorylation was detected under both dark and light incubation conditions. The protein kinases and phosphatases associated with the observed phosphorylation have not been identified and the biological role played by the phosphorylation remains undetermined.

Finally, the PsbP and PsbQ proteins appear to function as assembly and/or stability factors for PS II in higher plants.⁹ These findings result from RNAi suppression experiments used to examine the effects of the loss of PsbP in tobacco [169] and *Arabidopsis* [156]. The results obtained in these two experimental systems are quite similar. Moderately strong suppression of PsbP (5–10% wild-type levels) leads to slow photoautotrophic growth and a variety of functional defects associated with both the oxidizing and reducing sides of the photosystem [156,169,170]. The PS II reaction center proteins (CP47, CP43, D1 and D2) accumulate to relatively normal amounts, with only modest decreases being observed and with the accumulated proteins assembling into PS II dimers. LHC II–PS II supercomplexes, however, do not form and the reaction center proteins and the LHC proteins are not phosphorylated [170]. Significant defects in chloroplast and thylakoid architecture are also apparent in these plants [170,171]; however, the mechanism(s) giving rise to these alterations are unknown. Under stronger suppression of the PsbP protein (<1% wild-type levels), a significantly different phenotype is observed. While, as expected, the various PS II functional alterations are exacerbated and the ability to grow photoautotrophically is lost, the mutant's ability to accumulate the PS II core proteins CP47 and D2 is markedly reduced [156] and the thylakoid membranes are dramatically altered [171]. It is interesting to note that the loss of the reaction center components does not parallel the loss of PsbP; even low levels of PsbP (5–10% of wild type) support the near normal accumulation of these intrinsic components. It is only under severe depletion of PsbP that loss of the reaction center components is observed. In this respect, it is possible to hypothesize that PsbP may possess some “catalytic” function with respect to PS II assembly. The suppression of PsbP expression, while not affecting the accumulation of PsbO, leads to the complete loss of the PsbQ component, even in plants containing detectable levels of PsbP [156,169]. It is unclear at this time whether the loss of the PsbQ protein is due to decreased synthesis, inefficient transport or increased degradation. It should be noted, however, that large pools of unassembled mature PsbO, PsbQ and PsbP proteins normally exist in the thylakoid lumen without being degraded either in the presence [172,173] or in the absence [174,175] of assembled and functional PS II reaction centers. Consequently, the loss of the PsbQ component must be the direct result of the loss of expression of the PsbP protein and not a consequence of the reduced levels of PS II reaction centers observed when the PsbP component is strongly suppressed.

The PsbQ protein has also been examined using RNAi suppression techniques in both tobacco [169] and *Arabidopsis* [158]. Under normal growth light conditions, the complete suppression of PsbQ expression did not affect photoautotrophic growth, PS II assembly, or most PS II functional parameters (oxygen evolution rate, F_v/F_m , etc.) in either system. In *Arabidopsis* some modest defects were observed with respect to S-state distribution and the stability of the oxygen-evolving complex under dark incubation conditions [158]. A major alteration in phenotype was observed, however, when *Arabidopsis* mutants lacking the PsbQ proteins were placed under low light conditions ($5 \mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$). After three weeks these plants yellowed and died while wild-type plants were unaffected. The PsbQ-deficient plants lost the ability to assemble functional PS II reaction centers, with major losses of the CP47, CP43 and D2 components being observed. Interestingly the D1 protein was not seriously affected. The functional characteristics of the PS II reaction centers which did assemble were also seriously compromised [158]. A more detailed

examination of the assembly characteristics of PsbP- and PsbQ-deficient higher plants and a comparison between the higher plant, *C. reinhardtii* and cyanobacterial model systems is presented elsewhere [118].

8. PsbR

The PsbR protein is the most enigmatic of the extrinsic proteins associated with higher plant and green algal PS II; genes encoding this component are not found in the genomes of cyanobacteria or red algae. PsbR was first identified in the PS II complex of spinach [176]. Higher plants utilize a poorly understood targeting mechanism for this component. PrePsbR, which is translated on cytosolic ribosomes, has a very short transit sequence (only 41 amino acid residues in both *Arabidopsis* and spinach) relative to the other nuclear-encoded extrinsic proteins of the PS II complex. For instance, the PsbO-1 protein in *Arabidopsis* contains an 85 amino acid residue bipartite transit sequence. It has been suggested [177] that the shortened N-terminal sequence acts to target PsbR to the chloroplast stroma while a C-terminal hydrophobic domain of the protein functions as a non-cleavable signal for luminal transport. This domain would subsequently act as a hydrophobic anchor of the PsbR protein to the lumenal side of the thylakoid membrane, although this hypothesis has not been tested experimentally. The hydrophobic nature of the C-terminus of the PsbR protein apparently makes this component relatively insoluble and the protein tends to precipitate during isolation [178]. Interestingly, the protein is removed from PS II membranes by alkaline-Tris washing of PS II membranes [176,178]. This almost certainly indicates that the hydrophobic C-terminal domain is not organized as a transmembrane α -helix since proteins with a single transmembrane helix, such as cytochrome *f*, are not removed by this treatment.

It is unclear which subunits are the nearest neighbors of PsbR within the PS II complex. Since PsbR is not present in cyanobacterial PS II, no crystallographic information is available. The only experimental structural information is from the analysis of protein crosslinking experiments [179] using 3,3'-dithiobis[sulfosuccinimidylpropionate], a cleavable crosslinker which spans 12 Å. In this study, CP47 formed a crosslinked species with an unidentified 10 kDa polypeptide, yielding a 54–56 kDa product. The authors tentatively identified the 10 kDa component as the PsbR protein due to its apparent molecular weight and the disappearance of the crosslinked species when the PS II membranes were Tris-washed prior to the initiation of the crosslinking experiment. PsbR appears to be required for PsbP assembly into the PS II complex. Additionally, PsbJ, a low molecular mass plastid-encoded hydrophobic subunit, appears to be required for the stable assembly of PsbR [180] and, consequently, PsbP [181,182]. One possible model would be that PsbJ serves to anchor the PsbR protein to the PS II complex (possibly by interacting with the hydrophobic C-terminus of PsbR) and that PsbR participates along with PsbO in the formation of a binding domain for PsbP. If this model is correct, PsbR would be positioned at approximately the location of PsbV in cyanobacteria¹⁰ [5].

In the absence of crystallographic information, I-TASSER was used to predict the three-dimensional structure of the PsbR protein [183,184]. I-TASSER was the best performing structural prediction software platform on the CASP-7, CASP-8 and CASP-9 three-dimensional structural prediction trials. The top two models provided by I-TASSER are shown in Fig. 5. While such models must be considered tentative, in all of the predicted structures examined (top five models) the C-terminal domain is predicted to be organized as a hydrophobic helix-coil-helix motif. The longest helical domain averaged about fourteen amino acid residues in length. It is unlikely that such a structure could span the thylakoid membrane and it is

⁹ In cyanobacteria the loss of CyanoP or CyanoQ does not dramatically affect PS II assembly or stability.

¹⁰ In this regard it should be noted that the closest approach of PsbV to CP47 in cyanobacteria is less than 12 Å [5]. If PsbR occupies nearly the same position in higher plants, this could be consistent with the crosslinking evidence presented in [179].

possible that this anchors PsbR to the membrane (or to hydrophobic protein subunits such as PsbJ), which would be consistent with its release by alkaline–Tris treatment [176,178].

The function of the PsbR protein *in vivo* has been investigated using mutant plants in both potato [185] and *Arabidopsis* [180,186,187]. In these reports, the absence of the PsbR protein was reported to lead to impaired oxygen evolution and Q_A^- reoxidation. These defects may be directly related to the inability of the PsbP proteins to bind to PS II in the absence of the PsbR protein. In initial studies, the absence of PsbR did not appear to affect the accumulation of other PS II components. Specifically, in whole leaf extracts, the amounts of immunologically detectable PsbP and PsbQ proteins were not affected [185]. Later studies, however, examining thylakoids [180,186] or PS II membranes [187] isolated from mutants lacking the PsbR component, indicated that the PsbP and PsbQ proteins were strongly depleted. It was hypothesized that the absence of the PsbQ component was a secondary effect subsequent to the failure of the PsbP protein to bind in the absence of PsbR [180]. Similar electron transport defects have been documented in PS II which has been depleted of the PsbP component either genetically [156] or biochemically [157].

In *C. reinhardtii*, the PsbR protein is differentially phosphorylated [188]. Under State 1 conditions (aerobic dark incubation) no phosphorylation is observed while under State 2 conditions (anaerobic dark incubation) or incubation under moderate or high light, the protein is phosphorylated. The phosphorylation site is ^{43}S of the initial translation product (residue ^{15}S of the mature protein). In higher plants, this phosphorylation site is not conserved and using standard ^{32}P radiolabeling techniques, PsbR is not observed to be phosphorylated. Recently, however, two proteomic studies examining the phosphoproteome of *Arabidopsis* have detected putative PsbR phosphorylation using MS/MS techniques. The protein was reported to be phosphorylated under both light and dark conditions at ^{58}S (residue ^{30}S of the mature protein) [167,189]. The identity of the protein kinase and phosphatase associated with this phosphorylation site are unknown [190]. The function of the differential phosphorylation in *C. reinhardtii* and the possible phosphorylation in higher plants is unknown.

9. PsbU

9.1. Structure

The X-ray-derived crystal structure of bound PsbU has been determined in *T. vulcanus* [5,191] and *T. elongatus* [7,8,98]. The structure of PsbU from *T. vulcanus* at a resolution of 1.9 Å is shown in Fig. 1 in blue and possesses six alpha helices connected by five loops with little or no β -sheet structure being recognizable. No functional domains or other structural homology exist to other proteins in the current databases. The interacting partners in the photosystem include CP47, PsbO and PsbV.

PsbU was first identified in the cyanobacterium *Phormidium lamosum* as an ~9 kDa polypeptide [192] and was subsequently shown to be a stoichiometric component of cyanobacterial PS II [193]. The first report of this protein in a eukaryotic alga was in isolated PS II preparations from the red alga *Cyanidium caldarium* (henceforth, *C. caldarium*) [194]. Unlike PsbV, PsbU is nuclear-encoded in the red algae [195,196]. The PsbU subunit has been identified in most cyanobacteria, with the exception of *Prochlorococcus marinus* strains MED4 and SS120, although it is present in *P. marinus* MIT9313. This subunit has also been detected in chlorophyll *a/c*-containing algae [197,198]. In contrast, PsbU is absent from all chlorophyll *a/b*-containing eukaryotes.

The binding of PsbU to PS II was initially studied in isolated PS II complexes from *T. vulcanus* [193,199]. The PsbU protein was not able to bind to PS II in the absence of PsbO and PsbV. PsbU was observed to bind at low levels if only one of these polypeptides was present [199].

PsbU bound at approximately 40% its native level when only PsbV was present and 25% of its native level when only PsbO was present. Furthermore, oxygen evolution by isolated PS II complexes in which the three extrinsic proteins had been removed by a $CaCl_2$ treatment was greatly reduced and required reconstitution with PsbO, PsbV and PsbU to fully restore activity. Partial restoration of activity in the absence of PsbU, however, was observed [199].

Gene deletion and gene interruption studies with *Synechocystis* 6803 found the $\Delta psbV$ mutant and the $\Delta psbV:\Delta psbU$ double mutant had indistinguishable phenotypes, both consistent with PsbU requiring the presence of PsbV to functionally associate with PS II [200,201]. Interestingly, while the $\Delta psbV:\Delta psbU$ double mutant was able to grow photoautotrophically, the $\Delta psbO:\Delta psbU$ double mutant was not — at least when the cells were grown at pH 7.5. Photoautotrophy could be restored to the $\Delta psbO:\Delta psbU$ double mutant when cells were grown at pH 10 [201]. Although the mechanistic basis for this pH-dependent growth has proved difficult to pinpoint [202] it does appear consistent with the *in vitro* binding studies performed with *T. vulcanus*, since in those reconstitution studies, PsbU bound with twice the affinity to complexes having only PsbV compared with complexes having only PsbO [199].




Fig. 5. Putative structures for PsbR. The structure of PsbR was modeled using the program I-TASSER [183,184]. The top two candidate structures are illustrated. Both exhibited a helix–coil–helix motif at their C-termini. The longest helical segment averaged 12–15 amino acid residues in length in five candidate structures examined. This domain is probably too short to span the thylakoid membrane.

9.2. Function

Synechocystis 6803 cells lacking PsbU are photoautotrophic when grown under standard laboratory conditions in BG-11 growth medium and assemble PS II centers at a similar level to wild type [200,203,204]. While photoautotrophic growth rates of the mutant and wild type are similar, the $\Delta psbU$ cells have a more stable S_2 state [200,204,205], although the consequences of this for the viability of the cells are not immediately obvious. In the absence of PsbU, oxygen evolution rates are reduced by typically 20% or more relative to wild type and photoinactivation is observed during steady-state oxygen evolution measurements. This is consistent with the observation that $\Delta psbU$ mutants are susceptible to photodamage when exposed to high light, exhibiting a more rapid degradation of the D1 protein [200,203,204,206–208].

In addition to the above PS II-specific perturbations, the $\Delta psbU$ mutants are more susceptible to heat stress. Heat-inactivation of PS II is known to involve the loss of manganese from the oxygen-evolving complex [209]. Addition of PsbU to isolated thylakoid membranes from *Synechococcus* sp. PCC 7002 (henceforth *Synechococcus* 7002) that had been depleted of this subunit increased the heat stability of oxygen evolution while inactivation of *psbU* in *Synechococcus* 7002 prevented the ability of these mutants to acclimate to high temperature. In the absence of PsbU, the cells were unable to acquire cellular thermotolerance [210–212]. A similar result was obtained with a $\Delta psbU$ strain of *Synechocystis* 6803, although in this cyanobacterium the removal of PsbU only partially prevented acclimation to high temperature, whereas cells lacking PsbO or PsbV failed to develop cellular thermotolerance when grown at moderately high temperatures [206]. These results indicate that the stability of PS II, conferred by the lumenal extrinsic proteins, is important for cells to acquire thermotolerance by optimizing specific protein–protein or protein–lipid interactions; however, the mechanism is as yet unidentified.

Interestingly, removal of PsbU in *Synechocystis* 6803 also resulted in uncoupling of energy transfer from the phycobilisome to the photosystem. $\Delta psbU$ cells exhibited a decreased chlorophyll *a* variable fluorescence yield when excited with light absorbed by allophycocyanin but not when excited by light absorbed primarily by chlorophyll *a* [203,207]. Furthermore, picosecond fluorescence decay kinetics revealed changes in both allophycocyanin- and PS II-associated decay components, consistent with a decrease in the coupling of phycobilisomes to PS II and an increase in the number of closed PS II reaction centers in the dark-adapted $\Delta psbU$ mutant [207]. Thus removal of PsbU introduces PS II-specific structural changes on both sides of the thylakoid membrane.

Inactivation of PsbU has also been studied in *Synechococcus* sp. PCC 7942 (henceforth, *S. elongatus*) [208]. These experiments demonstrated an increased rate of D1 degradation under both low and high light conditions. In addition, the recombination reaction between the reduced primary quinone acceptor Q_A^- and the S_2 state in the presence of DCMU was inhibited by up to 5-fold in the mutant, which is perhaps consistent with the long-lived S_2 state observed in the $\Delta psbU$ mutant of *Synechocystis* 6803 [200,204]. However, the thermoluminescence signal in the $\Delta psbU$ mutant of *S. elongatus* was greatly diminished compared to that observed in *Synechocystis* 6803 [199]. This result is difficult to interpret and could possibly be a result of uncoupling of the phycobilisome from the reaction center, as reported to be the case in *Synechocystis* 6803 $\Delta psbU$ cells [207] or irreversible inactivation of PS II in the *S. elongatus* mutant at the sub-zero temperatures utilized in the thermoluminescence assay. The suggestion that the observed thermoluminescence signal might arise from the loss of additional protein subunits in the mutant seems less likely since oxygen evolution rates were reported to be similar in both the $\Delta psbU$ mutant and wild type [208]. However, the most striking result reported for the $\Delta psbU$ *S. elongatus* strain was the viability of the mutant cells when challenged by H_2O_2 or methyl viologen. Compared to wild type,

the mutant exhibited greater resistance and appeared to up-regulate antioxidant enzymes. This study suggests that perturbation of PS II activity by removing PsbU may increase the production of reactive oxygen species. Consequently these mutants may be more resistant when challenged by an external oxidative stress because their endogenous protective mechanisms are already activated [208].

The requirements for the chloride and calcium cofactors in PS II are modulated by the extrinsic proteins of the photosystem [213]; the effects of omitting these ions have been investigated in $\Delta psbU$ strains of *Synechocystis* 6803. Photoautotrophic growth in calcium-limiting medium was similar for wild type and the $\Delta psbU$ mutant; however, under chloride-limiting conditions, growth was retarded in the $\Delta psbU$ strain. Additionally, photoautotrophic growth was abolished for the $\Delta psbU$ strain when both Ca^{2+} and Cl^- were omitted from the growth medium [39,200,204]. In the 1.9 Å resolution structure of PS II from *T. vulcanus* two chloride ions have been identified in the vicinity of the Mn_4CaO_5 cluster of the oxygen-evolving complex and may contribute to the maintenance of the coordination environment of the cluster [5]; hence it is possible that structural rearrangements resulting from removing PsbU might affect the integrity of these chloride-binding sites.

The two chloride binding sites identified in PS II from *T. vulcanus* are positioned at the start of hydrogen bond networks arising from the Mn_4CaO_5 cluster and extending towards the bulk solution. The network through the Cl^-1 site is located in the interface of the D1, D2 and PsbO subunits, while the Cl^-2 site is located at the interface of D1, CP43 and PsbU subunits. These channels may function as proton exit channels or water inlet channels and hence removing PsbU may disrupt these processes, which in turn may contribute to the phenotype of the $\Delta psbU$ mutants. The presence of a putative oxygen exit channel in the vicinity of PsbU in *T. vulcanus* is shown in Fig. 6 and would correspond to the putative B2 oxygen exit channel suggested by Gabdulkhakov et al. [111]. The channel is formed by contributions from the D1, D2, CP43, CP47, PsbU and PsbV subunits with the mouth of the channel composed of residues from PsbU, PsbV and D2.

A similar role for PsbU has been suggested following analysis of the X-ray-derived PS II structure from *T. elongatus* at 2.9 Å resolution [111]. Theoretical calculations suggested the presence of several channels potentially associated with water access, proton exit and oxygen exit. The two channels suggested to play a role in oxygen exit received experimental support by co-crystallization with the noble gas krypton as well as dimethyl sulfoxide, which were both shown to localize to the putative channels. One of the exit channels was blocked by the C-terminal lysyl residue of PsbU (^{134}K in PsbU from *T. elongatus*) and the authors proposed that the C-terminus of PsbU might operate in a gating mechanism for this channel. While the lysyl residue at this position is not absolutely conserved, an arginyl residue is frequently present that would preserve the required hydrogen bonding network; however, some PsbU sequences do not retain a positive charge at their C-terminus, and therefore the significance of this proposal requires further experimental investigation. Indeed, mutation of this terminal lysyl residue *in vitro* employing PS II isolated from the red alga *C. caldarium* did not produce any detectable phenotype [214].

Although the C-terminal lysyl residue of PsbU is not always present, the adjacent tyrosyl residue (^{103}Y) is conserved in both prokaryotic and eukaryotic organisms. Mutation of this residue dramatically altered the calcium and chloride requirements for oxygen evolution and resembled a PsbU-minus phenotype even though the binding of PsbU was unaffected. Moreover, only phenylalanine can partially substitute for tyrosine at this position suggesting that an aromatic group is required [214,215]. Okumura and co-workers have proposed that ^{103}Y interacts with ^{340}P located near the C-terminus of the D1 protein and it is the disruption of this interaction that results in the elevated requirement for calcium and chloride in the absence of PsbU or upon mutation of the tyrosyl residue [214,215]. It is therefore plausible that the pleiotropic effects




Fig. 6. The presence of a putative oxygen exit channel in the vicinity of PsbU in *T. vulcanus* (PDB ID: 3ARC). The channel (gray) was predicted using the CAVER program [255,260]. Waters and a chloride (green) from the PDB ID: 3ARC file that are located in the channel are depicted as spheres. The calcium (yellow), manganese (purple) and oxygens (red) of the Mn_4CaO_5 cluster are shown as spheres. In addition the position of the conserved ^{103}Y :PsbU residue is shown. The hydrogen bond network connecting ^{103}Y :PsbU (PsbU:Y103) with ^{320}R :PsbC (CP43:R320) is indicated (dotted black lines). See text for additional details.

of removing PsbU from PS II manifest themselves through structural changes that include alteration of the chloride-binding site(s) which, in turn, can perturb the environment of the Mn_4CaO_5 cluster, as well as altering additional protein–protein or protein–lipid interactions within the photosystem. Additionally, ^{103}Y may indirectly interact with ^{320}R of CP43. ^{320}R :CP43 hydrogen bonds with ^{99}N :PsbU (moreover a second hydrogen bond between these two residues is also apparent via an intervening water: see Fig. 6) and this residue forms a hydrogen bond with ^{103}Y :PsbU. ^{320}R :CP43 also hydrogen bonds with ^{49}N :PsbV. Within this context, mutation of the analogous *Synechocystis* 6803 CP43 lysyl residue (^{305}R) yields a mutant (R305S) which, while growing at a normal rate in complete medium, exhibited a loss of oxygen evolution capability (70% of control), assembled 30% fewer PS II reaction centers and was more susceptible to photo-inactivation [216]. Under chloride-limiting conditions this mutant grew very slowly, evolved oxygen at only 20% of the control rate, and exhibited a long-lived S_2 state [217]. PS II particles isolated from this mutant lost the ability to bind PsbV [218] and PsbU (T. M. Bricker and C. Putnam-Evans, unpublished). These results indicate that this CP43 arginyl residue may be involved in the formation of a high-affinity binding site for both PsbU and PsbV.

Isolated PS II from *C. caldarium* has been used extensively to investigate the role of PsbU and this has established that the binding of PsbU moderates the requirements for calcium and chloride by minimizing the concentrations of these cofactors that are required for maximum activity [14,214]. Hence the requirements for calcium and chloride upon removal of PsbU in red algae and cyanobacteria are similar; however, while the PsbQ orthologue (PsbQ'), PsbU and PsbV from *C. caldarium* are able to bind to *T. vulcanus* PS II, leading to restoration of oxygen evolution, the cyanobacterial PsbU and PsbV from *T. vulcanus* only partially rebind to *C. caldarium* PS II and are unable to restore PS II activity [219]. Therefore, while the extrinsic protein environments of red algae and cyanobacteria are much alike they are not identical. Nevertheless, red algal PsbU can functionally bind to PS II from *T. vulcanus* in the presence of either the cyanobacterial or red algal PsbV proteins, although cyanobacterial PsbU cannot reconstitute red algal PS II even in the presence of cyanobacterial PsbV [220].

10. PsbV

PsbV, also referred to as cytochrome c_{550} , is a PS II luminal extrinsic subunit found in many cyanobacteria and red algae. It is a c-type monoheme cytochrome with a molecular weight of approximately 15 kDa and an acidic isoelectric point. The existence of a low-potential cytochrome in cyanobacterial species had long been known [221], but PsbV was first identified as a stoichiometric PS II component

in isolated core complexes from *T. vulcanus* [222]. Since then, PsbV has been the focus of extensive research aimed at understanding its contribution to PS II oxygen-evolving activity given the unusually low midpoint potential (E_m) of its heme group.

10.1. Structure

A number of structural studies have been performed on soluble PsbV from different cyanobacteria including *Synechocystis* 6803, *Arthrospira maxima*, and *T. elongatus* [223–225]. The PsbV heme is covalently bound to two cysteinyl residues (^{37}C and ^{40}C) and non-covalently coordinated by two histidyl axial ligands (^{41}H and ^{92}H) [223–225]. This bis-histidyl ligation scheme is somewhat unusual for c-type cytochromes. However, the rest of the protein structure is quite typical of other cytochromes, with a hydrophobic inner core of the protein forming a pocket for the prosthetic group and generally protecting it from the solvent, with the exception of an exposed edge including rings C and D along with the D propionate group.

PsbV has also been structurally resolved as a component of PS II crystal structures (see Fig. 1, green subunit) from thermophilic cyanobacteria [5,7,8,98,191], and the soluble and PS II-bound PsbV protein structures are quite similar. Within the context of cyanobacterial PS II, the PsbV protein is located near luminal loop regions of the CP43 (residues 386–397 and 412–417) and D1 (residues 305–315) proteins [5]. The ^{320}R residue of CP43 also comes to within 3 Å of residues ^{49}N and ^{51}S on PsbV. This is significant because site-directed modification of the analogous residue in *Synechocystis* 6803 (CP43 ^{305}R) exhibited markedly reduced PsbV binding to PS II [218]. In the current PS II structure calcium and chloride ions can each be observed at the periphery of PsbV, but the closest residues to these ions on PsbV are more than 4 Å away. The C and D rings of the heme group are shielded by the large luminal loop region of the CP43 protein, but there is a small opening in the protein surface that exposes the propionate oxygen of the D ring. The closest edge of the heme group is approximately 23 Å away from the nearest manganese atom of the Mn_4CaO_5 cluster, which is the nearest redox cofactor to the PsbV heme in PS II [5]. Residues of the PsbV protein (^{137}Y and ^{129}K) have also been implicated in a hydrogen bond network linking Y_z and the bulk solvent on the luminal side of the complex. This channel, formed by residues of the D1, CP43 and PsbV proteins, may serve as a proton exit channel for proton-coupled electron transfer via Y_z [5].

The binding properties of PsbV differ between cyanobacteria and red algae. Reconstitution experiments in *T. vulcanus* revealed that the PsbV protein could bind to PS II complexes independently of the PsbO protein, but this association did not support oxygen-evolving activity. The presence of both the PsbO and PsbU proteins was necessary for

full rebinding and maximal oxygen evolution rates [199]. PsbV was also identified in PS II preparations of the red alga *C. caldarium* [194,226]. In contrast, PsbV could not bind independently of the other extrinsic proteins in red algae [15]. Cross-reconstitution studies found the red algal PsbV protein to be functionally exchangeable between the cyanobacterial and red algal PS II systems. However, this was not true for the cyanobacterial PsbV protein [220]. Thus, while the PsbV proteins from cyanobacteria and red algae are quite similar (40–60% identical), the binding sites of the PS II membrane components are apparently quite different between the two systems [220].

10.2. Function

Subsequent mutant studies in cyanobacteria further solidified the link between the PsbV protein and PS II oxygen evolution. A *Synechocystis* 6803 mutant lacking PsbV ($\Delta psbV$) has reduced accumulation of PS II in the thylakoids (60% that of wild type) and lower oxygen evolution rates (40% relative to wild type) [227]. Consequently, the $\Delta psbV$ mutant exhibits severe growth phenotypes. The doubling time for photoautotrophic growth of $\Delta psbV$ is more than twice that of wild type, and $\Delta psbV$ cells cannot grow when calcium and chloride are eliminated from the growth medium [205,228]. These results suggest that PsbV aids in maintaining the proper ion environment within the oxygen-evolving complex. In this respect, PsbV functions analogously to PsbP in higher plants. Flash oxygen yield experiments have also shown an increase in the miss and double hit factors and slowed $S_3 \rightarrow [S_4] \rightarrow S_0$ transition in mutants lacking the PsbV protein [205]. The photoautotrophic growth and steady state oxygen evolution rates of a *T. elongatus* $\Delta psbV$ mutant were less severe relative to that of the *Synechocystis* 6803 $\Delta psbV$ mutant [229]. Nevertheless, the *T. elongatus* $\Delta psbV$ mutant was unable to grow in medium depleted of either calcium or chloride [229,230].

Additionally, the overall stability of PS II is reduced in the absence of PsbV. Dark treatment as well as heat reduces photosynthetic capacity more dramatically in the *Synechocystis* 6803 $\Delta psbV$ mutant [205]. This correlates with previous biochemical work that identified PsbV as the component responsible for heat stability of PS II oxygen evolution in *Synechococcus* 7002 [210]. However, the *T. elongatus* $\Delta psbV$ mutant did not display the PS II instabilities upon dark incubation or heat treatment as had been reported for the *Synechocystis* 6803 mutant [230]. The authors postulate that some of the functional defects observed in the *Synechocystis* 6803 $\Delta psbV$ mutant may also be attributed to weaker association of the PsbO protein to PS II in these cells. The $\Delta psbO\Delta psbV$ and $\Delta cyanoQ\Delta psbV$ double mutants in *Synechocystis* 6803 do not grow photoautotrophically in standard medium [39,227]. Therefore, the combined loss of PsbV and either PsbO or PsbQ has disastrous consequences for PS II stability and function.

Interestingly, several species of cyanobacteria (*T. elongatus*, *T. vulcanus*, *Acaryochloris marina* MBIC11017, *Arthrospira plantensis* NIES-39, and *Arthrospira maxima*) contain a second copy of the *psbV* gene, denoted *psbV2*. In these cyanobacterial genomes, the *psbV2* gene is adjacent to the *psbV* (or *psbV1*) gene, and, in some cases, these genes cluster with *petJ*, which encodes cytochrome c_6 [229]. While the PsbV2 protein has ~45% sequence identity with that of PsbV1, it lacks the second histidine residue that is normally the sixth axial ligand to the heme group, and it is likely that a tyrosine residue (^{86}Y , *T. elongatus* numbering) provides this ligand in the PsbV2 protein [225]. In *T. elongatus* cells, *psbV2* expression can be detected, albeit at very low levels (~1% that of *psbV1*), indicating that PsbV2 does not significantly contribute to PS II function under normal conditions [225]. Heterologous expression of *T. elongatus psbV2* in *Synechocystis* 6803 $\Delta psbV$ mutant cells restored wild-type photoautotrophic growth rates in complete medium. However, *psbV2* expression could not restore photoautotrophic growth in medium lacking calcium or chloride [229]. Notably, the *T. elongatus psbV1* gene was able to fully complement the *Synechocystis* $\Delta psbV$ mutant phenotypes.

Additionally, *Synechocystis* 6803 $\Delta psbV$ mutant cells heterologously expressing *psbV1* contained detectable amounts of heme corresponding to cytochrome c_{550} ; however, no heme was detected in cells expressing only *psbV2* [231]. While there is no experimental evidence that the heme group of PsbV participates in electron transfer reactions within PS II (see below), it is intriguing that the absence of the heme group in the PsbV2 protein correlates with an inability to fully complement the $\Delta psbV$ mutant phenotype.

10.3. Midpoint potential

The E_m of PsbV and its functional significance has been the subject of intense research. Unlike other c-type monoheme cytochromes, which have an E_m of +0 mV or greater [232], the E_m of soluble PsbV has been reported to be approximately –250 mV in numerous cyanobacterial species [221,230,233,234]. A variety of roles have been proposed for this low-potential cytochrome, including a redox factor in cyclic photosynthetic electron transfer around PS I [235], a ferredoxin electron acceptor for removal of excess electrons under anaerobic conditions [236] and a factor involved in hydrogen metabolism [237]. Because, as discussed above, there is overwhelming evidence that PsbV is associated with PS II function, the negative E_m value is a functional mystery, as participation in electron transfer reactions in the vicinity of the PS II oxygen-evolving complex would require a much more positive E_m value.

Numerous studies have been directed at determining the factors that govern the E_m properties of PsbV. Structural studies and analyses of the heme environment in soluble PsbV suggest that the heme environment of PsbV is quite similar to other c-type cytochromes with much higher E_m values [223–225,238]. The E_m of soluble PsbV is also pH-dependent, possibly due to the protonation state of a tyrosine residue (probably ^{82}Y , *T. elongatus* numbering) [239] or the heme propionate group near ^{49}N (*T. elongatus* numbering) [240]. Notably, mutation of ^{49}N in *Synechocystis* 6803 PsbV resulted in a small potential increase of 12–20 mV, but the pH dependence of the E_m value for this mutant was not reported [241].

More recent studies have focused on determining the E_m of PsbV bound to PS II. Because greater solvent exposure of the heme group tends to lower the E_m of cytochromes, the E_m of PsbV in the context of PS II may be different from that of the soluble protein. Indeed, several groups have reported slightly higher E_m values for PsbV (–110 or –80 mV) in the context of PS II [230,239,240]. Furthermore, the E_m of the PS II-bound form is pH-independent [239]. Kirilovsky and co-workers [230] characterized the E_m of mutants containing site-directed mutations of ^{92}H , the sixth axial ligand to the heme in *T. elongatus*. The H92M and H92C mutants displayed lower E_m values (–140 mV and –164, respectively) relative to wild type in the PS II-bound form (–110 mV). The proteins carrying the H92M and H92C substitutions were more readily lost during PS II purification relative to native PsbV, but H92M and H92C mutants did not display any PS II activity defects at the cellular level. Based on these results, the E_m of PsbV is not critical for PS II function under the conditions analyzed to date.

Recently, Guerrero and co-workers have reported an E_m for PS II-bound PsbV of +200 mV [242]. This vastly different E_m value was determined by reductive titration in the absence of the redox mediators typically used in such titrations. While a much lower E_m value (–200 mV) was observed for oxidative titration in the absence of redox mediators, the authors showed that treatment with dithionite, which is used to fully reduce PsbV prior to oxidative titration measurements, resulted in reduction of the Mn_4CaO_5 cluster and release of PsbV. Therefore, the authors postulate that the presence of redox mediators used in the titrations of previous reports partially reduce the Mn_4CaO_5 cluster of PS II, causing conformational changes which weaken the association of PsbV and lower its midpoint potential. If the E_m of PsbV is +200 mV when bound to PS II, it is possible that electron transfer reactions (ms–s timescale) could occur between the

heme group and the nearest redox-active cofactor, the Mn_4CaO_5 cluster (~23 Å) [242]. To date, there is no experimental evidence for electron transfer between the heme group of PsbV and the Mn_4CaO_5 cluster, but these findings will surely stimulate future studies along these lines.

11. CyanoQ

11.1. Structure

The association of a PsbQ orthologue in cyanobacterial PS II was discovered by Pakrasi and co-workers when analyzing the complete protein complement of isolated PS II complexes from *Synechocystis* 6803 [243]. The cyanobacterial protein, CyanoQ, is related to the authentic PsbQ-1 and PsbQ-2 proteins from *Arabidopsis* (At4g21280 and At4g05180, respectively) rather than the more divergent members of the PsbQ gene family that have auxiliary functions [18]. The X-ray crystal structure of CyanoQ (Fig. 4B) is composed of a four-helix bundle arranged in an up-down-up fold that resembles the PsbQ structure derived from spinach, despite retaining only 17% sequence identity [126–128]. However, the two proteins differ at their N-termini where, in the case of spinach PsbQ, there are two parallel β -strands that are required for binding to the photosystem [127,135]. The absence of this structural motif in CyanoQ suggests the mechanism for binding to cyanobacterial PS II is different from the binding mechanism of PsbQ to plant PS II. In addition, comparison of the electrostatic surfaces of the two proteins reveals CyanoQ possesses a dipole with a positively charged surface comprising helices 2 and 3 and a negatively charged face formed by helices 3 and 4; in contrast, PsbQ has only a positively charged surface formed by helices 2 and 3 [128]. The corresponding helices of the two proteins also differ in length; notably, helix 2 of CyanoQ is split to give a kink by a fully conserved dipeptide composed of ^{77}G and ^{78}P in *Synechocystis* 6803.

A further unique feature of CyanoQ is a motif for a signal-peptide cleavage site with a lipid-modified cysteinyl residue, ^{22}C . [13,244]. The equivalent lipidated cysteinyl is conserved in virtually all cyanobacteria. However, CyanoQ is not present in *Gloeobacter violaceus* or marine *Prochlorococcus* strains [13,20], and some strains of *Synechococcus* either lack a lipidation site or possess a modified lipobox [245]. Furthermore, there is no lipobox associated with the PsbQ-like proteins found in red and brown algae [128].

Both PsbQ and CyanoQ bind zinc; however, although both proteins crystallized with two bound zinc atoms, the binding sites were at different locations in the spinach and cyanobacterial structures. While PsbQ required the presence of zinc for crystallization, the structure of CyanoQ was solved both in the presence and absence of zinc [128]. In the CyanoQ structure, one of the zinc atoms was coordinated by ^{76}H and ^{116}D , which are located above a cavity on the molecular face between helices 2 and 3, and the size of the cavity increased in volume from 208 Å³ to 249 Å³ with zinc bound. Moreover, ^{76}H and ^{116}D and the majority of residues that make up the cavity are highly conserved. Therefore this feature might reflect a specific cation-mediated binding site and mechanism of interaction with another PS II subunit [128,245]. In contrast, the second zinc-binding site in the CyanoQ structure appears to be a crystallization artifact, as the ligands are not conserved.

Direct evidence for the binding of CyanoQ to PS II was obtained by isolating PS II complexes using a histidine tag placed on the C-terminus of the protein [12]. The isolation of PS II complexes has routinely been achieved by placing a histidine tag on one of the core antenna chlorophyll α -binding proteins, CP47 or CP43 [246,247]. The PS II complexes isolated using histidine-tagged CyanoQ were found to have a higher rate of oxygen evolution compared to PS II complexes isolated with the histidine tag on CP47. It has been suggested that this arises because the population of PS II isolated when the tag is on CP47 will

include some inactive PS II centers that are undergoing assembly and repair [12,248].

The PS II crystal structures from *T. elongatus* and *T. vulcanus* do not contain bound CyanoQ [5,7,98]. However, to investigate the binding of CyanoQ to PS II, *in silico* docking studies have been undertaken using the *T. elongatus* 3.0-Å resolution structure (PDB ID: 2AXT), positioning CyanoQ next to PsbV and below CP43 [245].

The result of *in silico* docking studies of CyanoQ to the 1.9-Å resolution structure from *T. vulcanus* (PDB ID: 3ARC) is presented in Fig. 7 and was carried out as described in Fagerlund and Eaton-Rye [245]. The CyanoQ structure from *T. elongatus* was used to model the CyanoQ protein, as no sequence information is available for CyanoQ from *T. vulcanus*. The proximity of CyanoQ to the mouth of the putative B1 oxygen exit channel proposed by Gabdulkhakov et al. [111] is evident. In Fig. 7 CyanoQ sits immediately below the exit of the channel with its putative cation-mediated binding site facing PsbV.

11.2. Function

Synechocystis cells lacking CyanoQ have been shown to exhibit reduced oxygen-evolving activity compared to wild type under CaCl_2 -depleted conditions [13], but photoautotrophic growth of ΔcyanoQ cells under either calcium-limiting or chloride-limiting conditions was only slightly, or not at all, retarded compared to wild type [13,39]. However, upon limitation of calcium, chloride and iron from the growth medium, ΔcyanoQ cells could not grow photoautotrophically [39]. Thus, CyanoQ is not absolutely required for optimal PS II activity unless additional stress conditions exist, a characteristic common to the cyanobacterial PS II extrinsic proteins.

The physiological role of CyanoQ has also been investigated through the phenotypic analysis of double mutants lacking other PS II extrinsic proteins [39]. Under calcium-limiting conditions, the photoautotrophic doubling time of the $\Delta\text{psbU}:\Delta\text{cyanoQ}$ mutant was extended by 30 h compared with ΔpsbU cells, whereas, photoautotrophic growth of the $\Delta\text{psbU}:\Delta\text{cyanoQ}$ strain showed little change under chloride-limiting conditions. However, the photoautotrophic doubling time of the $\Delta\text{psbO}:\Delta\text{cyanoQ}$ mutant in chloride-limiting media was extended by 10 h compared with ΔpsbO cells. In calcium-limiting conditions, strains lacking PsbO are not photoautotrophic. The double mutant lacking PsbV was not photoautotrophic in nutrient-replete conditions; however, the obligate photoheterotrophic phenotype of the $\Delta\text{psbV}:\Delta\text{cyanoQ}$ strain was restored to photoautotrophy by growing the cells at pH 10 [39]. As noted elsewhere a similar pH dependence has been observed with the $\Delta\text{psbO}:\Delta\text{psbU}$ double mutant [201,202,249] although the molecular mechanism underlying this pH effect is currently not known.

The absence of CyanoQ has a more pronounced effect on PS II activity and stability at the level of isolated PS II complexes. PS II complexes from ΔcyanoQ had greater oxygen-evolving activity in the presence of 20 mM CaCl_2 compared to 5 mM CaCl_2 , whereas the rates observed with PS II complexes isolated from wild type were similar at both CaCl_2 concentrations [13,40]. Furthermore, PS II complexes isolated from cells lacking CyanoQ were found to contain lower levels of PsbV than PS II complexes isolated from wild type [12,40]. Removal of PsbV is known to destabilize PS II and introduce a dependence on calcium and chloride for photoautotrophy (see section on the role of the PsbV protein). Hence, while the absence of CyanoQ is associated with an enhanced requirement for calcium and chloride (particularly in the double mutants lacking either PsbO or PsbU and in isolated PS II complexes) the reduced level of PsbV in the absence of CyanoQ is sufficient to maintain PS II activity that is similar to that seen in wild type in BG-11 growth medium. Additionally, PS II complexes lacking CyanoQ are more sensitive to the reducing agent hydroxylamine (NH_2OH) than wild type [40]. This implies that while the water oxidation machinery is largely protected from hydroxylamine in wild type, it appears to be partially




Fig. 7. Docking of *Thermosynechococcus elongatus* CyanoQ, modeled from the *Synechocystis* sp. PCC 6803 CyanoQ structure [128], to *Thermosynechococcus vulcanus* PS II. Orientation is looking down from the luminal side of the membrane. The putative oxygen exit channel (gray) was calculated using the program CAVER [255,260] and begins at Mn1 of the Mn_4CaO_5 cluster. Waters (red spheres) and glycerol from the PDB ID: 3ARC file are shown in the channel. The PS II subunits are colored: D1 (red), D2 (orange), CP43 (cyan), PsbE, (magenta), PsbF (blue) PsbJ (deep teal), and, PsbV (pale green). Calcium (yellow), chloride (green) and magnesium (purple) ions are shown as spheres. Helices of CyanoQ are rainbow-colored from the N-terminus (red).

exposed in the $\Delta cyanoQ$ mutant; consequently, the PS II complexes in this strain are more easily inactivated.

12. CyanoP

12.1. Structure

A PsbP orthologue was first identified in cyanobacteria in isolated PS II complexes from *Synechocystis* 6803 [243]. Subsequently, it has been shown through phylogenetic studies that PsbP-like proteins are found in all oxygenic photosynthetic organisms [18,250,251]. These PsbP-like proteins represent an extended family with eleven homologues identified in *Arabidopsis* belonging to three distinct groups: PS II-specific or authentic PsbP (two homologues), PsbP-like (two homologues), and those containing a “PsbP-like domain” (seven homologues) [123,251]. The homologue in *Synechocystis* 6803, referred to as CyanoP [123,248], is most closely related to the PsbP-like PPL1 protein of *Arabidopsis* that has been suggested to play a role in the repair of photodamaged PS II [252]. The relationship between the PsbP-like proteins and CyanoP has also been supported by phylogenetic studies [251].

CyanoP, like CyanoQ, possesses a lipobox for signal-peptide cleavage and lipidation at a Cys [13]. Lipidation of CyanoP has not yet been demonstrated, but CyanoP was not released from the thylakoid membrane by salt washes that release other PS II extrinsic proteins, consistent with lipid-mediated membrane attachment. Also consistent with lipidation, the N-terminus of the protein was blocked when sequenced by Edman degradation [14]. To date CyanoP has not been found in any X-ray-derived crystal structures of cyanobacterial PS II; however, a functional relationship with PS II has been inferred from the observation that CyanoP was not present in mutant strains of *Synechocystis* 6803 that lack PS II: hence it seems expression of CyanoP requires the presence of PS II [14].

The crystal structure of CyanoP from *T. elongatus* has been solved to 2.8 Å and is an antiparallel β -sheet sandwiched between α -helices (Fig. 3B) (PDB ID: 2XB3, [123]). This structure is similar to the structures solved for the tobacco and spinach PsbP proteins, despite low (~20%) sequence identity [120,121,123]. CyanoP was crystallized in the presence of zinc and it was shown the protein coordinated five zinc ions. Seven distinct contacts were made with the metal ion as a result of the protein's crystallization symmetry. It is possible that

some of these sites may be artifacts of the crystallization process; however, two are highly conserved and the binding of zinc at one of these two sites (zinc ion 4 in [123]) is at an equivalent position to the zinc ion located in the PsbP structure from spinach. Moreover, conserved residues from both of the conserved zinc-binding sites have been previously identified as potential ligands that may interact with manganese in tobacco PsbP [253] (however, see [118] for a critical analysis of this hypothesis). The physiological significance of a divalent metal-binding site on CyanoP is unknown, although conservation of coordinating residues across phyla suggests it may have an as of yet unidentified role.

The docking of CyanoP to the PS II structure from *T. elongatus* (PDB ID: 2AXT) has been modeled and positions CyanoP below D2 and away from the other extrinsic proteins. In addition, a putative interaction between the C-terminus of the α -subunit of cytochrome b_{559} (PsbE) and a cavity on the CyanoP surface was identified [245]. *In silico* docking to the *T. vulcanus* PS II structure (PDB ID: 3ARC) yielded similar results (Fig. 8). However, an additional interaction with a helix incorporating ^{277}S to ^{294}S of loop E of CP47 was predicted. A *Synechocystis* 6803 mutant, the CP47: $\Delta(K277-E283)$ strain, containing a 7 amino acid deletion in this region of loop E, has been reported to have impaired photoautotrophic growth and impaired oxygen evolution and reduced levels of assembled PS II centers [254]. Intriguingly, the CAVER program [255] predicted the presence of a proton exit channel leading directly from Y_D (^{160}O of D2) to the bulk phase at the docking site of CyanoP (Fig. 7). The required hydrogen bond network for this protein pathway may be provided by waters and the side chain of $^{180}R:D2$. This channel is composed of residues from CP47, D1, PsbV and PsbE and the mouth comprises residues from D1, PsbV, PsbE and D2.

13.2. Function

Synechocystis 6803 strains lacking CyanoP typically exhibited photoautotrophic growth rates in BG-11 that are similar to wild type; even under chloride- or calcium-limiting conditions only small effects on photoautotrophic growth have been observed [13,14,256]. However, $\Delta cyanoP$ cells grown in $CaCl_2$ -depleted medium were observed to exhibit a rate of oxygen evolution 63% of that found for the wild type grown under similar conditions [13]. Furthermore, oxygen evolution measurements of PS II complexes purified via a histidine tag on CP47




Fig. 8. Docking of *Thermosynechococcus elongatus* CyanoP to *Thermosynechococcus vulcanus* PS II (PDB ID: 3ARC). Orientation is looking down from the luminal side of the membrane. A putative proton exit channel (gray) predicted by the CAVER program [255,260] originating from Y_D ($^{160}Y:D2$) is shown. Waters (red spheres) from the PDB ID: 3ARC file are shown in the channel. The PS II subunits are colored: D1 (purple), D2 (orange), CP47 (brown), PsbE (magenta), and, PsbV (pale green). The CyanoP ribbon diagram is rainbow-colored from the N-terminus (red).

revealed that complexes isolated from Δ cyanoP cells had reduced oxygen-evolving activity compared to isolated PS II from wild type, particularly in the presence of reduced levels of calcium, suggesting CyanoP was necessary for optimal PS II activity in these preparations [13].

Since it has been suggested that PsbU and PsbV might have similar functions to PsbP and PsbQ in plants and green algae [199,257], the possibility that CyanoP is required when either PsbU or PsbV is absent was investigated by the construction of two double mutants, Δ psbU: Δ cyanoP and Δ psbV: Δ cyanoP [256]. The genetic deletion of CyanoP in combination with these other extrinsic components had little or no additional effects on photoautotrophic growth in normal as well as chloride- or calcium-limiting medium. A similar result was obtained with both the Δ psbO: Δ cyanoP and Δ cyanoQ: Δ cyanoP strains. Likewise, no specific effects on oxygen evolution attributable to the removal of CyanoP were observed in any of the double mutants. Moreover, there was no additive effect of removing CyanoP in the Δ psbO: Δ psbU: Δ cyanoP triple mutant compared to the corresponding double mutant strains [256].

Since results obtained with mutants lacking CyanoP produced small effects and these results varied among different laboratories, a comparative study of the three available Δ cyanoP strains was carried out [258]. Competitive growth of the wild type with the mutants indicated a selective advantage for the presence of CyanoP. Furthermore, flash-induced oxygen yield experiments displayed a decreased level of oxygen produced on the third flash, increased level on the fourth flash, and an earlier dampening of the oscillations relative to the wild type in Δ cyanoP cells. Additionally, thermoluminescence measurements demonstrated an apparent enhanced recombination between the S_3 state of the oxygen-evolving complex and the reduced secondary plastoquinone electron acceptor Q_B in the Δ cyanoP strains. These data indicate that the Δ cyanoP mutants exhibit a major $S_3Q_B^-$ recombination in addition to $S_2Q_B^-$ recombination, whereas the wild type predominately exhibits the thermoluminescence band corresponding to $S_2Q_B^-$ recombination. These results may implicate equilibration between Y_D (^{160}Y of D2) (or some other unspecified redox-active species) and the S_3 state favoring, for example, $Y_D^{red}S_3$ over $Y_D^{ox}S_2$ in PS II centers lacking CyanoP, thereby pointing to a role for CyanoP in the stabilization of charge separation in PS II [258]. It is therefore of interest that a putative proton exit channel linking Y_D to a possible docking site for CyanoP (Fig. 8) has been detected; however, the existence of this channel requires experimental verification.

Additionally, immunodetection of CyanoP in *Synechocystis* 6803 thylakoid membranes has indicated that CyanoP is present at a similar level to the PsbO protein [14], although this finding is controversial

[13]. Thus, oxygen flash yield experiments, thermoluminescence and western blotting suggest CyanoP is stoichiometrically present in PS II [14] in contrast to an earlier finding that it is associated only with ~3% of PS II complexes in the thylakoid membrane [13]. As detergent is required during the isolation of PS II complexes, it is possible that CyanoP is lost during complex purification [13,14,258]. However, despite the uncertainty regarding the stoichiometry of CyanoP, the role of this subunit is not essential under a wide range of conditions and it appears likely that the selective advantage conferred by the protein is linked to a regulatory role that remains to be fully elucidated.

13. Conclusions

As is evident in this review, over the last thirty years an enormous amount of effort has been focused on the elucidation of the structure and function of the extrinsic proteins of PS II. Much progress has clearly been made, but many questions remain unanswered. While phenomenologically the functions of these proteins have generally been identified, the mechanisms by which they influence PS II oxygen evolution remains elusive. While the structure of cyanobacterial PS II progresses towards atomic resolution (i.e. 1.2 Å), the organization of the CyanoP and CyanoQ subunits within the photosystem remains unclear. Experimental investigations into the structural and possible regulatory roles played by the extrinsic components in the numerous channels present in PS II must also be forthcoming. Additionally, while structures for isolated higher plant PsbP and PsbQ are available, no progress is evident in the elucidation of the overall structure of the higher plant photosystem. Clearly, much work remains.

Acknowledgements

Funding was provided by a United States Department of Energy grant # DE-FG02-98ER20310 to T.M.B. and L.K.F., from the USDA National Institute of Food and Agriculture Grant # 2008-35318-04605 to J.L.R., and by a Marsden Grant # 08-U00-043 to J.J.E.-R.

References

- [1] R.J. Debus, The manganese and calcium ions of photosynthetic oxygen evolution, *Biochim. Biophys. Acta* 1102 (1992) 269–352.
- [2] R.D. Britt, Oxygen evolution, in: D. Ort, C.F. Yocum (Eds.), *Oxygenic Photosynthesis: The Light Reactions*, Advances in Photosynthesis, vol. 4, Kluwer Academic Publishers, Dordrecht, 1996, pp. 137–164.

- [3] D.R. Ort, C.F. Yocum, Energy transfer and energy transduction in photosynthesis: an overview, in: D.R. Ort, C.F. Yocum (Eds.), *Oxygenic Photosynthesis: The Light Reactions*, Advances in Photosynthesis, vol. 4, Kluwer Academic Publishers, Dordrecht, 1996, pp. 1–9.
- [4] G. Renger, Light induced oxidative water splitting in photosynthesis: energetics, kinetics and mechanism, *Photochem. Photobiol. B: Biol.* 104 (2011) 35–43.
- [5] Y. Umena, K. Kawakami, J.-R. Shen, N. Kamiya, Crystal structure of oxygen-evolving Photosystem II at a resolution of 1.9 Å, *Nature* 473 (2011) 55–60.
- [6] C. Rosenberg, J. Christian, T.M. Bricker, C. Putnam-Evans, Site-directed mutagenesis of glutamate residues in the large extrinsic loop of the Photosystem II protein CP 43 affects oxygen-evolving activity and PS II assembly, *Biochemistry* 38 (1999) 15994–16000.
- [7] K.N. Ferreira, T.M. Iverson, K. Maghlaoui, J. Barber, S. Iwata, Architecture of the photosynthetic oxygen-evolving center, *Science* 303 (2004) 1831–1838.
- [8] A. Guskov, J. Kern, A. Gabdulkhakov, M. Broser, A. Zouni, W. Saenger, Cyanobacterial Photosystem II at 2.9-Å resolution and the roles of quinones, lipids, channels and chloride, *Nat. Struct. Mol. Biol.* 16 (2009) 334–342.
- [9] M. Miyao, N. Murata, Role of the 33 kDa polypeptide in preserving Mn in photosynthetic oxygen-evolution, *FEBS Lett.* 170 (1984) 350–354.
- [10] R.L. Burnap, L.A. Sherman, Deletion mutagenesis in *Synechocystis* sp. PCC 6803 indicates that the Mn-stabilizing protein of Photosystem II is not essential for oxygen evolution, *Biochemistry* 30 (1991) 440–446.
- [11] T.M. Bricker, Oxygen evolution in the absence of the 33 kDa manganese-stabilizing protein, *Biochemistry* 31 (1992) 4623–4628.
- [12] J.L. Roose, Y. Kashino, H.B. Pakrasi, The PsbQ protein defines cyanobacterial Photosystem II complexes with highest activity and stability, *Proc. Natl. Acad. Sci. U.S.A.* 104 (2007) 2548–2553.
- [13] L. Thornton, H. Ohkawa, J. Roose, Y. Kashino, N. Keren, H. Pakrasi, Homologs of plant PsbP and PsbQ proteins are necessary for regulation of Photosystem II activity in the cyanobacterium *Synechocystis* 6803, *Plant Cell* 16 (2004) 2164–2175.
- [14] Y. Ishikawa, W.P. Schroder, C. Funk, Functional analysis of the PsbP-like protein (*slh1418*) in *Synechocystis* sp. PCC 6803, *Photosynth. Res.* 84 (2005) 257–262.
- [15] I. Enami, S. Kikuchi, T. Fukuda, H. Ohta, J.-R. Shen, Binding and functional properties of four extrinsic proteins of Photosystem II from a red alga, *Cyanidium caldarium*, as studied by release-reconstitution experiments, *Biochemistry* 37 (1998) 2787–2793.
- [16] H. Ohta, T. Suzuki, M. Ueno, A. Okumura, S. Yoshihara, J.-R. Shen, I. Enami, Extrinsic proteins of Photosystem II: an intermediate member of the PsbQ protein family in red algal PS II, *Eur. J. Biochem.* 270 (2003) 4156–4163.
- [17] K. Ifuku, S. Ishihara, R. Shimamoto, K. Ido, F. Sato, Structure, function, and evolution of the PsbP protein family in higher plants, *Photosynth. Res.* 98 (2008) 427–437.
- [18] K. Ifuku, S. Ishihara, F. Satoh, Molecular functions of oxygen-evolving complex family proteins in photosynthetic electron flow, *J. Integr. Plant Biol.* 52 (2010) 723–734.
- [19] K. Ifuku, K. Ido, F. Sato, Molecular functions of PsbP and PsbQ proteins in the Photosystem II supercomplex, *J. Photochem. Photobiol. B: Biol.* 104 (2011) 158–164.
- [20] J. De Las Rivas, J. Barber, Analysis of the structure of the PsbO protein and its implications, *Photosynth. Res.* 81 (2004) 329–343.
- [21] T.M. Bricker, R.L. Burnap, The extrinsic proteins of Photosystem II, in: T. Wydrzynski, K. Satoh (Eds.), *Photosystem II: The Water/Plastoquinone Oxidoreductase of Photosynthesis*, Advances in Photosynthesis and Respiration, vol. 22, Springer, Dordrecht, 2005, pp. 95–120.
- [22] Q. Xu, J. Nelson, T.M. Bricker, Secondary structure of the 33 kDa, extrinsic protein of Photosystem II: a far-UV circular dichroism study, *Biochim. Biophys. Acta* 1188 (1994) 427–431.
- [23] J. De Las Rivas, J. Barber, Structure and thermal stability of Photosystem II reaction centers studied by infrared spectroscopy, *Biochemistry* 36 (1997) 8897–8903.
- [24] I.Z. Zubrzycki, L.K. Frankel, P.S. Russo, T.M. Bricker, Hydrodynamic studies on the manganese-stabilizing protein of Photosystem II, *Biochemistry* 37 (1998) 13553–13558.
- [25] B. Svensson, D.M. Tiede, B.A. Barry, Small-angle X-ray scattering studies of the manganese-stabilizing protein in Photosystem II, *J. Phys. Chem. B* 106 (2002) 8485–8488.
- [26] S.J. Darnell, D. Page, J.C. Mitchell, Automated decision-tree approach to predicting protein–protein interaction hot spots, *Proteins* 68 (2007) 813–823.
- [27] J.P. Gallivan, D.A. Dougherty, Cation– π interactions in structural biology, *Proc. Natl. Acad. Sci. U.S.A.* 96 (1999) 9459–9464.
- [28] C. Putnam-Evans, J. Wu, R. Burnap, J. Whitmarsh, T.M. Bricker, Site-directed mutagenesis of the CP 47 Protein of Photosystem II: Alteration of conserved charged residues in the domain ³⁶⁴E–⁴⁴⁴R, *Biochemistry* 35 (1996) 4046–4053.
- [29] C. Putnam-Evans, T.M. Bricker, Site-directed mutagenesis of the CPa-1 protein of Photosystem II: Alteration of the basic residue pair ^{384,385}R to ^{384,385}G leads to a defect associated with the oxygen-evolving complex, *Biochemistry* 31 (1992) 11482–11488.
- [30] M. Qian, S. Al-Khaldi, C. Putnam-Evans, T.M. Bricker, R.L. Burnap, Photoassembly of the Photosystem II (Mn)₄ cluster in site-directed mutants impaired in the binding of the manganese stabilizing protein, *Biochemistry* 36 (1997) 15244–15252.
- [31] F.K. Bentley, J. Eaton-Rye, The effect of removing Photosystem II extrinsic proteins on dimer formation and recovery from photodamage in *Synechocystis* sp. PCC 6803, in: J.F. Allen, E. Gantt, J.H. Golbeck, B. Osmond (Eds.), *Photosynthesis. Energy from the Sun: 14th International Congress on Photosynthesis*, Springer, Dordrecht, 2008, pp. 715–717.
- [32] M.G. Kuhn, W.F.J. Vermaas, Deletion mutations in a long hydrophilic loop in the Photosystem II chlorophyll-binding protein CP43 in the cyanobacterium *Synechocystis* sp. PCC 6803, *Plant Mol. Biol.* 23 (1993) 123–133.
- [33] J.J. Eaton-Rye, N. Murata, Evidence that the amino-terminus of the 33 kDa extrinsic protein is required for binding to the Photosystem II complex, *Biochim. Biophys. Acta* 977 (1989) 219–226.
- [34] H. Popelkova, M.M. Im, C.F. Yocum, N-terminal truncations of the manganese-stabilizing protein identify two amino acid sequences required for binding of the eukaryotic protein to Photosystem II and reveal the absence of one binding-related sequence in cyanobacteria, *Biochemistry* 41 (2002) 10038–10045.
- [35] A. Seidler, A.W. Rutherford, H. Michel, On the role of the N-terminus of the extrinsic 33 kDa protein in Photosystem II, *Plant Mol. Biol.* 31 (1996) 183–188.
- [36] A. Motoki, M. Usui, T. Shimazu, M. Hirano, S. Katoh, A domain of the Mn-stabilizing protein from *Synechococcus elongatus* involved in functional binding to Photosystem II, *J. Biol. Chem.* 277 (2002) 14747–14756.
- [37] Z. Li, R.L. Burnap, Mutations of basic arginine residue 334 in the D1 protein of Photosystem II lead to unusual S₂ state properties in *Synechocystis* sp. PCC 6803, *Photosynth. Res.* 72 (2002) 191–201.
- [38] S.F. Al-Khaldi, J. Coker, J.R. Shen, R.L. Burnap, Characterization of site-directed mutants in manganese-stabilizing protein (MSP) of *Synechocystis* sp. PCC6803 unable to grow photoautotrophically in the absence of cytochrome c-550, *Plant Mol. Biol.* 43 (2000) 33–41.
- [39] T.C. Summerfield, J.A. Shand, F.K. Bentley, J.J. Eaton-Rye, PsbQ (Slh1638) in *Synechocystis* sp. PCC 6803 is required for Photosystem II activity in specific mutants and in nutrient-limiting conditions, *Biochemistry* 44 (2005) 805–815.
- [40] Y. Kashino, N. Inoue-Kashino, J. Roose, H. Pakrasi, Absence of the PsbQ protein results in destabilization of the PsbV protein and decreased oxygen evolution activity in cyanobacterial Photosystem II, *J. Biol. Chem.* 281 (2006) 20834–20841.
- [41] T.M. Bricker, W.R. Odom, C.B. Queirolo, Close association of the 33 kDa extrinsic protein with the apoprotein of CPa-1 in Photosystem II, *FEBS Lett.* 231 (1988) 111–117.
- [42] I. Enami, T. Miyaoka, Y. Mochizuki, J.-R. Shen, K. Satoh, S. Katoh, Nearest neighbor relationships among constituent proteins of oxygen-evolving Photosystem II membranes: binding and function of the extrinsic 33 kDa protein, *Biochim. Biophys. Acta* 973 (1989) 35–40.
- [43] R. Nagao, T. Suzuki, A. Okumura, A. Niikura, M. Iwai, N. Dohmae, T. Tomo, J.-R. Shen, M. Ikeuchi, I. Enami, Topological analysis of the extrinsic PsbO, PsbP and PsbQ proteins in a green algal PSII complex by cross-linking with a water-soluble carbodiimide, *Plant Cell Physiol.* 51 (2010) 718–727.
- [44] K.-C. Han, J.-R. Shen, M. Ikeuchi, Y. Inoue, Chemical crosslinking studies of extrinsic proteins in cyanobacterial Photosystem II, *FEBS Lett.* 355 (1994) 121–124.
- [45] C.S. Hackett, P. Strittmatter, Covalent crosslinking of the active sites of vesicle-bound cytochrome b₅ and NADH cytochrome b₅ reductase, *J. Biol. Chem.* 259 (1984) 3275–3282.
- [46] W. Odom, T.M. Bricker, Interaction of CPa-1 with the manganese-stabilizing protein of Photosystem II: identification of domains cross-linked by 1-ethyl-3-[3-(dimethylamino)propyl]carbodiimide, *Biochemistry* 31 (1992) 5616–5620.
- [47] L.K. Frankel, T.M. Bricker, Interaction of CPa-1 with the manganese-stabilizing protein of Photosystem II: identification of domains on CPa-1 which are shielded from N-hydroxysuccinimide biotinylation by the manganese-stabilizing protein, *Biochemistry* 31 (1992) 11059–11063.
- [48] I. Enami, S. Ohta, S. Mitsuhashi, S. Takahashi, M. Ikeuchi, S. Katoh, Evidence from crosslinking for a close association of the extrinsic 33 kDa protein with the 9.4 kDa subunit of cytochrome b₅₅₉ and the 4.8 kDa product of the *psbI* gene in oxygen-evolving Photosystem II complexes from spinach, *Plant Cell Physiol.* 33 (1992) 291–297.
- [49] G.E. Means, R.E. Feeney, *Chemical Modification of Proteins*, Holden-Day, Inc., San Francisco, 1971.
- [50] J. Niels, J. Barber, Refinement of the structural model for the Photosystem II supercomplex of higher plants, *Biochim. Biophys. Acta* 1757 (2006) 353–361.
- [51] D. Slowik, M. Rossmann, P.V. Konarev, K.-D. Irrgang, W. Saenger, Structural investigation of PsbO from plant and cyanobacterial Photosystem II, *J. Mol. Biol.* 407 (2011) 125–137.
- [52] T.M. Bricker, L.K. Frankel, The structure and function of the 33 kDa extrinsic protein of Photosystem II. A critical review, *Photosynth. Res.* 56 (1998) 157–173.
- [53] H. Popelkova, C.F. Yocum, PsbO, the manganese-stabilizing protein: analysis of the structure–function relations that provide insights into its role in Photosystem II, *Photochem. Photobiol. B: Biol.* 104 (2011) 179–190.
- [54] Q. Xu, T.M. Bricker, Structural organization of proteins on the oxidizing side of Photosystem II. Two molecules of the 33-kDa manganese-stabilizing protein per reaction center, *J. Biol. Chem.* 267 (1992) 25816–25821.
- [55] C. Leuschner, T.M. Bricker, Interaction of the 33 kDa extrinsic protein with Photosystem II: rebinding of the 33 kDa extrinsic protein to Photosystem II membranes which contain four, two, or zero manganese per Photosystem II reaction center, *Biochemistry* 35 (1996) 4551–4557.
- [56] Y. Yamamoto, S. Nakayama, C.L. Cohn, D.W. Krogmann, Highly efficient purification of the 33-, 24-, and 18-kDa proteins in spinach Photosystem II by butanol/water phase partitioning and high-performance liquid chromatography, *Arch. Biochem. Biophys.* 255 (1987) 156–161.
- [57] A. Seidler, Introduction of a histidine tail at the N-terminus of a secretory protein expressed in *E. coli*, *Protein Eng.* 7 (1994) 1277–1280.

- [58] S.D. Betts, J.R. Ross, E. Pichersky, C.F. Yocum, Mutation Val235Ala weakens binding of the 33-kDa manganese-stabilizing protein of Photosystem II to one of two sites, *Biochemistry* 36 (1997) 4047–4053.
- [59] R. Murakami, K. Ifuku, A. Takabayashi, T. Shikanai, T. Endo, F. Sato, Functional dissection of two *Arabidopsis* PsbO proteins PsbO1 and PsbO2, *FEBS J.* 272 (2005) 2165–2175.
- [60] H. Popelkova, M.M. Im, J. D'Auria, S.D. Betts, N. Lydakis-Simantiris, C.F. Yocum, N-terminus of the Photosystem II manganese-stabilizing protein: effects of sequence elongation and truncation, *Biochemistry* 41 (2002) 2702–2711.
- [61] H. Popelkova, A. Commet, C.F. Yocum, Asp157 is required for the function of PsbO, the Photosystem II manganese-stabilizing protein, *Biochemistry* 48 (2009) 11920–11928.
- [62] L.K. Frankel, J.A. Cruz, T.M. Bricker, The role of carboxylic acid residues on the manganese-stabilizing protein in its binding to Photosystem II, *Biochemistry* 38 (1999) 14271–14278.
- [63] H. Popelkova, M.M. Im, C.F. Yocum, Binding of manganese-stabilizing protein to Photosystem II: identification of essential N-terminal threonine residues and domains that prevent nonspecific binding, *Biochemistry* 42 (2003) 6193–6200.
- [64] H. Popelkova, A. Commet, T. Kuntzleman, C.F. Yocum, Inorganic cofactor stabilization and retention: the unique functions of the two PsbO subunits of eukaryotic Photosystem II, *Biochemistry* 47 (2008) 12600–12953.
- [65] J.L. Roose, C.F. Yocum, H. Popelkova, Binding stoichiometry and affinity of the manganese-stabilizing protein affects redox reactions on the oxidizing side of Photosystem II, *Biochemistry* 50 (2011) 5988–5998.
- [66] R. Anati, N. Adir, Crystallization of dimers of the manganese-stabilizing protein of Photosystem II, *Photosynth. Res.* 64 (2000) 167–177.
- [67] E.J. Boekema, J.F.L. van Breeman, H. van Roon, J.P. Dekker, Conformational changes in Photosystem II supercomplexes upon removal of extrinsic subunits, *Biochemistry* 39 (2000) 12907–12915.
- [68] L.K. Frankel, T.M. Bricker, Interaction of the 33 kDa extrinsic protein with Photosystem II: identification of domains on the 33 kDa protein that are shielded from NHS biotinylation by Photosystem II, *Biochemistry* 34 (1995) 7492–7497.
- [69] H. Popelkova, A. Commet, C.F. Yocum, The double mutant DL6MW241F in PsbO, the Photosystem II manganese-stabilizing protein, yields insights into its evolution of its structure and function, *FEBS Lett.* 584 (2010) 4009–4014.
- [70] T.-A. Ono, Y. Inoue, Mn-preserving extraction of 33-, 24- and 16-kDa proteins from O₂-evolving PS II particles by divalent salt-washing, *FEBS Lett.* 164 (1983) 255–260.
- [71] T. Ono, Y. Inoue, S-State turnover in the O₂-evolving system of CaCl₂-washed Photosystem II particles depleted of three peripheral proteins as measured by thermoluminescence. Removal of 33 kDa protein inhibits S₃ to S₄ transition, *Biochim. Biophys. Acta* 806 (1985) 331–340.
- [72] M. Miyao, N. Murata, J. Lavorel, B. Maisson-peteri, A. Boussac, A.-L. Etienne, Effect of the 33-kDa protein on the S-state transitions in photosynthetic oxygen evolution, *Biochim. Biophys. Acta* 890 (1987) 151–159.
- [73] R. Murakami, K. Ifuku, A. Takabayashi, T. Shikanai, T. Endo, F. Sato, Characterization of an *Arabidopsis thaliana* mutant with impaired psbO, one of two genes encoding extrinsic 33-kDa proteins in Photosystem II, *FEBS Lett.* 523 (2002) 138–142.
- [74] H. Liu, L.K. Frankel, T.M. Bricker, Functional analysis of Photosystem II in a PsbO-1 deficient mutant in *Arabidopsis thaliana*, *Biochemistry* 46 (2007) 7607–7613.
- [75] Y. Allahverdiyeva, F. Mamedov, M. Holmström, M. Nurmi, B. Lundin, S. Styring, C. Spetea, E.-M. Aro, Comparison of the electron transport properties of the psbO1 and psbO2 mutants of *Arabidopsis thaliana*, *Biochim. Biophys. Acta* 1787 (2009) 1230–1237.
- [76] R.L. Burnap, J.-R. Shen, P.A. Jursinic, Y. Inoue, L.A. Sherman, Oxygen yield and thermoluminescence characteristics of a cyanobacterium lacking the manganese-stabilizing protein of Photosystem II, *Biochemistry* 31 (1992) 7404–7410.
- [77] I. Vass, K.M. Cool, D. Zsuzsanra, S.R. Mayes, J. Barber, Thermoluminescence and flash oxygen characterization of the IC2 deletion mutant of *Synechocystis* sp. PCC 6803 lacking the Photosystem II 33 kDa protein, *Biochim. Biophys. Acta* 1102 (1992) 195–201.
- [78] M.R. Razeghifard, T. Wydrzynski, R.J. Pace, R.L. Burnap, Y₂* reduction kinetics in the absence of the manganese-stabilizing protein of Photosystem II, *Biochemistry* 36 (1997) 14474–14478.
- [79] K. Lindberg, L.-E. Andreasson, On the binding and function of chloride in Photosystem II, in: P. Mathis (Ed.), *Photosynthesis: From Light to Biosphere: Proceedings of the Xth International Photosynthesis Congress*, vol. 2, Kluwer Academic Publishers, Dordrecht, 1995, pp. 401–404.
- [80] T. Ono, J.L. Zimmerman, Y. Inoue, A.W. Rutherford, EPR evidence for a modified S-state transition in chloride-depleted Photosystem II, *Biochim. Biophys. Acta* 851 (1986) 193–201.
- [81] S. Styring, M. Miyao, A.W. Rutherford, Formation and flash-dependent oscillation of the S₂-state multiline EPR signal in an oxygen-evolving Photosystem II preparation lacking the three extrinsic proteins in the oxygen-evolving system, *Biochim. Biophys. Acta* 767 (1987) 32–38.
- [82] H. Wincencjusz, H.J. van Gorkom, C.F. Yocum, The photosynthetic oxygen evolving complex requires chloride for its redox state S₂→S₃ and S₃→S₀ transitions but not for S₀→S₁ or S₁→S₂ transitions, *Biochemistry* 36 (1997) 3663–3670.
- [83] K. Lubbers, W. Drevenstedt, W. Junge, Chloride-depletion of photosynthetic water oxidase: no proton release during the second oxidation step, S²⁺→S³⁺, and a transmembrane radical pair recombination from the third on, *Biochim. Biophys. Acta* 336 (1993) 304–308.
- [84] T. Ono, T. Noguchi, Y. Inoue, M. Kusunoki, H. Yamaguchi, H. Oyanagi, XANES spectroscopy for monitoring intermediate reaction states of Cl⁻-depleted Mn cluster in photosynthetic water oxidation enzyme, *J. Am. Chem. Soc.* 117 (1995) 6386–6387.
- [85] A. Boussac, J.L. Zimmermann, A.W. Rutherford, EPR signals from modified charge accumulation states of the oxygen evolving enzyme in calcium deficient Photosystem II, *Biochemistry* 28 (1989) 8984–8989.
- [86] J.C. Depaula, P.M. Li, A.F. Miller, B.W. Wu, G.W. Brudwig, Effect of the 17-kilodalton and 23-kilodalton polypeptides, calcium, and chloride on electron-transfer in Photosystem II, *Biochemistry* 25 (1986) 6487–6494.
- [87] M. Haumann, W. Junge, Evidence for impaired hydrogen-bonding of tyrosine Y₂ in calcium-depleted Photosystem II, *Biochim. Biophys. Acta* 1411 (1999) 121–133.
- [88] T.M. Bricker, L.K. Frankel, The *Arabidopsis psbO1* mutant cannot efficiently utilize calcium in support of photosynthetic oxygen evolution, *J. Biol. Chem.* 283 (2008) 29022–29027.
- [89] G. Mavankal, D.C. McCain, T.M. Bricker, Effects of chloride on paramagnetic coupling of manganese in calcium chloride-washed Photosystem II preparations, *FEBS Lett.* 202 (1986) 235–239.
- [90] M. Miyao, N. Murata, Calcium ions can be substituted for the 24 kDa polypeptide in photosynthetic oxygen evolution, *FEBS Lett.* 168 (1984) 118–120.
- [91] T.A. Ono, Y. Inoue, Reconstitution of photosynthetic oxygen evolving activity by rebinding of 33 kDa protein to CaCl₂-extracted PS II particles, *FEBS Lett.* 166 (1984) 381–384.
- [92] J. Cole, M. Boska, N.V. Blough, K. Sauer, Reversible and irreversible effects of alkaline pH on Photosystem II electron-transfer reactions, *Biochim. Biophys. Acta* 848 (1986) 41–47.
- [93] W.J. Coleman, Govidjee, A model for the mechanism of chloride activation of oxygen evolution in Photosystem II, *Photosynth. Res.* 13 (1987) 199–223.
- [94] R. Wales, B.J. Newman, D. Pappin, J.C. Gray, The extrinsic 33 kDa protein of the oxygen-evolving complex of Photosystem II is a putative calcium-binding protein and is encoded by a multi-gene family in pea, *Plant Mol. Biol.* 12 (1989) 439–451.
- [95] A.N. Webber, J.C. Gray, Detection of calcium binding by Photosystem II polypeptides immobilized onto nitrocellulose membranes, *FEBS Lett.* 249 (1989) 79–82.
- [96] A. Seidler, The extrinsic proteins of Photosystem II, *Biochim. Biophys. Acta* 1277 (1996) 35–60.
- [97] A. Seidler, A.W. Rutherford, The role of the extrinsic 33 kDa protein in Ca⁺² binding in Photosystem II, *Biochemistry* 35 (1996) 12104–12110.
- [98] B. Loll, N. Kern, W. Saenger, A. Zouni, J. Biesiadka, Towards complete cofactor arrangement in the 3.0 Å resolution structure of Photosystem II, *Nature* 438 (2006) 1040–1044.
- [99] J.W. Murray, J. Barber, Identification of a calcium-binding site in the PsbO protein of Photosystem II, *Biochemistry* 45 (2006) 4128–4130.
- [100] J.B. Philbrick, B.A. Diner, B.A. Zilinskas, Construction and characterization of cyanobacterial mutants lacking the manganese-stabilizing protein of Photosystem II, *J. Biol. Chem.* 266 (1991) 13370–13376.
- [101] D.H. Engels, A. Lott, G.H. Schmid, E.K. Pistorious, Inactivation of the water-oxidizing enzyme in manganese stabilizing protein-free cells of the cyanobacteria *Synechococcus* PCC 7942 and *Synechocystis* PCC 6803 during dark incubation and conditions leading to photoactivation, *Photosynth. Res.* 42 (1994) 227–244.
- [102] S.P. Mayfield, P. Bennoun, J.D. Rochaix, Expression of the nuclear-encoded OEE1 protein is required for oxygen evolution and stability of Photosystem II particles in *Chlamydomonas reinhardtii*, *EMBO J.* 6 (1987) 313–318.
- [103] X. Yi, M. McChargue, S.M. Laborde, L.K. Frankel, T.M. Bricker, The manganese-stabilizing protein is required for Photosystem II assembly/stability and photoautotrophy in higher plants, *J. Biol. Chem.* 280 (2005) 16170–16174.
- [104] R.S. Hutchinson, J.J. Steenhuis, C.F. Yocum, M.R. Razeghifard, B.B. Barry, Deprotonation of the 33-kDa, extrinsic, manganese-stabilizing subunit accompanies photooxidation of the manganese in Photosystem II, *J. Biol. Chem.* 274 (1999) 31987–31995.
- [105] S.K. Hong, S.A. Pawlikowski, K.A. VanderMeulen, C.F. Yocum, The oxidation state of the Photosystem II manganese cluster influences the structure of the manganese-stabilizing protein, *Biochim. Biophys. Acta* 1504 (2001) 262–274.
- [106] H. Popelkova, S.D. Betts, N. Lydakis Simantiris, M.M. Im, E. Swenson, C.F. Yocum, Mutagenesis of basic residues R151 and R161 in manganese-stabilizing protein of Photosystem II causes inefficient binding of chloride to the oxygen-evolving complex, *Biochemistry* 45 (2006) 3107–3115.
- [107] J.L. Roose, C.F. Yocum, H. Popelkova, Function of PsbO, the Photosystem II manganese-stabilizing protein: probing the role of aspartic acid 157, *Biochemistry* 49 (2010) 6042–6051.
- [108] J.W. Murray, J. Barber, Structural characteristics of channels and pathways in Photosystem II including the identification of an oxygen channel, *J. Struct. Biol.* 159 (2007) 228–237.
- [109] F.M. Ho, Uncovering channels in Photosystem II by computer modelling: current progress, future prospects, and lessons from analogous systems, *Photosynth. Res.* 98 (2008) 503–522.
- [110] F.M. Ho, S. Styring, Access channels and methanol binding site to the CaMn₄ cluster in Photosystem II based on solvent accessibility simulations, with implications for substrate water access, *Biochim. Biophys. Acta* 1777 (2008) 140–153.
- [111] A. Gabdulkhakov, A. Guskov, M. Broser, J. Kern, F. Müh, W. Saenger, A. Zouni, Probing the accessibility of the Mn₄Ca cluster in Photosystem II; channels

- calculation, noble gas derivatization, and cocrystallization with DMSO, *Structure* 17 (2009) 1223–1234.
- [112] C. Spetea, T. Hundal, B. Lundin, M. Hedddad, I. Adamska, B. Andersson, Multiple evidence for nucleotide metabolism in the chloroplast thylakoid lumen, *Proc. Natl. Acad. Sci. U.S.A.* 101 (2004) 1409–1414.
- [113] B. Lundin, S. Thuswaldner, T. Shutova, S. Eshaghi, G. Samuelsson, J. Barber, B. Andersson, C. Spetea, Subsequent events to GTP binding by the plant PsbO protein: structural changes, GTP hydrolysis and dissociation from the Photosystem II complex, *Biochim. Biophys. Acta* 1767 (2007) 500–508.
- [114] B. Lundin, M. Hansson, B. Schoefs, A.V. Vener, C. Spetea, The *Arabidopsis* PsbO2 protein regulates dephosphorylation and turnover of the Photosystem II reaction center D1 protein, *Plant J.* 49 (2007) 528–539.
- [115] E.-M. Aro, I. Virgin, B. Andersson, Photoinhibition of Photosystem II. Inactivation, protein damage and turnover, *Biochim. Biophys. Acta* 1143 (1993) 113–134.
- [116] E.-M. Aro, M. Suorsa, A. Rokka, Y. Allahverdiyeva, V. Paakkari, A. Saleem, N. Batchkova, E. Rintamäki, Dynamics of Photosystem II: a proteomic approach to thylakoid protein complexes, *J. Exp. Bot.* 56 (2005) 347–356.
- [117] B. Lundin, M. Nurmi, M. Rojas-Stuetz, E.-M. Aro, I. Adamski, C. Spetea, Towards understanding the functional difference between the two PsbO isoforms in *Arabidopsis thaliana*—insights from phenotypic analyses of psbO knockout mutants, *Photosynth. Res.* 98 (2008) 405–414.
- [118] T.M. Bricker, L.K. Frankel, Auxiliary functions of the PsbO, PsbP and PsbQ proteins of higher plant Photosystem II: a critical analysis, *Photochem. Photobiol. B: Biol.* 104 (2011) 165–178.
- [119] T. Jansen, C. Rother, J. Steppuhn, H. Reinke, K. Beyreuther, C. Jansson, B. Andersson, R.G. Herrmann, Nucleotide sequence of cDNA clones encoding the complete 23 kDa and 16 kDa precursor proteins associated with the oxygen-evolving complex from spinach, *FEBS Lett.* 216 (1987) 234–240.
- [120] K. Ifuku, T. Nakatsu, H. Kato, F. Sato, Crystal structure of the PsbP protein of Photosystem II from *Nicotiana tabacum*, *EMBO Rep.* 5 (2004) 362–367.
- [121] J. Kohoutova, L.K. Smatanova, J. Brynda, M. Lapkouski, J.L. Revuelta, J.B. Arellano, R. Ettrich, Crystallization and preliminary crystallographic characterization of the extrinsic PsbP protein of Photosystem II from *Spinacia oleracea*, *Acta Crystallogr. F* 65 (2009) 111–115.
- [122] R. Ristvejova, M. Lapkouski, J. Kopecky, V.K. Hofbauerova, Z. Sofova, S. Gonzalez-Perez, I.K. Smatanova, J.L. Revuelta, J.B. Arellano, R. Ettrich, Structure of PsbP protein from *Spinacia oleracea* at 1.98 Å, doi:10.2210/pdb2vuv4/pdb.
- [123] F. Michoux, K. Takasaka, M. Boehm, P.J. Nixon, J.W. Murray, Structure of CyanoP at 2.8 Å: implications for the evolution and function of the PsbP subunit of Photosystem II, *Biochemistry* 49 (2010) 7411–7413.
- [124] M. Stewart, R.P. Baker, 1.9 Å resolution crystal structure of the *Saccharomyces cerevisiae* Ran-binding protein Mog1p, *J. Mol. Biol.* 299 (2000) 213–223.
- [125] M. Oki, T. Nishimoto, A protein required for nuclear-protein import, Mog1p, directly interacts with GTP-Gsp1p, the *Saccharomyces cerevisiae* ran homologue, *Proc. Natl. Acad. Sci. U.S.A.* 95 (1998) 15388–15393.
- [126] V. Calderone, M. Trabucchi, A. Vujicic, R. Battistutta, G.M. Giacometti, F. Andreucci, R. Barbato, G. Zanotti, Crystal structure of the PsbQ protein of Photosystem II from higher plants, *EMBO Rep.* 4 (2003) 900–905.
- [127] M. Balsera, J.B. Arellano, J.L. Revuelta, J. De Las Rivas, J.A. Hermoso, A 1.49 Å resolution crystal structure of PsbQ from Photosystem II of *Spinacia oleracea* reveals a PP1I structure in the N-terminal region, *J. Mol. Biol.* 350 (2005) 1051–1058.
- [128] S.A. Jackson, R.D. Fagerlund, S.M. Wilbanks, J.J. Eaton-Rye, Crystal structure of PsbQ from *Synechocystis* sp. PCC 6803 at 1.8 Å: implications for binding and function in cyanobacterial Photosystem II, *Biochemistry* 49 (2010) 2765–2767.
- [129] J. Ristvejova, V. Kopecky Jr., Z. Sovova, M. Balsera, J.B. Arellano, M. Green, R. Ettrich, Structure and dynamics of the N-terminal loop of PsbQ from Photosystem II of *Spinacia oleracea*, *Biochem. Biophys. Res. Commun.* 345 (2006) 287–291.
- [130] M. Miyao, N. Murata, The mode of binding of three extrinsic proteins of 33 kDa, 23 kDa and 18 kDa in the Photosystem II complex of spinach, *Biochim. Biophys. Acta* 977 (1989) 315–321.
- [131] K. Kavelaki, D.F. Ghanotakis, Effects of the manganese complex on the binding of the extrinsic proteins (17, 23 and 33 kDa) of Photosystem II, *Photosynth. Res.* 29 (1991) 149–155.
- [132] K. Ifuku, F. Sato, A truncated mutant of the extrinsic 23-kDa protein that absolutely requires the extrinsic 17-kDa protein for Ca^{2+} retention in Photosystem II, *Plant Cell Physiol.* 43 (2002) 1244–1249.
- [133] A. Tohri, N. Dohmae, T. Suzuki, H. Ohta, Y. Inoue, I. Enami, Identification of domains on the extrinsic 23 kDa protein possibly involved in electrostatic interaction with the extrinsic 33 kDa protein in spinach Photosystem II, *Eur. J. Biochem.* 271 (2004) 962–971.
- [134] T.M. Bricker, L.K. Frankel, Carboxylate groups on the manganese-stabilizing protein are required for efficient binding of the 24 kDa extrinsic protein to Photosystem II, *Biochemistry* 42 (2003) 2056–2061.
- [135] T. Kuwabara, T. Murata, M. Miyao, N. Murata, Partial degradation of the 18-kilodalton protein of the photosynthetic oxygen-evolving complex a study of a binding site, *Biochim. Biophys. Acta* 850 (1986) 146–155.
- [136] G.D. Meades Jr., A. McLachlan, L. Sallans, P.A. Limbach, T.M. Bricker, Association of the 17 kDa extrinsic protein with Photosystem II in higher plants, *Biochemistry* 44 (2005) 15216–15221.
- [137] T. Suzuki, J. Minagawa, T. Tomo, K. Sonoike, H. Ohta, I. Enami, Binding and functional properties of the extrinsic proteins in oxygen-evolving Photosystem II particles from a green alga, *Chlamydomonas reinhardtii* having his-tagged CP47, *Plant Cell Physiol.* 44 (2003) 76–84.
- [138] T. Suzuki, O. Tada, M. Makimura, A. Tohri, H. Ohta, Y. Yamamoto, Isolation and characterization of oxygen-evolving Photosystem II complexes retaining the PsbO, P and Q proteins from *Euglena gracilis*, *Plant Cell Physiol.* 45 (2004) 1168–1175.
- [139] Y. Yamamoto, F. Kubota, Specific release of the extrinsic 18-kDa protein from spinach Photosystem II particles by the treatment with NaCl and methanol and its application for large scale purification of the three extrinsic proteins of Photosystem II without chromatography, *Biochim. Biophys. Acta* 893 (1987) 579–583.
- [140] M. Bernier, R. Carpentier, The action of mercury on the binding of the extrinsic polypeptides associated with the water oxidizing complex of Photosystem II, *FEBS Lett.* 360 (1995) 251–254.
- [141] H. Yu, X. Xu, R. Britt, The 33 kDa protein can be removed without affecting the association of the 23 and 17 kDa proteins with the luminal side of PS II of spinach, *Biochemistry* 45 (2006) 3404–3411.
- [142] S. Caffarri, R. Kouril, S. Kereiche, E.J. Boekema, R. Croce, Functional architecture of higher plant Photosystem II supercomplexes, *EMBO J.* 28 (2009) 3052–3063.
- [143] H. Popelkova, C.F. Yocum, Current status of the role of Cl^- ion in the oxygen-evolving complex, *Photosynth. Res.* 93 (2007) 111–121.
- [144] M. Miqyas, H.J. van Gorkom, C.F. Yocum, The PS II calcium site revisited, *Photosynth. Res.* 92 (2007) 275–287.
- [145] H.-E. Akerlund, C. Jansson, B. Andersson, Reconstitution of oxygen evolution in high salt washed Photosystem II particles, *Biochim. Biophys. Acta* 681 (1982) 1–10.
- [146] M. Miyao, N. Murata, Partial disintegration and reconstitution of the photosynthetic oxygen evolution complex, *Biochim. Biophys. Acta* 725 (1983) 87–93.
- [147] M. Miyao, N. Murata, The Cl^- effect on photosynthetic oxygen evolution: interaction of Cl^- with 18-kDa, 24-kDa and 33-kDa proteins, *FEBS Lett.* 180 (1985) 303–308.
- [148] D.F. Ghanotakis, G.T. Babcock, C.F. Yocum, Calcium reconstitutes high rates of oxygen evolution in polypeptide depleted Photosystem II preparations, *FEBS Lett.* 167 (1984) 127–130.
- [149] D.F. Ghanotakis, G.T. Babcock, C.F. Yocum, On the role of water-soluble polypeptides (17, 23 kDa) calcium and chloride in photosynthetic oxygen evolution, *FEBS Lett.* 192 (1985) 1–3.
- [150] H.Y. Nakatani, Photosynthetic oxygen evolution does not require the participation of polypeptides of 16 and 24 kDa, *Biochem. Biophys. Res. Commun.* 120 (1984) 299–304.
- [151] A. Boussac, A.W. Rutherford, Ca^{2+} binding to the oxygen evolving enzyme varies with the redox state of the Mn enzyme, *Biochim. Biophys. Acta* 236 (1988) 432–436.
- [152] P. Adelroth, K. Lindberg, L.-E. Andreasson, Studies of Ca^{+2} binding in spinach Photosystem II using $^{45}\text{Ca}^{+2}$, *Biochemistry* 34 (1995) 9021–9027.
- [153] H. Wincencjusz, C.F. Yocum, H.J. van Gorkom, S-state dependence of chloride binding affinities and exchange dynamics in the intact and polypeptide depleted O_2 evolving complex of Photosystem II, *Biochemistry* 37 (1998) 8595–8604.
- [154] H. Wincencjusz, C.F. Yocum, H.J. van Gorkom, Activating anions that replace chloride in the oxygen-evolving complex of Photosystem II slow the kinetics of the terminal step in water oxidation and destabilize the S_2 and S_3 states, *Biochemistry* 38 (1999) 3719–3725.
- [155] D.F. Ghanotakis, J.N. Topper, C.F. Yocum, Structural organization of the oxidizing side of Photosystem II: exogenous reductants reduce and destroy the Mn-complex in Photosystem II membranes depleted of the 17 and 23 kDa polypeptides, *Biochim. Biophys. Acta* 767 (1984) 524–531.
- [156] X. Yi, H. Liu, S.R. Hargett, L.K. Frankel, T.M. Bricker, The PsbP protein is required for Photosystem II complex assembly/stability and photoautotrophy in *Arabidopsis thaliana*, *J. Biol. Chem.* 34 (2007) 24833–24841.
- [157] J.L. Roose, L.K. Frankel, T.M. Bricker, Documentation of significant electron transport defects on the reducing side of Photosystem II upon removal of the PsbP and PsbQ extrinsic proteins, *Biochemistry* 49 (2010) 36–41.
- [158] X. Yi, S.R. Hargett, L.K. Frankel, T.M. Bricker, The PsbQ protein is required in *Arabidopsis* for Photosystem II assembly/stability and photoautotrophy under low light conditions, *J. Biol. Chem.* 281 (2006) 26260–26267.
- [159] M. Tomita, K. Ifuku, F. Satoh, T. Noguchi, FTIR evidence that the PsbP extrinsic protein induces protein conformational changes around the oxygen-evolving Mn cluster in Photosystem II, *Biochemistry* 48 (2009) 6318–6325.
- [160] T. Ono, S. Izawa, Y. Inoue, Structural and functional modulation of the manganese cluster in Ca^{+2} -depleted Photosystem II by binding of the 24 kDa extrinsic protein, *Biochemistry* 31 (1992) 7648–7655.
- [161] K. Ifuku, T. Nakatsu, R. Shimamoto, Y. Yamamoto, S. Ishihara, H. Kato, F. Sato, Structure and function of the PsbP protein of Photosystem II from higher plants, *Photosynth. Res.* 84 (2005) 251–255.
- [162] K. Ifuku, F. Sato, Importance of the N-terminal sequence of the extrinsic 23 kDa polypeptide in Photosystem II in ion retention in oxygen evolution, *Biochim. Biophys. Acta* 1546 (2001) 196–204.
- [163] A. Villarejo, T. Shutova, O. Moskin, M. Forssen, V.V. Klimov, G. Samuelsson, A Photosystem II-associated carbonic anhydrase regulates the efficiency of photosynthetic oxygen evolution, *EMBO J.* 21 (2002) 1930–1938.
- [164] A.V. Shitov, O.V. Pobeguts, T.N. Smolova, S.I. Allakhverdiev, V.V. Klimov, Manganese-dependent carboxianhydrase activity of Photosystem II proteins, *Biochemistry (Mosc.)* 74 (2009) 509–517.
- [165] B.C. Tripathy, P. Mohanty, Zinc-inhibited electron transport of photosynthesis in isolated barley chloroplasts, *Plant Physiol.* 66 (1980) 1174–1178.
- [166] R.M. Chaloub, C.C.P. de Magalhães, C.P. dos Santos, Early toxic effects of zinc on PSII of *Synechococcus aquatilis* F. aquatilis (Cyanophyceae), *J. Phycol.* 41 (2005) 1162–1168.

- [167] S. Reiland, G. Messerli, K. Baerenfaller, B. Gerrits, A. Endler, J. Grossmann, W. Gruissem, S. Baginsky, Large-scale *Arabidopsis* phosphoproteome profiling reveals novel chloroplast kinase substrates and phosphorylation networks, *Plant Physiol.* 150 (2009) 889–903.
- [168] K. Lohrig, B. Müller, J. Davydova, D. Leister, D.A. Wolters, Phosphorylation site mapping of soluble proteins: bioinformatical filtering reveals potential plastidic phosphoproteins in *Arabidopsis thaliana*, *Planta* 229 (2009) 1123–1134.
- [169] K. Ifuku, J. Yamamoto, T.-a. Ono, S. Ishihara, F. Sato, PsbP protein, but not PsbQ protein, is essential for the regulation and stabilization of Photosystem II in higher plants, *Plant Physiol.* 139 (2005) 1175–1184.
- [170] K. Ido, K. Ifuku, Y. Yamamoto, S. Ishihara, A. Murakami, K. Takabe, C. Miyake, F. Sato, Knockdown of PsbP protein does not prevent assembly of the dimeric PSII core complex but impairs accumulation of Photosystem II supercomplexes in tobacco, *Biochim. Biophys. Acta* 1787 (2009) 873–881.
- [171] X. Yi, S. Hargett, L.K. Frankel, T.M. Bricker, The PsbP protein, but not the PsbQ protein, is required for normal thylakoid membrane architecture in *Arabidopsis thaliana*, *FEBS Lett.* 583 (2009) 2142–2147.
- [172] W.F. Ettinger, S.M. Theg, Physiologically active chloroplasts contain pools of unassembled extrinsic proteins of the photosynthetic oxygen-evolving enzyme complex in the thylakoid lumen, *J. Cell Biol.* 115 (1991) 321–328.
- [173] A. Hashimoto, W.F. Ettinger, Y. Yamamoto, S.M. Theg, Assembly of newly imported oxygen-evolving complex subunits in isolated chloroplasts: sites of assembly and binding mechanism, *Plant Cell* 9 (1997) 441–452.
- [174] C. de Vitry, J. Olive, D. Drapier, M. Recouvreur, F.A. Wollman, Post-translational events leading to the assembly of Photosystem II protein complex a study using photosynthesis mutants from *Chlamydomonas reinhardtii*, *J. Cell Biol.* 109 (1989) 991–1006.
- [175] S.P. Mayfield, M. Schirmer-Rahire, G.Z. Frandk, J.D. Rochaix, Analysis of the genes of the OEE1 and OEE3 proteins of the Photosystem II complex from *Chlamydomonas reinhardtii*, *Plant Mol. Biol.* 12 (1989) 683–693.
- [176] U. Ljungberg, H.-E. Åkerlund, C. Larsson, B. Andersson, Identification of polypeptides associated with the 23 and 33 kDa proteins of photosynthetic oxygen evolution, *Biochim. Biophys. Acta* 767 (1984) 145–152.
- [177] A. Webber, L.C. Packman, J.C. Gray, A 10 kDa polypeptide associated with the oxygen-evolving complex of Photosystem II has a putative C-terminal non-cleavable thylakoid transfer domain, *FEBS Lett.* 242 (1989) 435–438.
- [178] U. Ljungberg, H.E. Åkerlund, B. Andersson, Isolation and characterization of the 10-kDa and 22-kDa polypeptides of higher-plant Photosystem 2, *Eur. J. Biochem.* 158 (1986) 477–478.
- [179] R. Harrer, R. Bassi, M.G. Testi, C. Schafer, Nearest-neighbor analysis of a Photosystem II complex from *Marchantia polymorpha* L. (liverwort), which contains reaction center and antenna proteins, *Eur. J. Biochem.* 255 (1998) 196–205.
- [180] M. Suorsa, S. Sirpio, Y. Allahverdiyeva, V. Paakkari, F. Mamedov, S. Styring, E. Aro, PsbR, a missing link in the assembly of the oxygen-evolving complex of plant Photosystem II, *J. Biol. Chem.* 281 (2006) 145–150.
- [181] M. Hager, M. Hermann, K. Biehler, A. Krieger-Liszky, R. Bock, Lack of the small plastid-encoded PsbJ polypeptide results in a defective water-splitting apparatus of Photosystem II, reduced photosystem I levels, and hypersensitivity to light, *J. Biol. Chem.* 277 (2002) 14031–14039.
- [182] M. Suorsa, R.E. Regel, V. Paakkari, N. Battchikova, R.G. Herrmann, E.-M. Aro, Protein assembly of Photosystem II and accumulation of subcomplexes in the absence of low molecular mass subunits PsbL and PsbJ, *Eur. J. Biochem.* 271 (2003) 96–107.
- [183] Y.B.B. Zhang, I-TASSER server for protein 3D structure prediction, *BMC Bioinformatics* 9 (2008) 40–48.
- [184] A. Roy, A. Kucukural, Y. Zhang, I-TASSER: a unified platform for automated protein structure and function prediction, *Nat. Protoc.* 5 (2010) 725–738.
- [185] J. Stockhaus, M. Hofer, G. Renger, P. Westhoff, T. Wydrzynski, L. Willmitzer, Antisense RNA efficiently inhibits formation of the 10 kD polypeptide of Photosystem II in transgenic potato plants: analysis of the role of the 10 kD protein, *EMBO J.* 9 (1990) 3013–3021.
- [186] Y. Allahverdiyeva, F. Mamedov, M. Suorsa, S. Styring, I. Vass, E.-M. Aro, Insights into the function of PsbR protein in *Arabidopsis thaliana*, *Biochim. Biophys. Acta* 1767 (2007) 677–685.
- [187] H. Liu, L.K. Frankel, T.M. Bricker, Characterization and complementation of a psbR mutant in *Arabidopsis thaliana*, *Arch. Biochem. Biophys.* 489 (2009) 34–40.
- [188] M.V. Turkina, J. Kargul, A. Blanco-Rivero, A. Villarejo, J.B. Barber, A.V. Vener, Environmentally modulated phosphoproteome of photosynthetic membranes in the green alga *Chlamydomonas reinhardtii*, *Mol. Cell. Proteomics* 5 (2006) 1412–1425.
- [189] N. Sugiyama, H. Nakagami, K. Mochida, A. Daudi, M. Tomita, K. Shirasu, Y. Ishihama, Large-scale phosphorylation mapping reveals the extent of tyrosine phosphorylation in *Arabidopsis*, *Mol. Syst. Biol.* 4 (2008) 193.
- [190] P. Pesaresi, M. Pribil, T. Wunder, D. Leister, Dynamics of reversible protein phosphorylation in thylakoids of flowering plants: the roles of STN7, STN8 and TAP38, *Biochim. Biophys. Acta* 1807 (2011) 887–896.
- [191] N. Kamiya, J.-R. Shen, Crystal structure of oxygen-evolving Photosystem II from *Thermosynechococcus vulcanus* at 3.7 Å resolution, *Proc. Natl. Acad. Sci. U.S.A.* 100 (2003) 98–103.
- [192] A.C. Stewart, U. Ljungberg, H.-E. Åkerlund, B. Anderson, Studies on the polypeptide composition of the cyanobacterial oxygen-evolving complex, *Biochim. Biophys. Acta* 808 (1985) 353–362.
- [193] J.-R. Shen, M. Ikeuchi, Y. Inoue, Stoichiometric association of extrinsic cytochrome c_{550} and 12 kDa protein with a highly purified oxygen evolving Photosystem II core complex from *Synechococcus vulcanus*, *FEBS Lett.* 301 (1992) 145–149.
- [194] I. Enami, H. Murayama, H. Ohta, M. Kamo, K. Nakazato, J.R. Shen, Isolation and characterization of a Photosystem II complex from the red alga *Cyanidium caldarium*: association of cytochrome c_{550} and a 12 kDa protein with the complex, *Biochim. Biophys. Acta* 1232 (1995) 208–216.
- [195] H. Ohta, A. Okumura, S. Okuyama, A. Akiyama, M. Iwai, S. Yoshihara, J.-R. Shen, M. Kamo, I. Enami, Cloning, expression of the psbU gene, and functional studies of the recombinant 12-kDa protein of Photosystem II from a red alga *Cyanidium caldarium*, *Biochem. Biophys. Res. Commun.* 260 (1999) 245–250.
- [196] I. Enami, T. Suzuki, O. Tada, Y. Nakada, K. Nakamura, A. Tohri, H. Ohta, I. Inoue, J.-R. Shen, Distribution of the extrinsic proteins as a potential marker for the evolution of photosynthetic oxygen-evolving Photosystem II, *FEBS J.* 272 (2005) 5020–5030.
- [197] R. Nagao, A. Moriguchi, T. Tomo, A. Niikura, S. Nakajima, T. Suzuki, A. Okumura, M. Iwai, J.-R. Shen, M. Ikeuchi, I. Enami, Binding and functional properties of five extrinsic proteins in oxygen-evolving Photosystem II from a marine centric diatom, *Chaetoceros gracilis*, *J. Biol. Chem.* 258 (2010) 29191–29199.
- [198] B. Thangaraj, C.M. Ryan, P. Souda, K. Krause, K.F. Faull, A.P.M. Weber, P. Fromme, J.P. Whitelegge, Data-directed top-down Fourier-transform mass spectrometry of a large integral membrane protein complex: Photosystem II from *Galdieria sulphuraria*, *Proteomics* 10 (2010) 3644–3656.
- [199] J.-R. Shen, Y. Inoue, Binding and functional properties of two new extrinsic components, cytochrome c_{550} and a 12-kDa protein, in cyanobacterial Photosystem II, *Biochemistry* 32 (1993) 1825–1832.
- [200] J.-R. Shen, M. Ikeuchi, Y. Inoue, Analysis of the psbU gene encoding the 12-kDa extrinsic protein of Photosystem II and studies on its role by deletion mutagenesis in *Synechocystis* sp. PCC 6803, *J. Biol. Chem.* 272 (1997) 17821–17826.
- [201] J.J. Eaton-Rye, J.A. Shand, W.S. Nicoll, pH-dependent photoautotrophic growth of specific Photosystem II mutants lacking lumenal extrinsic polypeptides in *Synechocystis* sp. PCC 6803, *FEBS Lett.* 543 (2003) 148–153.
- [202] T.C. Summerfield, J.J. Eaton-Rye, L.A. Sherman, Global gene expression of a Δ PsbO: Δ PsbU mutant and a spontaneous revertant in the cyanobacterium *Synechocystis* sp. PCC 6803, *Photosynth. Res.* 94 (2007) 265–274.
- [203] S. Clarke, J. Eaton-Rye, Mutation of 363-Phe in the Photosystem II protein CP47 impairs photoautotrophic growth, alters the chloride requirement, and prevents photosynthesis in the absence of either PSII-O or PSII-V in *Synechocystis* sp. PCC 6803, *Biochemistry* 38 (1999) 2707–2715.
- [204] N. Inoue-Kashino, Y. Kashino, K. Satoh, I. Terashima, H.B. Pakrasi, PsbU provides a stable architecture for the oxygen-evolving system in cyanobacterial Photosystem II, *Biochemistry* 44 (2005) 12214–12228.
- [205] J.-R. Shen, M. Qian, Y. Inoue, R.L. Burnap, Functional characterization of *Synechocystis* sp. PCC 6803 Δ psbU and Δ psbV mutants reveals important roles of cytochrome c_{550} in cyanobacterial oxygen evolution, *Biochemistry* 37 (1998) 1551–1558.
- [206] A. Kimura, J.J. Eaton-Rye, E.H. Morita, Y. Nishiyama, H. Hayashi, Protection of the oxygen-evolving machinery by the extrinsic proteins of Photosystem II is essential for development of cellular thermotolerance in *Synechocystis* sp. PCC 6803, *Plant Cell Physiol.* 43 (2002) 932–938.
- [207] J. Veerman, F.K. Bentley, J.J. Eaton-Rye, C.W. Mullineaux, S. Vasil'ev, D. Bruce, The PsbU subunit of Photosystem II stabilizes energy transfer and primary photochemistry in the phycobilisome-Photosystem II assembly of *Synechocystis* sp. PCC 6803, *Biochemistry* 44 (2005) 16939–16948.
- [208] I. Balint, J. Bhattacharya, A. Perelman, D. Schatz, Y. Moskovitz, N. Keren, R. Schwarz, Inactivation of the extrinsic subunit of Photosystem II, PsbU, in *Synechococcus* PCC 7942 results in elevated resistance to oxidative stress, *FEBS Lett.* 580 (2006) 2117–2122.
- [209] P. Mohanty, V.D. Kreslavski, D.A. Los, V.V. Klimov, M. Mimuro, R. Carpentier, S.I. Allakhverdiev, Heat stress: susceptibility, recovery and regulation, in: J.J. Eaton-Rye, B.C. Tripathy, T.D. Sharkey (Eds.), *Photosynthesis: Plastid Biology, Energy Conversion and Plastid Biology, Advances in Photosynthesis and Respiration*, vol. 34, Springer, Dordrecht, in press.
- [210] Y. Nishiyama, H. Hayashi, T. Watanabe, N. Murata, Photosynthetic oxygen evolution is stabilized by cytochrome c_{550} against heat inactivation in *Synechococcus* sp. PCC 7002, *Plant Physiol.* 105 (1994) 1313–1319.
- [211] Y. Nishiyama, D.A. Los, H. Hayashi, N. Murata, Thermal protection of the oxygen evolving machinery by PsbU, an extrinsic protein of Photosystem II, in *Synechococcus* species PCC 7002, *Plant Physiol.* 115 (1997) 1473–1480.
- [212] Y. Nishiyama, D.A. Los, N. Murata, PsbU, a protein associated with Photosystem II, is required for the acquisition of cellular thermotolerance in *Synechococcus* species PCC 7002, *Plant Physiol.* 120 (1999) 301–308.
- [213] C.F. Yocum, The calcium and chloride requirements for the O_2 evolving complex, *Coord. Chem. Rev.* 252 (2008) 296–305.
- [214] A. Okumura, H. Ohta, Y. Inoue, I. Enami, Identification of functional domains of the extrinsic 12 kDa protein in red algal PSII by limited proteolysis and directed mutagenesis, *Plant Cell Physiol.* 42 (2001) 1331–1337.
- [215] A. Okumura, M. Sano, T. Suzuki, H. Tanaka, R. Nagao, K. Nakazato, M. Iwai, H. Adachi, J.-R. Shen, I. Enami, Aromatic structure of tyrosine-92 in the extrinsic PsbU protein of red algal Photosystem II is important for its functioning, *FEBS Lett.* 581 (2007) 5255–5258.
- [216] N. Knoepfle, T.M. Bricker, C. Putnam-Evans, Site-directed mutagenesis of basic arginine residues 305 and 342 in the CP 43 protein of Photosystem II affects oxygen-evolving activity in *Synechocystis* 6803, *Biochemistry* 38 (1999) 1582–1588.

- [217] A. Young, M. McChargue, T.M. Bricker, L.K. Frankel, C. Putnam-Evans, Alterations of the oxygen-evolving apparatus in an ³⁰⁵Arg to ³⁰⁵S mutant in the CP43 protein of Photosystem II from *Synechocystis* sp. PCC 6803 under chloride-limiting conditions, *Biochemistry* 41 (2002) 15747–15753.
- [218] T.M. Bricker, A. Young, L.K. Frankel, C. Putnam-Evans, Introduction of the ³⁰⁵Arg → ³⁰⁵Ser mutation in the large extrinsic loop E of the CP43 protein of *Synechocystis* sp. PCC 6803 leads to the loss of cytochrome *c*₅₅₀ binding to Photosystem II, *Biochim. Biophys. Acta* 1556 (2002) 92–96.
- [219] I. Enami, S. Yoshihara, A. Tohri, A. Okumura, H. Ohta, J.R. Shen, Cross-reconstitution of various extrinsic proteins and Photosystem II complexes from cyanobacteria, red algae and higher plants, *Plant Cell Physiol.* 41 (2000) 1354–1364.
- [220] I. Enami, M. Iwai, A. Akiyama, T. Suzuki, A. Okumura, T. Katoh, O. Tada, H. Ohta, J.-R. Shen, Comparison of binding and functional properties of two extrinsic components, cyt *c*₅₅₀ and a 12 kDa protein, in cyanobacterial PSII with those in red algal PSII, *Plant Cell Physiol.* 44 (2003) 820–827.
- [221] R.W. Holton, J. Myers, Cytochromes of a blue-green alga: extraction of a c-type with a strongly negative redox potential, *Science* 142 (1963) 234–235.
- [222] J.-R. Shen, M. Ikeuchi, Y. Inoue, Stoichiometric association of extrinsic cytochrome *c*₅₅₀ and the 12 kDa protein with a highly purified oxygen-evolving PS II core complex from *Synechococcus vulcanus*, *FEBS Lett.* 301 (1992) 145–149.
- [223] C. Frazao, F.J. Enguita, R. Coelho, G.M. Sheldrick, J.A. Navarro, M. Hervas, M.A. De la Rosa, M.A. Carrondo, Crystal structure of low-potential cytochrome *c*₅₄₉ from *Synechocystis* sp. PCC 6803 at 1.21 Å resolution, *J. Biol. Inorg. Chem.* 6 (2001) 324–332.
- [224] M.R. Sawaya, D.W. Krogmann, A. Serag, K.K. Ho, T.O. Yeates, C.A. Kerfeld, Structures of cytochrome *c*₅₄₉ and cytochrome *c*₆ from the cyanobacterium *Arthrospira maxima*, *Biochemistry* 40 (2001) 9215–9225.
- [225] C.A. Kerfeld, M.R. Sawaya, H. Bottin, K.T. Tran, M. Sugiura, D. Cascio, A. Desbois, T.O. Yeates, D. Kirilovsky, A. Boussac, Structural and EPR characterization of the soluble form of cytochrome *c*₅₅₀ and of the *psbV2* gene product from the cyanobacterium *Thermosynechococcus elongatus*, *Plant Cell Physiol.* 44 (2003) 697–706.
- [226] L. Bumba, H. Havelkova-Dousova, M. Husak, F. Vacha, Structural characterization of Photosystem II complex from red alga *Porphyridium cruentum* retaining extrinsic subunits of the oxygen-evolving complex, *Eur. J. Biochem.* 271 (2004) 2967–2975.
- [227] J.-R. Shen, R.L. Burnap, Y. Inoue, An independent role of cytochrome *c*-550 in cyanobacterial Photosystem II as revealed by double-deletion mutagenesis of the *psbO* and *psbV* genes in *Synechocystis* sp. PCC 6803, *Biochemistry* 34 (1995) 12661–12668.
- [228] T. Morgan, J. Shand, S. Clarke, J. Eaton-Rye, Specific requirements for cytochrome *c*-550 and the manganese-stabilizing protein in photoautotrophic strains of *Synechocystis* sp. PCC 6803 with mutations in the domain ³⁵¹Gly to ⁴³⁶Thr of the chlorophyll binding protein CP47, *Biochemistry* 37 (1998) 14437–14449.
- [229] H. Katoh, S. Itoh, J.R. Shen, M. Ikeuchi, Functional analysis of *psbV* and a novel c-type cytochrome gene *psbV2* of the thermophilic cyanobacterium *Thermosynechococcus elongatus* Strain BP-1, *Plant Cell Physiol.* 42 (2001) 599–607.
- [230] D. Kirilovsky, M. Roncel, A. Boussac, A. Wilson, J.L. Zurita, J.M. Ducruet, H. Bottin, M. Sugiura, J.M. Ortega, A.W. Rutherford, Cytochrome *c*₅₅₀ in the cyanobacterium *Thermosynechococcus elongatus*: study of redox mutants, *J. Biol. Chem.* 279 (2004) 52869–52880.
- [231] H. Katoh, N. Hagino, A.R. Grossman, T. Ogawa, Genes essential to iron transport in the cyanobacterium *Synechocystis* sp. strain PCC 6803, *J. Bacteriol.* 183 (2001) 2779–2784.
- [232] G.W. Pettigrew, G.R. Moore, Cytochromes c: Biological Aspects, Springer, Berlin Heidelberg New York, 1987.
- [233] J. Alam, J. Sprinkle, M.A. Hermodson, D.W. Krogmann, Characterization of cytochrome *c*-550 from cyanobacteria, *Biochim. Biophys. Acta* 766 (1984) 317–321.
- [234] C.W. Hoganson, G. Lagenfeldt, L.-E. Andréasson, EPR and redox potentiometric studies of cytochrome *c*-549 of *Anacystis nidulans*, *Biochim. Biophys. Acta* 1016 (1990) 203–206.
- [235] P.F. Kienzl, G.A. Peschke, Cytochrome *c*-549 — an endogenous cofactor of cyclic photophosphorylation in the cyanobacterium *Anacystis nidulans*? *FEBS Lett.* 162 (1983) 76–80.
- [236] D.W. Krogmann, S. Smith, Low potential cytochrome *c*₅₅₀ function in cyanobacteria and algae, *Curr. Res. Photosynth.* 2 (1990) 687–690.
- [237] L.Z. Morand, R.H. Cheng, D.W. Krogmann, K.K. Ho, Soluble electron transfer catalysts of cyanobacteria, in: D.A. Bryant (Ed.), *The Molecular Biology of Cyanobacteria*, Advances in Photosynthesis, vol. 1, Kluwer Academic Publishers, Dordrecht, 1994, pp. 381–407.
- [238] J.S. Vrettos, M.J. Reifler, O. Kievet, K.V. Lakshmi, J.C. de Paula, G.W. Brudvig, Factors that determine the unusually low reduction potential of cytochrome *c*₅₅₀ in cyanobacterial Photosystem II, *J. Biol. Inorg. Chem.* 6 (2001) 708–716.
- [239] M. Roncel, A. Boussac, J.L. Zurita, H. Bottin, M. Sugiura, D. Kirilovsky, J.M. Ortega, Redox properties of the Photosystem II cytochromes *b*₅₅₉ and *c*₅₅₀ in the cyanobacterium *Thermosynechococcus elongatus*, *J. Biol. Inorg. Chem.* 8 (2003) 206–216.
- [240] H. Ishikita, E.W. Knapp, Redox potential of cytochrome *c*₅₅₀ in the cyanobacterium *Thermosynechococcus elongatus*, *FEBS Lett.* 579 (2005) 3190–3194.
- [241] H. Andrews, Z. Li, A. Altuve-Blanco, M. Rivera, R.L. Burnap, Expression, mutagenesis, and characterization of recombinant low-potential cytochrome *c*₅₅₀ of Photosystem II, *Biochemistry* 44 (2005) 6092–6100.
- [242] F. Guerrero, A. Sedoud, D. Kirilovsky, A.W. Rutherford, J.M. Ortega, M. Roncel, A high redox potential form of cytochrome *c*₅₅₀ in Photosystem II from *Thermosynechococcus elongatus*, *J. Biol. Chem.* 286 (2011) 5985–5994.
- [243] Y. Kashino, W.M. Lauber, J.A. Carroll, Q. Wang, J. Whitmarsh, K. Satoh, H.B. Pakrasi, Proteomic analysis of a highly active Photosystem II preparation from the cyanobacterium *Synechocystis* sp. PCC 6803 reveals the presence of novel polypeptides, *Biochemistry* 41 (2002) 8004–8012.
- [244] T. Ujihara, I. Sakurai, N. Mizusawa, H. Wada, A method for analyzing lipid-modified proteins with mass spectrometry, *Anal. Biochem.* 374 (2008) 429–431.
- [245] R.D. Fagerlund, J.J. Eaton-Rye, The lipoproteins of cyanobacterial Photosystem II, *J. Photochem. Photobiol. B: Biol.* 104 (2011) 191–203.
- [246] T.M. Bricker, J. Morvant, N. Masri, H. Sutton, L.K. Frankel, Isolation of a highly active Photosystem II preparation from *Synechocystis* 6803 using a histidine-tagged mutant of CP 47, *Biochim. Biophys. Acta* 1409 (1998) 50–57.
- [247] M.M. Nowaczyk, R. Hebel, E. Schlodder, H.E. Meyer, B. Warscheid, M. Rogner, Psb27, a cyanobacterial lipoprotein, is involved in the repair cycle of Photosystem II, *Plant Cell* 18 (2006) 3121–3131.
- [248] P.J. Nixon, F. Michoux, J. Yu, M. Boehm, J. Komenda, Recent advances in understanding the assembly and repair of Photosystem II, *Ann. Bot.* 106 (2010) 1–16.
- [249] J.J. Eaton-Rye, Requirements for different combinations of the extrinsic proteins in specific cyanobacterial Photosystem II mutants, *Photosynth. Res.* 84 (2005) 275–281.
- [250] J. De Las Rivas, M. Baklsera, J. Barber, Evolution of oxygenic photosynthesis: genome wide analysis of the OEC proteins, *Trends Plant Sci.* 9 (2004) 18–25.
- [251] N. Sato, Phylogenomic and structural modeling analyses of the PsbP superfamily reveal multiple small segment additions in the evolution of Photosystem II-associated PsbP protein in green plants, *Mol. Phylogenet. Evol.* 56 (2010) 176–186.
- [252] S. Ishihara, A. Takabayashi, K. Ido, T. Endo, K. Ifuku, F. Sato, Distinct functions for the two PsbP-like proteins PPL1 and PPL2 in the chloroplast lumen of *Arabidopsis*, *Plant Physiol.* 145 (2007) 668–679.
- [253] N. Bondarava, A. Krieger-Liszka, Manganese binding to the 23 kDa extrinsic protein of Photosystem II, *Biochim. Biophys. Acta* 1767 (2007) 583–588.
- [254] E. Haag, J.J. Eaton-Rye, G. Renger, W.F.J. Vermaas, Functionally important domains of the large hydrophilic loop of CP 47 as probed by oligonucleotide-directed mutagenesis in *Synechocystis* sp. PCC 6803, *Biochemistry* 32 (1993) 4444–4454.
- [255] P. Beneš, E. Chovancová, B. Kozlíková, A. Pavelka, O. Strnad, J. Brezovský, V. Šustr, M. Klvaňa, T. Szabó, A. Gora, M. Zamborský, L. Biedermannová, P. Medek, J. Damborský, J. Sochor, CAVER 2.1, Software (2010).
- [256] T.C. Summerfield, R.T. Winter, J.J. Eaton-Rye, Investigation of a requirement for the PsbP-like protein in *Synechocystis* sp. PCC 6803, *Photosynth. Res.* 84 (2005) 263–268.
- [257] J.-R. Shen, W. Vermaas, Y. Inoue, The role of cytochrome *c*-550 as studied through reverse genetics and mutant characterization in *Synechocystis* sp. PCC 6803, *J. Biol. Chem.* 270 (1995) 6901–6907.
- [258] D. Sveshnikov, C. Funk, W.P. Schröder, The PsbP-like protein (sl11418) of *Synechocystis* sp. PCC 6803 stabilises the donor side of Photosystem II, *Photosynth. Res.* 93 (2007) 101–109.
- [259] W.L. DeLano, The PyMOL molecular graphics system, Software (2002).
- [260] M. Petřek, M. Otyepka, P. Banáš, P. Košinová, J. Koča, CAVER: a new tool to explore routes from protein clefts, pockets and cavities, *BMC Bioinformatics* 7 (2006) 316.