

9-1-2015

## Stochastic dynamics of determinantal processes by integration by parts


Laurent Decreusefond

Ian Flint

Nicolas Privault

Giovanni Luca Torrisi

Follow this and additional works at: <https://digitalcommons.lsu.edu/cosa>

 Part of the [Analysis Commons](#), and the [Other Mathematics Commons](#)

---

### Recommended Citation

Decreusefond, Laurent; Flint, Ian; Privault, Nicolas; and Torrisi, Giovanni Luca (2015) "Stochastic dynamics of determinantal processes by integration by parts," *Communications on Stochastic Analysis*: Vol. 9 : No. 3 , Article 5.

DOI: 10.31390/cosa.9.3.05

Available at: <https://digitalcommons.lsu.edu/cosa/vol9/iss3/5>

## STOCHASTIC DYNAMICS OF DETERMINANTAL PROCESSES BY INTEGRATION BY PARTS

LAURENT DECREUSEFOND, IAN FLINT, NICOLAS PRIVAULT,  
AND GIOVANNI LUCA TORRISI

ABSTRACT. We derive an integration by parts formula for functionals of determinantal processes on compact sets, completing the arguments of [4]. This is used to show the existence of a configuration-valued diffusion process which is non-colliding and admits the distribution of the determinantal process as reversible law. In particular, this approach allows us to build a concrete example of the associated diffusion process, providing an illustration of the results of [4] and [30].

### 1. Introduction

Determinantal processes are point processes that exhibit repulsion, and were introduced to represent the configuration of fermions, cf. [17, 21, 25]. They are known to be connected with the zeros of analytic functions (cf. [13] and references therein) as well as with the theory of random matrices (cf. [3]). More recently, in [8, 28, 29], different authors have used determinantal processes to model phenomena arising in telecommunication networks.

The Markov process associated to the Ornstein-Uhlenbeck operator on the Poisson space has been constructed in [24]. In [1], using the Dirichlet forms theory (see [9, 15]), is constructed the diffusion whose symmetrizing measure is the law of a Poisson process over a Riemannian manifold. This result has been extended to Gibbs processes on  $\mathbb{R}^d$  in [2]. Using such methods, the Dirichlet form and diffusion process associated to determinantal processes has been constructed in [30].

In this paper, by completing the arguments of [4] we prove an integration by parts formula for functionals of determinantal processes on compact subsets of  $\mathbb{R}^d$ , and we recover the closability of the associated Dirichlet form. This provides a novel proof of the existence of interacting diffusion processes properly associated to determinantal processes. In addition, our approach based on integration by parts exhibits the generator of the diffusion process, and allows in turn to provide an explicit example of a diffusion process satisfying our hypotheses.

---

Received 2015-8-5; Communicated by the editors.

2010 *Mathematics Subject Classification*. Primary 60G55; Secondary ; 60J60; 60G60; 60H07; 60K35.

*Key words and phrases*. Dirichlet forms, diffusion processes, integration by parts, Malliavin calculus, determinantal processes.

This research is supported by ANR Masterie ANR-10-BLAN-121 and MOE Tier 2 grant MOE2012-T2-2-033 ARC 3/13.

As a preliminary step, we derive an integration by parts formula for functionals of a determinantal process on a compact set  $D \subset \mathbb{R}^d$ , by completing the result established in [4]. In comparison with [4], the integration by parts formula on compact sets is extended to closed gradient and divergence operators by the use of a different set of test functionals, cf. (3.1) and Theorem 3.9. Our construction of the diffusion processes follows the lines of [1], and it differs from the one of [26] which is based on sample-path identities. Our gradient and divergence operators also differ from those of [19], which also deals with compact subsets of  $\mathbb{R}^d$ . Nevertheless the integration by parts formula of Theorem 3.9 can also be applied to density estimation and sensitivity analysis for functionals of determinantal processes along the same lines.

In Theorem 4.2 we construct the Dirichlet form corresponding to a determinantal process on a compact set  $D \subset \mathbb{R}^d$ . In Theorem 5.1 we show the existence of the diffusion properly associated to a determinantal process on a compact set  $D \subset \mathbb{R}^d$ . Note that, as in the other constructions (cf. [1, 2, 30]), the associated diffusion process admits the distribution of the determinantal process as a reversible law. We prove the non-collision property of the diffusion in Theorem 5.3.

Finally, in Section 6, we provide an example of a determinantal process satisfying our integration by parts formula, and for which the aforementioned properly associated diffusion process exists.

Some definitions related to point processes theory and more in particular to determinantal processes are recalled in Section 2 based on [5, 6, 13, 14, 18]. Some notions from the Dirichlet forms theory are given in Sections 4 and 5 based on [9, 15]. We also refer the reader to [7, 22] for the required background on functional analysis.

## 2. Preliminaries

*Locally finite point processes.* Let  $S$  be a Polish space, and denote by  $\mathcal{B}(S)$  the associated Borel  $\sigma$ -algebra. For any subset  $B \subset S$ , let  $\sharp B$  denote the cardinality of  $B$ , setting  $\sharp B = \infty$  if  $B$  is not finite. We denote by  $N_{lf}$  the set of locally finite point configurations on  $S$ :

$$N_{lf} := \{B \subset S : \sharp(B \cap D) < \infty, \text{ for any compact } D \subset S\}.$$

We identify locally finite configurations with  $\mathbb{N}$ -valued simple Radon measures, equip  $N_{lf}$  with the vague topology (see Appendix 2 in [5]), and we denote the corresponding Borel  $\sigma$ -algebra by  $\mathcal{N}_{lf}$ . We recall that a non-negative simple Radon measure is a Radon measure which is less than or equal to 1 on singletons. We define similarly the set of finite point configurations on  $S$  as  $N_f := \{B \subset S : \sharp B < \infty\}$ , which is naturally equipped with the trace  $\sigma$ -algebra  $\mathcal{N}_f = \mathcal{N}_{lf}|_{N_f}$ . For any measurable set  $B \subset S$ , let  $N_f^B$  be the space of finite configurations on  $B$ , and  $\mathcal{N}_f^B$  the associated trace- $\sigma$ -algebra.

By a locally finite and simple point process  $\mathbf{X}$  on  $S$  we mean a measurable mapping defined on some probability space  $(\Omega, \mathcal{F}, P)$  taking values on  $(N_{lf}, \mathcal{N}_{lf})$ . We denote by  $\mathbf{X}(B)$  the number of points of  $\mathbf{X}$  in a measurable set  $B \subset S$ , i.e.  $\mathbf{X}(B) := \sharp(\mathbf{X} \cap B)$ , and by  $\mathbf{X}^B := \mathbf{X} \cap B = \{X_1, \dots, X_{\mathbf{X}(B)}\}$  the restriction to  $B$

of the point process  $\mathbf{X} \equiv \{X_n\}_{1 \leq n \leq \mathbf{X}(S)}$ . In the following, we shall denote by  $\mathbb{P}$  the law of  $\mathbf{X}$  and by  $\mathbb{P}_B$  the law of  $\mathbf{X}^B$ .

The correlation functions of  $\mathbf{X}$ , with respect to (*w.r.t.*) a given Radon measure  $\nu$  on  $(S, \mathcal{B}(S))$ , are (if they exist) symmetric measurable functions  $\rho_n : S^n \rightarrow \mathbb{R}_+$  such that

$$\mathbb{E} \left[ \prod_{i=1}^n \mathbf{X}(B_i) \right] = \int_{B_1 \times \dots \times B_n} \rho_n(x_1, \dots, x_n) \nu(dx_1) \cdots \nu(dx_n),$$

for any family of mutually disjoint bounded subsets  $B_1, \dots, B_n$  of  $S$ ,  $n \geq 1$ . We require in addition that  $\rho_n(x_1, \dots, x_n) = 0$  whenever  $x_i = x_j$  for some  $1 \leq i \neq j \leq n$ . When  $\rho_1$  exists, the measure  $\rho_1(x) \nu(dx)$  is known as the intensity measure of  $\mathbf{X}$ .

As in [11], we define for any Radon measure  $\nu$  on  $(S, \mathcal{B}(S))$  the  $\nu$ -sample measure  $L^\nu$  on  $(N_f, \mathcal{N}_f)$  by

$$\int_{N_f} f(\alpha) L^\nu(d\alpha) := \sum_{n \geq 0} \frac{1}{n!} \int_{S^n} f(\{x_1, \dots, x_n\}) \nu(dx_1) \cdots \nu(dx_n), \quad (2.1)$$

for any measurable  $f : N_f \rightarrow \mathbb{R}_+$ . For any compact subset  $D \subset S$ , the Janossy densities of  $\mathbf{X}$  *w.r.t.*  $\nu$  are (if they exist) measurable symmetric functions  $j_D^n : D^n \rightarrow \mathbb{R}$  satisfying, for all measurable  $f : N_f^D \rightarrow \mathbb{R}_+$ ,

$$\mathbb{E} [f(\mathbf{X}^D)] = \sum_{n \geq 0} \frac{1}{n!} \int_{D^n} f(\{x_1, \dots, x_n\}) j_D^n(x_1, \dots, x_n) \nu(dx_1) \cdots \nu(dx_n), \quad (2.2)$$

i.e. defining  $j_D(\mathbf{x}) := j_D^{|\mathbf{x}|}(x_1, \dots, x_{|\mathbf{x}|})$  for  $\mathbf{x} = \{x_1, \dots, x_{|\mathbf{x}|}\} \in N_f^D$ ,  $j_D$  is the density of  $\mathbb{P}_D$  with respect to  $L_D^\nu$  (the restriction to  $\mathcal{N}_f^D$  of  $L^\nu$ ), when  $\mathbb{P}_D \ll L_D^\nu$ .

*Kernels and integral operators.* Let  $\nu$  be a Radon measure on  $(S, \mathcal{B}(S))$ . For any compact set  $D \subset S$ , we denote by  $L^2(D, \nu)$  the Hilbert space of complex-valued square integrable functions *w.r.t.* the restriction of the Radon measure  $\nu$  to  $D$ , equipped with the inner product

$$\langle f, g \rangle_{L^2(D, \nu)} := \int_D f(x) \overline{g(x)} \nu(dx), \quad f, g \in L^2(D, \nu)$$

where  $\bar{z}$  denotes the complex conjugate of a complex  $z \in \mathbb{C}$ . By definition, an integral operator  $\mathcal{T} : L^2(S, \nu) \rightarrow L^2(S, \nu)$  with kernel  $T : S^2 \rightarrow \mathbb{C}$  is a bounded operator defined by

$$\mathcal{T}f(x) := \int_S T(x, y) f(y) \nu(dy), \quad \text{for } \nu\text{-almost all } x \in S.$$

Letting  $\mathcal{P}_D$  denote the projection operator from  $L^2(S, \nu)$  onto  $L^2(D, \nu)$ , we set  $\mathcal{T}_D = \mathcal{P}_D \mathcal{T} \mathcal{P}_D$  and note that its kernel is  $T_D(x, y) := \mathbf{1}_D(x) T(x, y) \mathbf{1}_D(y)$ , for  $\nu$ -almost all  $x, y \in S$ . It can be shown that  $\mathcal{T}_D$  is a compact operator. The operator  $\mathcal{T}$  is said to be Hermitian or self-adjoint if

$$T(x, y) = \overline{T(y, x)}, \quad \text{for } \nu^{\otimes 2}\text{-almost all } (x, y) \in S^2. \quad (2.3)$$

Equivalently, this means that the integral operator  $\mathcal{T}_D$  is self-adjoint for any compact set  $D \subset S$ . If  $\mathcal{T}_D$  is self-adjoint, by the spectral theorem we have that  $L^2(D, \nu)$

has an orthonormal basis  $(\varphi_j^D)_{j \geq 1}$  of eigenfunctions of  $\mathcal{T}_D$ . The corresponding eigenvalues  $(\lambda_j^D)_{j \geq 1}$  have finite multiplicity (except possibly the zero eigenvalue) and the only possible accumulation point of the eigenvalues is the eigenvalue zero. Then, the kernel  $T_D$  of  $\mathcal{T}_D$  can be written as

$$T_D(x, y) = \sum_{j \geq 1} \lambda_j^D \varphi_j^D(x) \overline{\varphi_j^D(y)}, \tag{2.4}$$

for  $\nu$ -almost all  $x, y \in D$ . We say that an operator  $\mathcal{T}$  is positive (respectively non-negative) if its spectrum is included in  $(0, +\infty)$  (respectively  $[0, +\infty)$ ). For two operators  $\mathcal{T}$  and  $\mathcal{U}$ , we will say that  $\mathcal{T} > \mathcal{U}$  (respectively  $\mathcal{T} \geq \mathcal{U}$ ) in the operator ordering if  $\mathcal{T} - \mathcal{U}$  is a positive operator (respectively a non-negative operator).

We say that a self-adjoint integral operator  $\mathcal{T}_D$ , with kernel  $T_D$  as in (2.4), is of trace class if  $\sum_{j \geq 1} |\lambda_j^D| < \infty$ , and we define the trace of the operator  $\mathcal{T}_D$  as  $\text{Tr}(\mathcal{T}_D) := \sum_{j \geq 1} \lambda_j^D$ . If  $\mathcal{T}_D$  is of trace class for every compact subset  $D \subset S$ , then we say that  $\mathcal{T}$  is locally of trace class. It is easily seen that  $\mathcal{T}^n$  is locally of trace class, for all  $n \geq 2$ , if  $\mathcal{T}$  is locally of trace class. Finally, we define the Fredholm determinant of  $\mathbf{Id} + \mathcal{T}_D$ , when  $\|\mathcal{T}_D\|_{op} < 1$ , as

$$\text{Det}(\mathbf{Id} + \mathcal{T}_D) := \exp \left( \sum_{n \geq 1} \frac{(-1)^{n-1}}{n} \text{Tr}(\mathcal{T}_D^n) \right). \tag{2.5}$$

Here,  $\mathbf{Id}$  denotes the identity operator on  $L^2(S, \nu)$  and  $\|\cdot\|_{op}$  denotes the operator norm.

*Determinantal processes on  $S$ .* Let  $\mu$  be a Radon measure on  $(S, \mathcal{B}(S))$ . A locally finite and simple point process  $\mathbf{X}$  on  $S$  is said to be a determinantal process with kernel  $K$  and reference measure  $\mu$  if its correlation functions *w.r.t.*  $\mu$  exist and satisfy

$$\rho_n(x_1, \dots, x_n) = \det(K(x_i, x_j))_{1 \leq i, j \leq n},$$

for any  $n \geq 1$  and  $\mu$ -a.e.  $x_1, \dots, x_n \in S$ , where  $K(\cdot, \cdot)$  is a measurable function. Throughout this paper we shall work under the following hypothesis.

**(H1):** *The integral operator  $\mathcal{K}$  on  $L^2(S, \mu)$  with kernel  $K$  is locally of trace class, self-adjoint, and  $0 \leq \mathcal{K} < \mathbf{Id}$ .*

The existence and uniqueness in law of a determinantal process with kernel  $K$  is guaranteed under **(H1)** by the results in [17, 21, 23]. See also Lemma 4.2.6 and Theorem 4.5.5 in [13].

We define the global interaction operator  $\mathcal{J} := (\mathbf{Id} - \mathcal{K})^{-1}\mathcal{K}$ . We also define the local interaction operator  $\mathcal{J}[D] := (\mathbf{Id} - \mathcal{K}_D)^{-1}\mathcal{K}_D$ , and we emphasize that  $\mathcal{J}[D]$  is not the projection of  $\mathcal{J}$  onto  $L^2(D, \mu)$ . We refer the reader to [11] for the following properties of  $\mathcal{J}[D]$ . First,  $\mathcal{J}[D]$  is a self-adjoint integral operator and, letting  $J[D]$  denote its kernel, as a consequence of (2.4) we have

$$J[D](x, y) = \sum_{j \geq 1} \frac{\lambda_j^D}{1 - \lambda_j^D} \varphi_j^D(x) \overline{\varphi_j^D(y)}, \tag{2.6}$$

for  $\mu$ -almost all  $x, y \in D$ . Second,  $\mathcal{J}[D]$  is a trace class operator. Third, denoting by  $\det J[D](\{x_1, \dots, x_n\}) := \det (J[D](x_i, x_j))_{1 \leq i, j \leq n}$ , the function

$$(x_1, \dots, x_n) \mapsto \det J[D](\{x_1, \dots, x_n\})$$

is  $\mu^{\otimes n}$ -a.e. non-negative and symmetric in  $x_1, \dots, x_n$ , and we set

$$\det J[D](\{x_1, \dots, x_n\}) := \det J[D](x_1, \dots, x_n).$$

The local interaction operator is related to the Janossy densities of a determinantal process by the following proposition.

**Proposition 2.1** ([21]). *Under (H1), for any compact  $D \subset S$ , the Janossy densities  $j_D^n(\mathbf{x})$  of  $\mathbf{X}$  are given by*

$$j_D^n(x_1, \dots, x_n) = \text{Det}(\mathbf{Id} - \mathcal{K}_D) \det J[D](x_1, \dots, x_n), \quad x_1, \dots, x_n \in D, \quad n \geq 1. \tag{2.7}$$

Moreover,  $P(\mathbf{X}^D = \emptyset)$  is given by  $j_D^0(\emptyset) = \text{Det}(\mathbf{Id} - \mathcal{K}_D)$ .

### 3. Differential Calculus and Integration by Parts

In this section we derive an integration by parts formula for functionals of a determinantal point process, and we extend it by closability. Hereafter we assume that  $S$  is a domain of  $\mathbb{R}^d$  equipped with the Euclidean distance,  $\mu$  is a Radon measure on  $(S, \mathcal{B}(S))$  and  $D \subset S$  is a fixed compact set. We denote by  $\|\cdot\|$  the Euclidean norm on  $\mathbb{R}^d$ , by  $x \cdot y$  the usual inner product of  $x, y \in \mathbb{R}^d$ , and by  $x^{(i)}$  the  $i$ -th component of  $x \in \mathbb{R}^d$ ,  $i = 1, \dots, d$ .

*Differential calculus.* We denote by  $\mathcal{C}^\infty(D, \mathbb{R}^d)$  the set of all  $\mathcal{C}^\infty$ -vector fields  $v : D \rightarrow \mathbb{R}^d$  and by  $\mathcal{C}^\infty(S^k)$  the set of all  $\mathcal{C}^\infty$ -functions on  $S^k$ .

**Definition 3.1.** A function  $F : N_f^D \rightarrow \mathbb{R}$  is said to be in  $\mathcal{S}_D$  if

$$F(\mathbf{x}) = f_0 \mathbf{1}_{\{\mathbf{x}(D)=0\}} + \sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} f_k(x_1, \dots, x_k), \quad \mathbf{x} = \{x_1, \dots, x_{\mathbf{x}(D)}\} \in N_f^D, \tag{3.1}$$

where  $f_0 \in \mathbb{R}$  is a constant,  $n \geq 1$  is an integer and, for any  $k = 1, \dots, n$ ,  $f_k \in \mathcal{C}^\infty(D^k)$  is a symmetric function.

The set of test functions  $\mathcal{S}_D$  is dense in  $L_D^2 := L^2(N_f^D, \mathbb{P}_D)$  (indeed, it contains the space  $\tilde{\mathcal{S}}_D$  defined in Definition 4.1 below which is dense in  $L_D^2$ , see e.g. [15] p.54). The gradient of  $F \in \mathcal{S}_D$  as in (3.1) is defined by

$$\nabla_x^{N_{if}} F(\mathbf{x}) := \sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} \sum_{i=1}^k \mathbf{1}_{\{x_i\}}(x) \nabla_{x_i} f_k(x_1, \dots, x_k), \quad x \in D, \quad \mathbf{x} \in N_f^D, \tag{3.2}$$

where  $\nabla_y$  denotes the usual gradient on  $\mathbb{R}^d$  with respect to  $y$ . For  $v \in \mathcal{C}^\infty(D, \mathbb{R}^d)$ , we also let

$$\nabla_v^{N_{if}} F(\mathbf{x}) := \sum_{k=1}^{\mathbf{x}(D)} \nabla_{x_k}^{N_{if}} F(\mathbf{x}) \cdot v(x_k) = \sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} \sum_{i=1}^k \nabla_{x_i} f_k(x_1, \dots, x_k) \cdot v(x_i), \tag{3.3}$$

where we recall that the symbol  $\cdot$  denotes the inner product on  $\mathbb{R}^d$ .

*Quasi-invariance.* Next we recall some results from [4], with some complements that make the proofs more precise. Let  $\text{Diff}_0(D)$  be the set of all diffeomorphisms from  $D$  into itself. For  $\phi \in \text{Diff}_0(D)$  and a Radon measure  $\nu$  on  $D$ ,  $\nu_\phi$  denotes the image measure of  $\nu$  by  $\phi$ . For such a  $\phi$ , we define the map  $\mathcal{I}_\phi : L^2(D, \nu_\phi) \rightarrow L^2(D, \nu)$ ,  $\mathcal{I}_\phi(f) = f \circ \phi$ , whose inverse is given by  $\mathcal{I}_\phi^{-1} = \mathcal{I}_{\phi^{-1}}$ . Note that  $\mathcal{I}_\phi$  is an isometry. Given an operator  $\mathcal{T}$  on  $L^2(D, \nu)$ , we define the operator on  $L^2(D, \nu_\phi)$  by  $\mathcal{T}^\phi = \mathcal{I}_\phi^{-1} \mathcal{T} \mathcal{I}_\phi$ . Lastly, for any  $\mathbf{x} = \{x_n\}_{1 \leq n \leq \mathbf{x}(S)} \in N_{lf}$ , we define the map  $\Phi : N_{lf} \rightarrow N_{lf}$ ,  $\Phi(\{x_n\}_{1 \leq n \leq \mathbf{x}(S)}) = \{\phi(x_n)\}_{1 \leq n \leq \mathbf{x}(S)}$ . The following lemmas are proved in [4] and [27]:

**Lemma 3.2.** *Assume (H1), and take  $\phi \in \text{Diff}_0(D)$ . The following properties hold.*

- a)  $\mathcal{K}_D^\phi$  and  $\mathcal{J}[D]^\phi$  are integral operators on  $L^2(D, \mu_\phi)$  with respective kernels  $K_D^\phi(x, y) = K(\phi^{-1}(x), \phi^{-1}(y))$  and  $J[D]^\phi(x, y) = J[D](\phi^{-1}(x), \phi^{-1}(y))$ .
- b)  $\mathcal{K}_D^\phi$  is of trace class and  $\text{Tr}(\mathcal{K}_D^\phi) = \text{Tr}(\mathcal{K}_D)$ .
- c)  $\text{Det}(\mathbf{Id} - \mathcal{K}_D^\phi) = \text{Det}(\mathbf{Id} - \mathcal{K}_D)$ . This translates into the fact that  $P(\mathbf{X}(D) = 0) = P(\Phi(\mathbf{X})(D) = 0)$  which is expected since  $\phi$  is a diffeomorphism.
- d)  $\mathcal{J}[D]^\phi = \mathcal{J}^\phi[D] := (\mathbf{Id} - \mathcal{K}_D^\phi)^{-1} \mathcal{K}_D^\phi$  is the local interaction operator associated with  $\mathcal{K}^\phi$ .

**Lemma 3.3.** *Assume (H1), and let  $\phi \in \text{Diff}_0(D)$ . Then,  $\Phi(\mathbf{X}^D)$  is a determinantal process with integral operator  $\mathcal{K}_D^\phi$  and reference measure  $\mu_\phi$ .*

To prove the quasi-invariance of the determinantal measure restricted to a compact set  $D \subset S$  with respect to the group of diffeomorphisms on  $D$ , we state one last result.

**Lemma 3.4.** *Under (H1), we have  $\det J[D](\mathbf{x}) > 0$ , for  $\mathbb{P}_D$ -a.e.  $\mathbf{x} \in N_f^D$ . However, it does not in general hold that  $\det J[D](\mathbf{x}) > 0$ , for  $L_D^\mu$ -a.e.  $\mathbf{x} \in N_f^D$ .*

*Proof.* Recall that under (H1), we have  $\mathbb{P}_D \ll L_D^\mu$  and

$$j_D(\mathbf{x}) = \frac{d\mathbb{P}_D}{dL_D^\mu}(\mathbf{x}) = \text{Det}(\mathbf{Id} - \mathcal{K}_D) \det J[D](\mathbf{x}),$$

for  $\mathbf{x} \in N_f^D$ . Since  $j_D$  is a density, we clearly have  $j_D(\mathbf{x}) > 0$ , for  $\mathbb{P}_D$ -a.e.  $\mathbf{x} \in N_f^D$ . Hence, since  $\|\mathcal{K}_D\| < 1$ , we have  $\det J[D](\mathbf{x}) > 0$ , for  $\mathbb{P}_D$ -a.e.  $\mathbf{x} \in N_f^D$ . As to the concluding part of the lemma, we notice that in general, one does not have that  $\mathbb{P}_D$  is equivalent to  $L_D^\mu$ . Indeed, consider for example the case where the rank of  $\mathcal{K}_D$  is less than or equal to  $N \geq 1$ . Then,  $j_D^{N+1}(x_1, \dots, x_{N+1}) = 0$ , for  $\mu^{\otimes(N+1)}$ -a.e.  $(x_1, \dots, x_{N+1}) \in D^{N+1}$  (since  $\mathbf{X}^D$  has less than  $N + 1$  points almost surely, see [23] for details). It suffices to define the set  $A := \{B \subset D : \#B = N + 1\}$  which verifies  $\mathbb{P}(A) = 0$  but  $L_D^\mu(A) = \mu(D)/(N + 1)!$ .  $\square$

*Remark 3.5.* If we assume that, for any  $n \geq 1$ , the function  $(x_1, \dots, x_n) \mapsto \det J[D](x_1, \dots, x_n)$  is strictly positive  $\mu^{\otimes n}$ -a.e. on  $D^n$ , then we have that  $\mathbb{P}_D$  and  $L_D^\mu$  are equivalent, and it follows that  $\det J[D](\mathbf{x}) > 0$ , for  $L_D^\mu$ -a.a.  $\mathbf{x} \in N_f^D$ .

The next Proposition 3.6 is similar to its analog in [4], however the proof given there implicitly uses the fact that  $\det J[D](\mathbf{x}) > 0$ , for  $L_D^\mu$ -a.a.  $\mathbf{x} \in \mathbb{N}_f^D$ , which has been shown to be false in general. In order to prove Proposition 3.6, we assume the following technical condition.

**(H2)** : *The Radon measure  $\mu$  is absolutely continuous w.r.t. the Lebesgue measure  $\ell$  on  $D$ , with Radon-Nikodym derivative  $\rho = \frac{d\mu}{d\ell}$  which is strictly positive and continuously differentiable on  $D$ .*

Under **(H2)**, for any  $\phi \in \text{Diff}_0(D)$ ,  $\mu_\phi$  is absolutely continuous with respect to  $\mu$  with density given by

$$p_\phi^\mu(x) = \frac{d\mu_\phi(x)}{d\mu(x)} = \frac{\rho(\phi^{-1}(x))}{\rho(x)} \text{Jac}(\phi^{-1})(x), \tag{3.4}$$

where  $\text{Jac}(\phi^{-1})(x)$  is the Jacobian of  $\phi^{-1}$  at a point  $x \in S$ . We draw attention to the fact that it is indeed  $\text{Jac}(\phi^{-1})(x)$  that appears in (3.4), which differs from equation (2.11) of [1]. We are now in a position to state and prove the main result of this section.

**Proposition 3.6.** *Assume **(H1)**, **(H2)**, and let  $\phi \in \text{Diff}_0(D)$ , for any measurable non-negative  $f$  on  $D$  we have*

$$\begin{aligned} & \mathbb{E} \left[ \exp \left( - \sum_{k=1}^{\mathbf{x}(D)} f \circ \phi(X_k) \right) \right] \\ &= \mathbb{E} \left[ \exp \left( - \sum_{k=1}^{\mathbf{x}(D)} f(X_k) + \sum_{k=1}^{\mathbf{x}(D)} \ln(p_\phi^\mu(X_k)) \right) \frac{\det J^\phi[D](\mathbf{X}^D)}{\det J[D](\mathbf{X}^D)} \right]. \end{aligned}$$

*Proof.* For any measurable non-negative  $f$  on  $D$ , we have

$$\begin{aligned} & \mathbb{E} \left[ \exp \left( - \sum_{k=1}^{\mathbf{x}(D)} f(X_k) + \sum_{k=1}^{\mathbf{x}(D)} \ln(p_\phi^\mu(X_k)) \right) \frac{\det J^\phi[D](\mathbf{X}^D)}{\det J[D](\mathbf{X}^D)} \right] \\ &= \sum_{n \geq 0} \frac{1}{n!} \int_{D^n} e^{-\sum_{k=1}^n f(x_k)} \\ & \quad \times \text{Det}(\mathbf{Id} - \mathcal{K}_D^\phi) \det J^\phi[D](x_1, \dots, x_n) \mu_\phi(dx_1) \cdots \mu_\phi(dx_n), \end{aligned}$$

where we have used (2.7), (3.4) and Lemma 3.2, c). Then, we conclude by Lemma 3.3. Indeed,  $\text{Det}(\mathbf{Id} - \mathcal{K}_D^\phi) \det J^\phi[D](x_1, \dots, x_n)$  is the Janossy density of  $\Phi(\mathbf{X}^D)$  with respect to  $\mu_\phi$ .  $\square$

*Integration by parts and closability.* We close this section with the statement and proof of the integration by parts formula for determinantal processes which is based on the closed gradient and divergence operators, cf. Theorem 3.9 below. The integration by parts formula is proved on the set of test functionals  $\mathcal{S}_D$  introduced in Definition 3.1, extending and making more precise the argument and proof of Theorem 10 page 289 of [4].



**(H3)** : The function  $(x_1, \dots, x_n) \mapsto \det J[D](x_1, \dots, x_n)$  is continuously differentiable on  $D^n$ .

Assuming that **(H1)** and **(H3)** hold, the potential energy is defined by

$$U[D](\mathbf{x}) := -\log \det J[D](\mathbf{x}).$$

We insist that since  $\det J[D](\mathbf{x}) > 0$  for  $\mathbb{P}_D$ -a.e.  $\mathbf{x} \in N_f^D$ ,  $U[D]$  is well defined for  $\mathbb{P}_D$ -a.e.  $\mathbf{x} \in N_f^D$ . For any  $v \in C^\infty(D, \mathbb{R}^d)$  and  $\mathbf{x} = \{x_1, \dots, x_{\mathbf{x}(D)}\} \in N_f^D$ , we set

$$\begin{aligned} \nabla_v^{N_{if}} U[D](\mathbf{x}) &:= -\sum_{k=1}^{\infty} \mathbf{1}_{\{\mathbf{x}(D)=k\}} \sum_{i=1}^k \frac{\nabla_{x_i} \det J[D](x_1, \dots, x_k)}{\det J[D](x_1, \dots, x_k)} \cdot v(x_i) \\ &= \sum_{k=1}^{\infty} \mathbf{1}_{\{\mathbf{x}(D)=k\}} \sum_{i=1}^k U_{i,k}(x_1, \dots, x_k) \cdot v(x_i). \end{aligned} \quad (3.5)$$

Under Conditions **(H1)** and **(H2)** we define the vector field  $\beta^\mu(x) := \nabla \rho(x)/\rho(x)$ , as well as

$$B_v^\mu(\mathbf{x}) := \sum_{k=1}^{\mathbf{x}(D)} (-\beta^\mu(x_k) \cdot v(x_k) + \operatorname{div} v(x_k)),$$

where  $\operatorname{div}$  denotes the adjoint of the gradient  $\nabla$  on  $D$ .

**Lemma 3.7.** *Assume **(H1)**, **(H2)** and **(H3)**. Then, for any  $F, G \in \mathcal{S}_D$  and  $v \in C^\infty(D, \mathbb{R}^d)$ , we have*

$$\mathbb{E}[G(\mathbf{X}^D) \nabla_v^{N_{if}} F(\mathbf{X}^D)] = \mathbb{E}[F(\mathbf{X}^D) \nabla_v^{N_{if}*} G(\mathbf{X}^D)], \quad (3.6)$$

where

$$\nabla_v^{N_{if}*} G(\mathbf{x}) := -\nabla_v^{N_{if}} G(\mathbf{x}) + G(\mathbf{x}) \left( -B_v^\mu(\mathbf{x}) + \nabla_v^{N_{if}} U[D](\mathbf{x}) \right), \quad \mathbf{x} \in N_f^D.$$

*Proof.* For  $v \in C^\infty(D, \mathbb{R}^d)$ , consider the flow  $\phi_t^v : D \rightarrow D$ ,  $t \in \mathbb{R}$ , where for a fixed  $x \in D$ , the curve  $t \mapsto \phi_t^v(x)$  is defined as the solution to the Cauchy problem

$$\frac{d}{dt} \phi_t^v(x) = v(\phi_t^v(x)), \quad \phi_0^v(x) = x.$$

We define the mapping  $\Phi_t^v : N_f^D \rightarrow N_f^D$  by  $\Phi_t^v(\mathbf{x}) := \{\phi_t^v(x) : x \in \mathbf{x}\}$ . Following [1], for a function  $R : N_f^D \rightarrow \mathbb{R}$ , we define the gradient  $\nabla_v^{N_{if}} R(\mathbf{x})$  as the directional derivative along  $v$  i.e. for  $\mathbf{x} \in N_f^D$ ,

$$\nabla_v^{N_{if}} R(\mathbf{x}) := \left. \frac{d}{dt} R(\Phi_t^v(\mathbf{x})) \right|_{t=0},$$

provided that the derivative exists. It is easy to check that formulas (3.3) and (3.5) are consistent with this definition. Note that by (3.4), the image measure  $\mu_{\phi_t^v}$  is absolutely continuous with respect to  $\mu$  on  $D$ , with Radon-Nikodym derivative

$$\frac{\rho(\phi_{-t}^v(x))}{\rho(x)} \operatorname{Jac}(\phi_{-t}^v)(x).$$

Note also that

$$\operatorname{Jac}(\phi_t^v)(x) = \exp \left( -\int_0^t \operatorname{div} v(\phi_z^v(x)) dz \right), \quad (3.7)$$

and therefore

$$\begin{aligned} \frac{d}{dt} \left( \frac{\rho(\phi_{-t}^v(x))}{\rho(x)} \text{Jac}(\phi_{-t}^v)(x) \right) &= -\exp\left(-\int_0^t \text{div } v(\phi_z^v(x)) dz\right) \\ &\quad \times \left[ \frac{\nabla \rho(\phi_{-t}^v(x))}{\rho(x)} \cdot v(\phi_{-t}^v(x)) + \frac{\rho(\phi_{-t}^v(x))}{\rho(x)} \text{div } v(\phi_{-t}^v(x)) \right]. \end{aligned}$$

Using Proposition 3.6, exchanging  $d/dt$  with  $\mathbb{E}$  (this exchange will be justified later on after (3.12)) and setting  $\text{Jac}^{\phi_t^v} := \text{Jac}(\phi_t^v)$ , we have

$$\mathbb{E} \left[ G(\mathbf{X}^D) \frac{d}{dt} F(\Phi_t^v(\mathbf{X}^D)) \right] \tag{3.8}$$

$$\begin{aligned} &= \mathbb{E} \left[ \left( \frac{d}{dt} G(\Phi_{-t}^v(\mathbf{X}^D)) \right) F(\mathbf{X}^D) \left( \prod_{k=1}^{\mathbf{X}(D)} \frac{\rho(\phi_{-t}^v(X_k))}{\rho(X_k)} \text{Jac}^{\phi_{-t}^v}(X_k) \right) \right. \\ &\quad \left. \times \frac{\det J[D](\phi_{-t}^v(X_1), \dots, \phi_{-t}^v(X_{\mathbf{X}(D)}))}{\det J[D](X_1, \dots, X_{\mathbf{X}(D)})} \right] \end{aligned}$$

$$+ \mathbb{E} \left[ \left( \frac{d}{dt} \prod_{k=1}^{\mathbf{X}(D)} \frac{\rho(\phi_{-t}^v(X_k))}{\rho(X_k)} \text{Jac}^{\phi_{-t}^v}(X_k) \right) F(\mathbf{X}^D) G(\Phi_{-t}^v(\mathbf{X}^D)) \right] \tag{3.9}$$

$$\times \frac{\det J[D](\phi_{-t}^v(X_1), \dots, \phi_{-t}^v(X_{\mathbf{X}(D)}))}{\det J[D](X_1, \dots, X_{\mathbf{X}(D)})} \tag{3.10}$$

$$\begin{aligned} &+ \mathbb{E} \left[ \left( \frac{d}{dt} \frac{\det J[D](\phi_{-t}^v(X_1), \dots, \phi_{-t}^v(X_{\mathbf{X}(D)}))}{\det J[D](X_1, \dots, X_{\mathbf{X}(D)})} \right) F(\mathbf{X}^D) G(\Phi_{-t}^v(\mathbf{X}^D)) \right. \\ &\quad \left. \times \prod_{k=1}^{\mathbf{X}(D)} \frac{\rho(\phi_{-t}^v(X_k))}{\rho(X_k)} \text{Jac}^{\phi_{-t}^v}(X_k) \right], \end{aligned} \tag{3.11}$$

for any  $t \in \mathbb{R}$  and  $F, G \in \mathcal{S}_D$ . The claimed integration by parts formula follows by evaluating the above relation at  $t = 0$ . Specifically, we use the relation

$$\begin{aligned} &\frac{d}{dt} \frac{\det J[D](\phi_{-t}^v(X_1), \dots, \phi_{-t}^v(X_{\mathbf{X}(D)}))}{\det J[D](X_1, \dots, X_{\mathbf{X}(D)})} \\ &= - \sum_{i=1}^{\mathbf{X}(D)} \frac{\nabla_{X_i} \det J[D](\phi_{-t}^v(X_1), \dots, \phi_{-t}^v(X_{\mathbf{X}(D)}))}{\det J[D](X_1, \dots, X_{\mathbf{X}(D)})} \cdot v(\phi_{-t}^v(X_i)) \end{aligned} \tag{3.12}$$

to evaluate (3.11). Using the definition of  $\mathcal{S}_D$ , one checks that for  $\mathbb{P}_D$ -a.e.  $\mathbf{x} \in N_f^D$ ,

$$t \mapsto G(\mathbf{x}) \frac{d}{dt} F(\Phi_t^v(\mathbf{x})),$$

is uniformly bounded by a positive constant in an neighborhood of zero. By the assumptions **(H2)** and **(H3)** and the form (3.1) of the functionals in  $\mathcal{S}_D$ , one may easily check that (3.9), (3.10) and (3.11) can be uniformly bounded in an

neighborhood of zero by  $\mathbb{P}_D$ -integrable functions. This justifies the exchange of  $d/dt$  and  $\mathbb{E}$  claimed previously. We check this fact only for (3.11). Take

$$F(\mathbf{x}) = f_0 \mathbf{1}_{\{\mathbf{x}(D)=0\}} + \sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} f_k(x_1, \dots, x_k), \quad \mathbf{x} = \{x_1, \dots, x_{\mathbf{x}(D)}\} \in N_f^D,$$

of the form (3.1). By (3.12) we easily see that, up to a positive constant, the modulus of the random variable

$$\begin{aligned} & \left( \frac{d}{dt} \frac{\det J[D](\phi_{-t}^v(X_1), \dots, \phi_{-t}^v(X_{\mathbf{x}(D)}))}{\det J[D](X_1, \dots, X_{\mathbf{x}(D)})} \right) F(\mathbf{X}^D) G(\Phi_{-t}^v(\mathbf{X}^D)) \\ & \qquad \qquad \qquad \times \prod_{k=1}^{\mathbf{x}(D)} \frac{\rho(\phi_{-t}^v(X_k))}{\rho(X_k)} \text{Jac}^{\phi_{-t}^v}(X_k) \end{aligned}$$

is bounded above by

$$\begin{aligned} & \sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} \left( \prod_{i=1}^k \frac{\rho(\phi_{-t}^v(X_i))}{\rho(X_i)} \text{Jac}^{\phi_{-t}^v}(X_i) \right) \\ & \qquad \times \sum_{i=1}^k \left| \frac{\nabla_{X_i} \det J[D](\phi_{-t}^v(X_1), \dots, \phi_{-t}^v(X_k))}{\det J[D](X_1, \dots, X_k)} \cdot v(\phi_{-t}^v(X_i)) \right|. \end{aligned} \quad (3.13)$$

By assumptions **(H2)**, **(H3)** and equation (3.7), it follows that, up to a positive constant, (3.13) is bounded above by

$$\sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} \frac{\rho(X_1)^{-1} \cdots \rho(X_k)^{-1}}{\det J[D](X_1, \dots, X_k)},$$

for any  $t$  in a neighborhood of zero. We conclude the proof noticing that the mean of this r.v. is finite by definition of the Janossy densities, and since  $\det J[D](\mathbf{x}) > 0$ , for  $\mathbb{P}_D$ -a.e.  $\mathbf{x} \in N_f^D$ .  $\square$

*Remark 3.8.* We remark that there is a sign change in (3.6), as compared to the results of [4], which is justified by the corrected formula for (3.4).

Next, we extend the integration by parts formula by closability to a larger class of functionals. For  $v \in \mathcal{C}^\infty(D, \mathbb{R}^d)$ , we consider the closability of the linear operators  $\nabla_v^{N_{lf}} : \mathcal{S}_D \rightarrow L_D^2$  and  $\nabla_v^{N_{lf}^*} : \mathcal{S}_D \rightarrow L_D^2$ . In the following, we denote by  $\bar{A}$  the minimal closed extension of a closable linear operator  $A$ , and by  $\text{Dom}(\bar{A})$  the domain of  $\bar{A}$ . We refer the reader to [7] for all these notions of functional analysis.

In Theorem 3.9 below we assume, in addition to **(H1)**, **(H2)** and **(H3)**, the following condition.

**(H4):** For any  $n \geq 1$ ,  $1 \leq i, j \leq n$ , and  $1 \leq h, k \leq d$ , we have

$$\int_{D^n} \left| \frac{\partial_{x_i^{(h)}} \det J[D](x_1, \dots, x_n) \partial_{x_j^{(k)}} \det J[D](x_1, \dots, x_n)}{\det J[D](x_1, \dots, x_n)} \right| \mathbf{1}_{\{\det J[D](x_1, \dots, x_n) > 0\}} \mu(dx_1) \cdots \mu(dx_n) < \infty.$$

**Theorem 3.9.** Assume **(H1)** – **(H4)**.

(i) For any  $v \in C^\infty(D, \mathbb{R}^d)$ , the linear operators  $\nabla_v^{N_{lf}}$  and  $\nabla_v^{N_{lf}^*}$  are well-defined, i.e.

$$\nabla_v^{N_{lf}}(\mathcal{S}_D) \subset L_D^2 \quad \text{and} \quad \nabla_v^{N_{lf}^*}(\mathcal{S}_D) \subset L_D^2,$$

and closable.

(ii) For any  $v \in C^\infty(D, \mathbb{R}^d)$ , we have

$$\mathbb{E} \left[ G(\mathbf{X}^D) \overline{\nabla_v^{N_{lf}} F(\mathbf{X}^D)} \right] = \mathbb{E} \left[ F(\mathbf{X}^D) \overline{\nabla_v^{N_{lf}^*} G(\mathbf{X}^D)} \right]$$

for all  $F \in \text{Dom}(\overline{\nabla_v^{N_{lf}}})$ ,  $G \in \text{Dom}(\overline{\nabla_v^{N_{lf}^*}})$ .

*Remark 3.10.* Note that under **(H1)**, **(H2)** and **(H3)**, condition **(H4)** is satisfied if, for any  $n \geq 1$ , the function

$$(x_1, \dots, x_n) \mapsto \det J[D](x_1, \dots, x_n),$$

is strictly positive on the compact  $D^n$ .

*Proof of Theorem 3.9.* (i) Let  $v \in C_c^\infty(D, \mathbb{R}^d)$  and  $F \in \mathcal{S}_D$ . For ease of notation, throughout this proof we write  $\nabla_v$  in place of  $\nabla_v^{N_{lf}}$  and  $\nabla_v^*$  in place of  $\nabla_v^{N_{lf}^*}$ . We clearly have  $|\nabla_v F(\mathbf{x})| \leq C$  for some constant  $C > 0$ ,  $\mathbb{P}_D$ -a.e., and therefore  $\nabla_v(\mathcal{S}_D) \subset L_D^2$ . The claim  $\nabla_v^*(\mathcal{S}_D) \subset L_D^2$  follows if we check that

$$\|G(\mathbf{x}) \nabla_v U[D](\mathbf{x})\|_{L_D^2} < \infty \quad \text{and} \quad \|G(\mathbf{x}) B_v^\mu(\mathbf{x})\|_{L_D^2} < \infty$$

for any  $G \in \mathcal{S}_D$ . The latter relation easily follows noticing that  $|G(\mathbf{x}) B_v^\mu(\mathbf{x})| \leq C$  for some constant  $C > 0$ ,  $\mathbb{P}_D$ -a.e.. Taking  $G$  of the form (3.1), by (3.5), for some positive constant  $C > 0$ , we have

$$\begin{aligned} \|G \nabla_v U[D]\|_{L_D^2}^2 &\leq C \text{Det}(\mathbf{Id} - \mathcal{K}_D) \sum_{k=1}^m \frac{1}{k!} \sum_{1 \leq i, j \leq k} \int_{D^k} \mathbf{1}_{\{\det J[D](x_1, \dots, x_k) > 0\}} \\ &\frac{\nabla_{x_i} \det J[D](x_1, \dots, x_k) \cdot v(x_i) \nabla_{x_j} \det J[D](x_1, \dots, x_k) \cdot v(x_j)}{\det J[D](x_1, \dots, x_k)} \mu(dx_1) \cdots \mu(dx_k) \\ &< \infty, \end{aligned}$$

where the latter integral is finite by **(H4)**.

To conclude, we only need to show that  $\nabla_v$  is closable (the closability of  $\nabla_v^*$  can be proved similarly). Let  $(F_n)_{n \geq 1}$  be a sequence in  $\mathcal{S}_D$  converging to 0 in  $L_D^2$  and such that  $\nabla_v F_n$  converges to  $V$  in  $L_D^2$  as  $n$  goes to infinity. We need to show that  $V = 0$   $\mathbb{P}_D$ -a.e.. We have

$$\begin{aligned} \left| \langle G, V \rangle_{L_D^2} \right| &= \lim_{n \rightarrow \infty} |\mathbb{E}[G(\mathbf{X}^D) \nabla_v F_n(\mathbf{X}^D)]| = \lim_{n \rightarrow \infty} |\mathbb{E}[F_n(\mathbf{X}^D) \nabla_v^* G(\mathbf{X}^D)]| \quad (3.14) \\ &\leq \|\nabla_v^* G\|_{L_D^2} \lim_{n \rightarrow \infty} \|F_n\|_{L_D^2} = 0, \quad G \in \mathcal{S}_D. \end{aligned}$$

Here, the second inequality in (3.14) follows by the integration by parts formula (3.6). The conclusion follows noticing that  $\langle G, V \rangle_{L_D^2} = 0$  for all  $G \in \mathcal{S}_D$  implies  $V = 0$   $\mathbb{P}_D$ -a.e. due to the density of  $\mathcal{S}_D$  in  $L_D^2$ .

(ii) By (i) both operators  $\nabla_v$  and  $\nabla_v^*$  are closable. Take  $F \in \text{Dom}(\overline{\nabla_v})$ ,  $G \in \text{Dom}(\overline{\nabla_v^*})$  and let  $(F_n)_{n \geq 1}$ ,  $(G_n)_{n \geq 1}$  be sequences in  $\mathcal{S}_D$  such that  $F_n$  converges to

$F, G_n$  converges to  $G$ ,  $\nabla_v F_n$  converges to  $\overline{\nabla}_v F$  and  $\nabla_v^* G_n$  converges to  $\overline{\nabla}_v^* G$  in  $L^2_D$  as  $n$  goes to infinity. By Lemma 3.7 the integration by parts formula applies to r.v.'s in  $\mathcal{S}_D$ , therefore we have  $\mathbb{E}[G_n(\mathbf{X}^D)\nabla_v F_n(\mathbf{X}^D)] = \mathbb{E}[F_n(\mathbf{X}^D)\nabla_v^* G_n(\mathbf{X}^D)]$  for all  $n \geq 1$ . The claim follows if we prove that

$$\lim_{n \rightarrow \infty} \mathbb{E}[G_n(\mathbf{X}^D)\nabla_v F_n(\mathbf{X}^D)] = \mathbb{E}[G(\mathbf{X}^D)\overline{\nabla}_v F(\mathbf{X}^D)]$$

and

$$\lim_{n \rightarrow \infty} \mathbb{E}[F_n(\mathbf{X}^D)\nabla_v^* G_n(\mathbf{X}^D)] = \mathbb{E}[F(\mathbf{X}^D)\overline{\nabla}_v^* G(\mathbf{X}^D)].$$

We only show the first limit above; the second limit being proved similarly. We have

$$\begin{aligned} & |\mathbb{E}[G_n(\mathbf{X}^D)\nabla_v F_n(\mathbf{X}^D)] - \mathbb{E}[G(\mathbf{X}^D)\overline{\nabla}_v F(\mathbf{X}^D)]| \\ & \leq \|G_n\|_{L^2_D} \|\nabla_v F_n - \overline{\nabla}_v F\|_{L^2_D} + \|G_n - G\|_{L^2_D} \|\overline{\nabla}_v F\|_{L^2_D}, \end{aligned}$$

which tends to 0 as  $n$  goes to infinity. □

#### 4. Dirichlet Forms Corresponding to Determinantal Processes on $D \subset S$

In this section we construct the Dirichlet form associated to a determinantal process, cf. Theorem 4.2 below. We start by recalling some definitions related to bilinear forms (see [15] for details). Let  $H$  be a Hilbert space with inner product  $\langle \cdot, \cdot \rangle$  and  $\mathcal{A} : \text{Dom}(\mathcal{A}) \times \text{Dom}(\mathcal{A}) \rightarrow \mathbb{R}$  a bilinear form defined on a dense subspace  $\text{Dom}(\mathcal{A})$  of  $H$ , the domain of  $\mathcal{A}$ . The form  $\mathcal{A}$  is said to be symmetric if  $\mathcal{A}(F, G) = \mathcal{A}(G, F)$ , for any  $F, G \in \text{Dom}(\mathcal{A})$ , and non-negative definite if  $\mathcal{A}(F, F) \geq 0$ , for any  $F \in \text{Dom}(\mathcal{A})$ . Let  $\mathcal{A}$  be symmetric and non-negative definite,  $\mathcal{A}$  is said to be closed if  $\text{Dom}(\mathcal{A})$  equipped with the norm

$$\|F\|_{\mathcal{A}} := \sqrt{\mathcal{A}(F, F) + \langle F, F \rangle}, \quad F \in \text{Dom}(\mathcal{A}),$$

is a Hilbert space. A symmetric and non-negative definite bilinear form  $\mathcal{A}$  is said to be closable if, for any sequence  $(F_n)_{n \geq 1} \subset \text{Dom}(\mathcal{A})$  such that  $F_n$  goes to 0 in  $H$  and  $(F_n)_{n \geq 1}$  is Cauchy *w.r.t.*  $\|\cdot\|_{\mathcal{A}}$  it holds that  $\mathcal{A}(F_n, F_n)$  converges to 0 in  $\mathbb{R}$  as  $n$  goes to infinity. Let  $\mathcal{A}$  be closable and denote by  $\text{Dom}(\overline{\mathcal{A}})$  the completion of  $\text{Dom}(\mathcal{A})$  *w.r.t.* the norm  $\|\cdot\|_{\mathcal{A}}$ . It turns out that  $\mathcal{A}$  is uniquely extended to  $\text{Dom}(\overline{\mathcal{A}})$  by the closed, symmetric and non-negative definite bilinear form

$$\overline{\mathcal{A}}(F, G) = \lim_{n \rightarrow \infty} \mathcal{A}(F_n, G_n), \quad (F, G) \in \text{Dom}(\overline{\mathcal{A}}) \times \text{Dom}(\overline{\mathcal{A}}),$$

where  $\{(F_n, G_n)\}_{n \geq 1}$  is any sequence in  $\text{Dom}(\mathcal{A}) \times \text{Dom}(\mathcal{A})$  such that  $(F_n, G_n)$  converges to  $(F, G) \in \text{Dom}(\overline{\mathcal{A}}) \times \text{Dom}(\overline{\mathcal{A}})$  *w.r.t.* the norm  $\|\cdot\|_{\overline{\mathcal{A}}} + \|\cdot\|_{\overline{\mathcal{A}}}$ . Suppose  $H = L^2(B, \beta)$  where  $(B, \mathcal{F}, \beta)$  is a  $\sigma$ -finite measure space. A symmetric, non-negative definite and closed bilinear form  $\mathcal{A}$  is said to be a Dirichlet form if

$$\mathcal{A}(\min\{F^+, 1\}, \min\{F^+, 1\}) \leq \mathcal{A}(F, F), \quad F \in \text{Dom}(\mathcal{A}),$$

where  $F^+$  denotes the positive part of  $F$ . Suppose that  $B$  is a Hausdorff topological space,  $\mathcal{F} = \mathcal{B}(B)$  the corresponding Borel  $\sigma$ -algebra, and let  $\mathcal{A}$  be a Dirichlet form. Throughout this paper, we shall use the following notions. The generator  $\mathcal{H}^{\text{gen}}$  of

the Dirichlet form is the unique symmetric non-negative definite operator on  $H$  such that

$$\mathcal{A}(F, G) = \langle (-\mathcal{H}^{\text{gen}} F), G \rangle,$$

for  $F, G \in \text{Dom}(-\mathcal{H}^{\text{gen}}) \subset \text{Dom}(\mathcal{A})$ . The symmetric semi-group of  $\mathcal{A}$  is the linear operator on  $H$  defined by  $T_t F := \exp(-t\mathcal{H}^{\text{gen}})F$ ,  $t > 0$ .

After these general considerations, we move to the situation at hand. Assume **(H1)**, **(H2)** and **(H3)**. We recall that  $D$  is a fixed compact set of  $S$  which is in turn included in  $\mathbb{R}^d$ . We denote by  $\check{N}_f^D$  the set of  $\mathbb{N}$ -valued Radon measures on  $D$ . We equip  $\check{N}_f^D$  with the vague topology and the corresponding Borel  $\sigma$ -algebra  $\mathcal{B}(\check{N}_f^D)$ . Note that  $N_f^D$  is contained in  $\check{N}_f^D$ . In the following, we consider the subspace  $\tilde{\mathcal{S}}_D$  of  $\mathcal{S}_D$  made of cylindrical functions.

**Definition 4.1.** A function  $F : N_f^D \rightarrow \mathbb{R}$  is said to be in  $\tilde{\mathcal{S}}_D$  if it is of the form

$$F(\mathbf{x}) = f \left( \sum_{k=1}^{\mathbf{x}(D)} \varphi_1(x_k), \dots, \sum_{k=1}^{\mathbf{x}(D)} \varphi_m(x_k) \right) \mathbf{1}_{\{\mathbf{x}(D) \leq n\}}, \quad \mathbf{x} = \{x_1, \dots, x_{\mathbf{x}(D)}\} \in N_f^D,$$

for some integers  $m, n \geq 1$ ,  $\varphi_1, \dots, \varphi_m \in C^\infty(D)$ ,  $f \in C_b^\infty(\mathbb{R}^m)$ .

Note that, as already mentioned,  $\tilde{\mathcal{S}}_D$  is dense in  $L_D^2$ . We consider the bilinear map  $\mathcal{E}_D$  defined on  $\tilde{\mathcal{S}}_D \times \tilde{\mathcal{S}}_D$  by

$$\mathcal{E}_D(F, G) := \mathbb{E} \left[ \sum_{i=1}^{\mathbf{x}(D)} \nabla_{X_i}^{N_{if}} F(\mathbf{X}^D) \cdot \nabla_{X_i}^{N_{if}} G(\mathbf{X}^D) \right].$$

For  $F \in \mathcal{S}_D$  of the form (3.1), i.e.

$$F(\mathbf{x}) = f_0 \mathbf{1}_{\{\mathbf{x}(D)=0\}} + \sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} f_k(x_1, \dots, x_k), \quad \mathbf{x} = \{x_1, \dots, x_{\mathbf{x}(D)}\} \in N_f^D$$

we also define the Laplace operator  $\mathcal{H}_D$  by

$$\begin{aligned} \mathcal{H}_D F(\mathbf{x}) = & \sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} \sum_{i=1}^k (-\beta^\mu(x_i) \cdot \nabla_{x_i} f_k(x_1, \dots, x_k) \\ & - \Delta_{x_i} f_k(x_1, \dots, x_k) + U_{i,k}(x_1, \dots, x_k) \cdot \nabla_{x_i} f_k(x_1, \dots, x_k)), \end{aligned}$$

where  $\Delta = -\text{div} \nabla$  denotes the Laplacian. The next theorem provides the Dirichlet form associated to a determinantal process.

**Theorem 4.2.** *Under conditions **(H1)** – **(H4)** we have:*

(i) *The Laplace operator  $\mathcal{H}_D : \tilde{\mathcal{S}}_D \rightarrow L_D^2$  is linear, symmetric, non-negative definite and well-defined, i.e.  $\mathcal{H}_D(\tilde{\mathcal{S}}_D) \subset L_D^2$ . In particular the operator square root  $\mathcal{H}_D^{1/2}$  of  $\mathcal{H}_D$  exists.*

(ii) *The bilinear form  $\mathcal{E}_D : \tilde{\mathcal{S}}_D \times \tilde{\mathcal{S}}_D \rightarrow \mathbb{R}$  is symmetric, non-negative definite and well-defined, i.e.  $\mathcal{E}_D(\tilde{\mathcal{S}}_D \times \tilde{\mathcal{S}}_D) \subset \mathbb{R}$ .*

(iii)  *$\mathcal{H}_D^{1/2}$  and  $\mathcal{E}_D$  are closable and the following relation holds:*

$$\overline{\mathcal{E}_D}(F, G) = \mathbb{E} \left[ \overline{\mathcal{H}_D^{1/2}} F(\mathbf{X}^D) \overline{\mathcal{H}_D^{1/2}} G(\mathbf{X}^D) \right], \quad \forall F, G \in \text{Dom}(\overline{\mathcal{H}_D^{1/2}}). \quad (4.1)$$

(iv) The bilinear form  $(\overline{\mathcal{E}_D}, \text{Dom}(\overline{\mathcal{H}_D^{1/2}}))$  is a Dirichlet form.

The proof of the theorem is based on the following lemma.

**Lemma 4.3.** Under conditions **(H1)** – **(H4)**, for any  $F, G \in \tilde{\mathcal{S}}_D$ , we have

$$\mathbb{E} \left[ \sum_{i=1}^{\mathbf{x}(D)} \nabla_{X_i}^{N_{if}} F(\mathbf{X}^D) \cdot \nabla_{X_i}^{N_{if}} G(\mathbf{X}^D) \right] = \mathbb{E}[G(\mathbf{X}^D) \mathcal{H}_D F(\mathbf{X}^D)] \tag{4.2}$$

$$= \mathbb{E}[\mathcal{H}_D^{1/2} F(\mathbf{X}^D) \mathcal{H}_D^{1/2} G(\mathbf{X}^D)]. \tag{4.3}$$

*Proof of Theorem 4.2.* (i) By Relation (4.2) in Lemma 4.3 we easily deduce that, for any  $F, G \in \tilde{\mathcal{S}}_D$  we have

$$\mathbb{E}[G(\mathbf{X}^D) \mathcal{H}_D F(\mathbf{X}^D)] = \mathbb{E}[F(\mathbf{X}^D) \mathcal{H}_D G(\mathbf{X}^D)] \quad \text{and} \quad \mathbb{E}[F(\mathbf{X}^D) \mathcal{H}_D F(\mathbf{X}^D)] \geq 0.$$

Therefore,  $\mathcal{H}_D$  is symmetric and non-negative definite. It remains to check that, under the foregoing assumptions,  $\mathcal{H}_D$  is well-defined. Let  $F \in \tilde{\mathcal{S}}_D$  be of the form

$$\begin{aligned} F(\mathbf{x}) &= \sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} f \left( \sum_{i=1}^k \varphi_1(x_i), \dots, \sum_{i=1}^k \varphi_m(x_i) \right) \\ &= \sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} f_k(x_1, \dots, x_k) \end{aligned}$$

for some integers  $m, n \geq 1$ ,  $\varphi_1, \dots, \varphi_m \in \mathcal{C}^\infty(D)$ ,  $f \in \mathcal{C}_b^\infty(\mathbb{R}^m)$ . For the well-definiteness of  $\mathcal{H}_D$  we only need to check that, for  $\mathbf{x} = \{x_1, \dots, x_{\mathbf{x}(D)}\} \in \mathbb{N}_f^D$ ,

$$F_1(\mathbf{x}) := \mathbf{1}_{\{\mathbf{x}(D)=k\}} (\beta^\mu(x_i))^{(j)}, \quad \text{and} \quad F_2(\mathbf{x}) := \mathbf{1}_{\{\mathbf{x}(D)=k\}} (U_{i,k}(x_1, \dots, x_k))^{(j)},$$

are in  $L_D^2$  for all  $k \in \{1, \dots, n\}$ ,  $i \in \{1, \dots, k\}$  and  $j \in \{1, \dots, d\}$ . One may easily check that  $F_1 \in L_D^2$  due to **(H2)** and  $F_2 \in L_D^2$  due to **(H4)**.

(ii) The symmetry and non-negative definiteness of  $\mathcal{E}_D$  follow from Lemma 4.3. It remains to check that, under the foregoing assumptions,  $\mathcal{E}_D$  is well-defined. By Step (i), for any  $F \in \tilde{\mathcal{S}}_D$ , we have  $\mathcal{H}_D F \in L_D^2$ . We conclude the proof by noting that, by Lemma 4.3, for any  $F, G \in \tilde{\mathcal{S}}_D$  and some positive constant  $c > 0$ , we have

$$|\mathcal{E}_D(F, G)| = |\mathbb{E}[G(\mathbf{X}^D) \mathcal{H}_D F(\mathbf{X}^D)]| \leq c \|\mathcal{H}_D F(\mathbf{X}^D)\|_{L_D^2} < \infty.$$

(iii) We first show that  $\mathcal{E}_D$  is closable. By Lemma 3.4 page 29 in [15], we have to check that if  $(F_n)_{n \geq 1} \subset \tilde{\mathcal{S}}_D$  is such that  $F_n$  converges to 0 in  $L_D^2$ , then  $\mathcal{E}_D(G, F_n)$  converges to 0, for any  $G \in \tilde{\mathcal{S}}_D$ . This easily follows by Lemma 4.3, the Cauchy-Schwarz inequality and the fact that  $\mathcal{H}_D G$  is square integrable (see Step (i)). The closability of  $\mathcal{H}_D^{1/2}$  follows by the closability of  $\mathcal{E}_D$ , relation (4.3) in Lemma 4.3 and Remark 3.2 (i) page 29 in [15]. Finally, we prove relation (4.1). Take  $F, G \in \text{Dom}(\overline{\mathcal{H}_D^{1/2}})$  and let  $(F_n)_{n \geq 1}, (G_n)_{n \geq 1}$  be sequences in  $\tilde{\mathcal{S}}_D$  such that  $F_n$  converges to  $F$ ,  $G_n$  converges to  $G$ ,  $\mathcal{H}_D^{1/2} F_n$  converges to  $\overline{\mathcal{H}_D^{1/2}} F$ , and  $\mathcal{H}_D^{1/2} G_n$  converges to  $\overline{\mathcal{H}_D^{1/2}} G$  in  $L_D^2$  as  $n$  goes to infinity. By Lemma 4.3 we have

$$\mathcal{E}_D(F_n, G_n) = \mathbb{E}[\mathcal{H}_D^{1/2} F_n(\mathbf{X}^D) \mathcal{H}_D^{1/2} G_n(\mathbf{X}^D)], \quad \text{for all } n \geq 1.$$

The claim follows if we prove that

$$\lim_{n \rightarrow \infty} \mathbb{E}[\mathcal{H}_D^{1/2} F_n(\mathbf{X}^D) \mathcal{H}_D^{1/2} G_n(\mathbf{X}^D)] = \mathbb{E}[\overline{\mathcal{H}_D^{1/2}} F(\mathbf{X}^D) \overline{\mathcal{H}_D^{1/2}} G(\mathbf{X}^D)].$$

Indeed we have:

$$\begin{aligned} & |\mathbb{E}[\mathcal{H}_D^{1/2} G_n(\mathbf{X}^D) \mathcal{H}_D^{1/2} F_n(\mathbf{X}^D)] - \mathbb{E}[\overline{\mathcal{H}_D^{1/2}} G(\mathbf{X}^D) \overline{\mathcal{H}_D^{1/2}} F(\mathbf{X}^D)]| \\ & \leq \|\mathcal{H}_D^{1/2} G_n\|_{L_b^2} \|\mathcal{H}_D^{1/2} F_n - \overline{\mathcal{H}_D^{1/2}} F\|_{L_b^2} \\ & \quad + \|\mathcal{H}_D^{1/2} G_n - \overline{\mathcal{H}_D^{1/2}} G\|_{L_b^2} \|\overline{\mathcal{H}_D^{1/2}} F\|_{L_b^2} \rightarrow 0, \quad \text{as } n \rightarrow \infty. \end{aligned}$$

(iv) The bilinear form  $(\overline{\mathcal{E}}_D, \text{Dom}(\overline{\mathcal{H}_D^{1/2}}))$  defined by (4.1) is clearly symmetric, non-negative definite, and closed. We conclude the proof by applying Proposition 4.10 page 35 in [15]. First, note that  $\tilde{\mathcal{S}}_D$  is dense in  $\text{Dom}(\overline{\mathcal{H}_D^{1/2}})$  (w.r.t. the norm  $\|\cdot\|_{\overline{\mathcal{E}}_D}$ ). By Exercise 2.7 page 47 in [15], for any  $\varepsilon > 0$  there exists an infinitely differentiable function  $\varphi_\varepsilon : \mathbb{R} \rightarrow [-\varepsilon, 1 + \varepsilon]$  (which shall not be confused with the functions  $\varphi_1, \dots, \varphi_m$  involved in Definition 4.1) such that  $\varphi_\varepsilon(t) = t$  for any  $t \in [0, 1]$ ,  $0 \leq \varphi_\varepsilon(t) - \varphi_\varepsilon(s) \leq t - s$  for all  $t, s \in \mathbb{R}$ ,  $t \geq s$ ,  $\varphi_\varepsilon(t) = 1 + \varepsilon$  for  $t \in [1 + 2\varepsilon, \infty)$  and  $\varphi_\varepsilon(t) = -\varepsilon$  for  $t \in (-\infty, -2\varepsilon]$ . Note that  $|\varphi'_\varepsilon(t)|^2 \leq 1$  for any  $\varepsilon > 0$ ,  $t \in \mathbb{R}$  and  $\varphi_\varepsilon$  is in  $\mathcal{C}_b^\infty$ , for any  $\varepsilon > 0$ . Consider the function

$$F(\mathbf{x}) = f\left(\sum_{k=1}^{\mathbf{x}(D)} \varphi_1(x_k), \dots, \sum_{k=1}^{\mathbf{x}(D)} \varphi_m(x_k)\right) \mathbf{1}_{\{\mathbf{x}(D) \leq n\}}, \quad \mathbf{x} = \{x_1, \dots, x_{\mathbf{x}(D)}\} \in \mathbb{N}_f^D,$$

for some integers  $m, n \geq 1$ ,  $\varphi_1, \dots, \varphi_m \in \mathcal{C}^\infty(D)$ ,  $f \in \mathcal{C}^\infty(\mathbb{R}^m)$ . Note that  $\varphi_\varepsilon \circ F \in \tilde{\mathcal{S}}_D$ . Indeed we have

$$\varphi_\varepsilon \circ F(\mathbf{x}) = \varphi_\varepsilon\left(f\left(\sum_{k=1}^{\mathbf{x}(D)} \varphi_1(x_k), \dots, \sum_{k=1}^{\mathbf{x}(D)} \varphi_m(x_k)\right)\right) \mathbf{1}_{\{\mathbf{x}(D) \leq n\}},$$

because  $\varphi_\varepsilon(0) = 0$ . Next, we have

$$\begin{aligned} \overline{\mathcal{E}}_D(\varphi_\varepsilon \circ F, \varphi_\varepsilon \circ F) &= \mathbb{E}\left[\sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} \sum_{i=1}^k \|\nabla_{X_i}^{N_{if}} \varphi_\varepsilon \circ F(\mathbf{X}^D)\|^2\right] \\ &= \mathbb{E}\left[\sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} \sum_{i=1}^k \left\| \sum_{j=1}^m \varphi'_\varepsilon \circ f\left(\sum_{l=1}^k \varphi_1(X_l), \dots, \sum_{l=1}^k \varphi_m(X_l)\right) \right. \right. \\ & \quad \left. \left. \times \partial_j f\left(\sum_{l=1}^k \varphi_1(X_l), \dots, \sum_{l=1}^k \varphi_m(X_l)\right) \nabla \varphi_j(X_i) \right\|^2\right] \\ &\leq \mathbb{E}\left[\sum_{k=1}^n \mathbf{1}_{\{\mathbf{x}(D)=k\}} \sum_{i=1}^k \left\| \sum_{j=1}^m \partial_j f\left(\sum_{l=1}^k \varphi_1(X_l), \dots, \sum_{l=1}^k \varphi_m(X_l)\right) \nabla \varphi_j(X_i) \right\|^2\right] \\ &= \overline{\mathcal{E}}_D(F, F), \end{aligned} \tag{4.4}$$



where in (4.4) we used the fact that  $|\varphi'_\varepsilon(t)|^2 \leq 1$ ,  $t \in \mathbb{R}$ . By this inequality we have, for any  $F \in \tilde{\mathcal{S}}_D$ ,

$$\liminf_{\varepsilon \rightarrow 0} \overline{\mathcal{E}}_D(F \pm \varphi_\varepsilon \circ F, F \mp \varphi_\varepsilon \circ F) \geq 0$$

and the proof is completed (since, as required by Proposition 4.10 page 35 in [15], we checked condition (4.6) page 34 in [15]). Indeed, for any  $\varepsilon > 0$ , by the above inequality and Lemma 4.3, we have

$$\begin{aligned} & \overline{\mathcal{E}}_D(F \pm \varphi_\varepsilon \circ F, F \mp \varphi_\varepsilon \circ F) \\ &= \mathbb{E}[(F(\mathbf{X}^D) - \varphi_\varepsilon \circ F(\mathbf{X}^D))\mathcal{H}_D(F(\mathbf{X}^D) + \varphi_\varepsilon \circ F(\mathbf{X}^D))] \\ &\geq \mathbb{E}[F(\mathbf{X}^D)\mathcal{H}_D\varphi_\varepsilon \circ F(\mathbf{X}^D) - \varphi_\varepsilon \circ F(\mathbf{X}^D)\mathcal{H}_DF(\mathbf{X}^D)] = 0. \end{aligned}$$

□

*Proof of Lemma 4.3.* Let  $F, G \in \tilde{\mathcal{S}}_D$  be defined for  $\mathbf{x} = \{x_1, \dots, x_{\mathbf{x}(D)}\} \in \mathbb{N}_f^D$ , respectively, as

$$\begin{aligned} F(\mathbf{x}) &= f \left( \sum_{k=1}^{\mathbf{x}(D)} \varphi_1(x_k), \dots, \sum_{k=1}^{\mathbf{x}(D)} \varphi_{m_1}(x_k) \right) \mathbf{1}_{\{\mathbf{x}(D) \leq n_1\}}, \\ G(\mathbf{x}) &= g \left( \sum_{k=1}^{\mathbf{x}(D)} \gamma_1(x_k), \dots, \sum_{k=1}^{\mathbf{x}(D)} \gamma_{m_2}(x_k) \right) \mathbf{1}_{\{\mathbf{x}(D) \leq n_2\}}, \end{aligned}$$

for some integers  $m_1, m_2, n_1, n_2 \geq 1$ ,  $\varphi_1, \dots, \varphi_{m_1}, \gamma_1, \dots, \gamma_{m_2} \in \mathcal{C}^\infty(D)$ ,  $f \in \mathcal{C}_b^\infty(\mathbb{R}^{m_1})$ ,  $g \in \mathcal{C}_b^\infty(\mathbb{R}^{m_2})$ . Define

$$F_i(\mathbf{x}) = \partial_i f \left( \sum_{k=1}^{\mathbf{x}(D)} \varphi_1(x_k), \dots, \sum_{k=1}^{\mathbf{x}(D)} \varphi_{m_1}(x_k) \right) \mathbf{1}_{\{\mathbf{x}(D) \leq n_1\}},$$

and  $v_i(x) = \nabla \varphi_i(x)$ ,  $x \in D$ . By a direct computation we find

$$\mathcal{H}_DF(\mathbf{x}) = \sum_{i=1}^{m_1} \left( -\nabla_{v_i}^{N_{if}} F_i(\mathbf{x}) + (B_{v_i}^\mu(\mathbf{x}) + \nabla_{v_i}^{N_{if}} U[D](\mathbf{x})) F_i(\mathbf{x}) \right) = \sum_{i=1}^{m_1} \nabla_{v_i}^{N_{if}^*} F_i(\mathbf{x}).$$

So, by Lemma 3.7 and since  $\tilde{\mathcal{S}}_D \subset \mathcal{S}_D$ , using obvious notation we have

$$\begin{aligned} & \mathbb{E}[G(\mathbf{X}^D)\mathcal{H}_DF(\mathbf{X}^D)] \\ &= \sum_{i=1}^{m_1} \mathbb{E} \left[ G(\mathbf{X}^D) \nabla_{v_i}^{N_{if}^*} F_i(\mathbf{X}^D) \right] = \sum_{i=1}^{m_1} \mathbb{E} \left[ F_i(\mathbf{X}^D) \nabla_{v_i}^{N_{if}} G(\mathbf{X}^D) \right] \\ &= \mathbb{E} \left[ \sum_{l=1}^{\mathbf{X}(D)} \sum_{i=1}^{m_1} F_i(\mathbf{X}^D) \nabla \varphi_i(X_l) \cdot \sum_{j=1}^{m_2} G_j(\mathbf{X}^D) \nabla \gamma_j(X_l) \right] \\ &= \mathbb{E} \left[ \sum_{i=1}^{\mathbf{X}(D)} \nabla_{X_i}^{N_{if}} F(\mathbf{X}^D) \cdot \nabla_{X_i}^{N_{if}} G(\mathbf{X}^D) \right]. \end{aligned}$$

Finally, since  $\mathcal{H}_D$  is symmetric and non-negative definite the square root operator  $\mathcal{H}_D^{1/2}$  is well-defined. Relation (4.3) follows by the properties of  $\mathcal{H}_D^{1/2}$ .  $\square$

Let  $\mathcal{FC}_b^\infty(D)$  denote the set of functionals  $F : \mathbb{N}_f^D \rightarrow \mathbb{R}$  of the form

$$F(\mathbf{x}) = f \left( \sum_{k=1}^{\mathbf{x}(D)} \varphi_1(x_k), \dots, \sum_{k=1}^{\mathbf{x}(D)} \varphi_m(x_k) \right), \quad \mathbf{x} = \{x_1, \dots, x_{\mathbf{x}(D)}\} \in \mathbb{N}_f^D,$$

for some integer  $m \geq 1$ ,  $\varphi_1, \dots, \varphi_m \in \mathcal{C}^\infty(D)$ ,  $f \in \mathcal{C}_b^\infty(\mathbb{R}^m)$ . For  $F \in \mathcal{FC}_b^\infty(D)$ , we naturally define

$$\nabla_x^{\mathbb{N}_f} F(\mathbf{x}) := \sum_{l=1}^m \partial_l f \left( \sum_{k=1}^{\mathbf{x}(D)} \varphi_1(x_k), \dots, \sum_{k=1}^{\mathbf{x}(D)} \varphi_m(x_k) \right) \nabla \varphi_l(x),$$

$\mathbf{x} = \{x_1, \dots, x_{\mathbf{x}(D)}\} \in \mathbb{N}_f^D$ . We conclude this section with the following proposition.

**Proposition 4.4.** *Under (H1) – (H4), we have that  $\mathcal{FC}_b^\infty(D) \subset \text{Dom}(\overline{\mathcal{H}_D^{1/2}})$  and*

$$\bar{\mathcal{E}}_D(F, G) = \mathbb{E} \left[ \sum_{i=1}^{\mathbf{X}(D)} \nabla_{X_i}^{\mathbb{N}_f} F(\mathbf{X}^D) \cdot \nabla_{X_i}^{\mathbb{N}_f} G(\mathbf{X}^D) \right], \quad F, G \in \mathcal{FC}_b^\infty(D). \quad (4.5)$$

*Proof.* For  $F \in \mathcal{FC}_b^\infty(D)$ , we clearly have  $F^n(\mathbf{x}) := F(\mathbf{x}) \mathbf{1}_{\{|\mathbf{x}|(D) \leq n\}} \in \tilde{\mathcal{S}}_D$ , for any positive integer  $n$ . By a straightforward computation,  $F^m \rightarrow F$  in  $L_D^2$  as  $n$  goes to infinity. By the standard construction of the smallest closed extension of  $\mathcal{E}_D$  (see e.g. [9]), to get that  $F \in \text{Dom}(\bar{\mathcal{E}}_D)$  and that  $\bar{\mathcal{E}}_D(F, F) = \lim_{n \rightarrow \infty} \mathcal{E}_D(F^n, F^n)$ , it suffices to prove that  $\mathcal{E}_D(F^n - F^m, F^n - F^m)$  tends to zero as  $m, n$  go to infinity. This easily follows by the dominated convergence theorem. Indeed for some positive  $C > 0$  and  $m < n$ ,

$$\begin{aligned} \mathcal{E}_D(F^n - F^m, F^n - F^m) &= \mathbb{E} \left[ \sum_{i=1}^{\mathbf{X}(D)} \|\nabla_{X_i}^{\mathbb{N}_f} F(\mathbf{X}^D)\|^2 \mathbf{1}_{\{m < \mathbf{X}(D) \leq n\}} \right] \\ &\leq C \mathbb{E} [\mathbf{X}(D) \mathbf{1}_{\{m < \mathbf{X}(D) \leq n\}}] \xrightarrow{m, n \rightarrow \infty} 0. \end{aligned}$$

Now, noticing that by the monotone convergence theorem,

$$\mathcal{E}_D(F^n, F^n) = \mathbb{E} \left[ \sum_{i=1}^{\mathbf{X}(D)} \|\nabla_{X_i}^{\mathbb{N}_f} F(\mathbf{X}^D)\|^2 \mathbf{1}_{\{\mathbf{X}(D) \leq n\}} \right] \rightarrow \mathbb{E} \left[ \sum_{i=1}^{\mathbf{X}(D)} \|\nabla_{X_i}^{\mathbb{N}_f} F(\mathbf{X}^D)\|^2 \right],$$

we have (4.5) with  $G = F$  and we conclude by polarization.  $\square$

### 5. Diffusions Associated to Determinantal Processes on $D \subset S$

We start recalling some notions, see Chapters IV and V in [15]. Given  $\pi$  in the set  $\mathcal{P}(\check{\mathbb{N}}_f^D)$  of the probability measures on  $(\check{\mathbb{N}}_f^D, \mathcal{B}(\check{\mathbb{N}}_f^D))$ , we call a  $\pi$ -stochastic process with state space  $\check{\mathbb{N}}_f^D$  the collection

$$\mathbf{M}_{D, \pi} = (\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, (\mathbf{M}_t)_{t \geq 0}, (\mathbb{P}_{\mathbf{x}})_{\mathbf{x} \in \check{\mathbb{N}}_f^D}, \mathbb{P}_\pi)$$

where  $\mathcal{F} := \bigvee_{t \geq 0} \mathcal{F}_t$  is a  $\sigma$ -algebra on the set  $\Omega$ ,  $(\mathcal{F}_t)_{t \geq 0}$  is the  $\mathbb{P}_\pi$ -completed filtration generated by the process  $\mathbf{M}_t : \Omega \rightarrow \check{\mathbb{N}}_f^D$ ,  $\mathbb{P}_\mathbf{x}$  is a probability measure on  $(\Omega, \mathcal{F})$  for all  $\mathbf{x} \in \check{\mathbb{N}}_f^D$ , and  $\mathbb{P}_\pi$  is the probability measure on  $(\Omega, \mathcal{F})$  defined by

$$\mathbb{P}_\pi(A) := \int_{\check{\mathbb{N}}_f^D} \mathbb{P}_\mathbf{x}(A) \pi(d\mathbf{x}), \quad A \in \mathcal{F}.$$

A collection  $(\mathbf{M}_{D,\pi}, (\theta_t)_{t \geq 0})$  is called a  $\pi$ -time homogeneous Markov process with state space  $\check{\mathbb{N}}_f^D$  if  $\theta_t : \Omega \rightarrow \Omega$  is a shift operator, i.e.  $\mathbf{M}_s \circ \theta_t = \mathbf{M}_{s+t}$ ,  $s, t \geq 0$ ; the map  $\mathbf{x} \mapsto \mathbb{P}_\mathbf{x}(A)$  is measurable for all  $A \in \mathcal{F}$ ; the time homogeneous Markov property

$$\mathbb{P}_\mathbf{x}(\mathbf{M}_t \in A | \mathcal{F}_s) = \mathbb{P}_{\mathbf{M}_{t-s}}(\mathbf{M}_{t-s} \in A), \quad \mathbb{P}_\mathbf{x} - a.s., \quad A \in \mathcal{F}, \quad 0 \leq s \leq t, \quad \mathbf{x} \in \check{\mathbb{N}}_f^D$$

holds. A  $\pi$ -time homogeneous Markov process  $(\mathbf{M}_{D,\pi}, (\theta_t)_{t \geq 0})$  with state space  $\check{\mathbb{N}}_f^D$  is said to be  $\pi$ -tight on  $\check{\mathbb{N}}_f^D$  if  $(\mathbf{M}_t)_{t \geq 0}$  is right-continuous with left limits  $\mathbb{P}_\pi$ -almost surely;  $\mathbb{P}_\mathbf{x}(\mathbf{M}_0 = \mathbf{x}) = 1$ ,  $\mathbf{x} \in \check{\mathbb{N}}_f^D$ ; the filtration  $(\mathcal{F}_t)_{t \geq 0}$  is right continuous; the following strong Markov property holds:

$$\mathbb{P}_{\pi'}(\mathbf{M}_{t+\tau} \in A | \mathcal{F}_\tau) = \mathbb{P}_{\mathbf{M}_\tau}(\mathbf{M}_t \in A)$$

$\mathbb{P}_{\pi'}$ -almost surely for all  $\mathcal{F}_t$ -stopping time  $\tau$ ,  $\pi' \in \mathcal{P}(\check{\mathbb{N}}_f^D)$ ,  $A \in \mathcal{F}$  and  $t \geq 0$ , cf. Theorem IV.1.15 in [15]. In addition, a  $\pi$ -tight process on  $\check{\mathbb{N}}_f^D$  is said to be a  $\pi$ -special standard process on  $\check{\mathbb{N}}_f^D$  if for any  $\pi' \in \mathcal{P}(\check{\mathbb{N}}_f^D)$  which is equivalent to  $\pi$  and all  $\mathcal{F}_t$ -stopping times  $\tau$ ,  $(\tau_n)_{n \geq 1}$  such that  $\tau_n \uparrow \tau$  then  $\mathbf{M}_{\tau_n}$  converges to  $\mathbf{M}_\tau$ ,  $\mathbb{P}_{\pi'}$ -almost surely.

In the following,  $\mathbb{E}_\mathbf{x}$  denotes the expectation under  $\mathbb{P}_\mathbf{x}$ ,  $\mathbf{x} \in \check{\mathbb{N}}_f^D$ .

**Theorem 5.1.** *Assume (H1)–(H4). Then there exists a  $\mathbb{P}_D$ -tight special standard process  $(\mathbf{M}_{D,\mathbb{P}_D}, (\theta_t)_{t \geq 0})$  on  $\check{\mathbb{N}}_f^D$  such that:*

(1)  $\mathbf{M}_{D,\mathbb{P}_D}$  is a diffusion, in the sense that:

$$\mathbb{P}_\mathbf{x}(\{\omega \in \Omega : t \mapsto \mathbf{M}_t(\omega) \text{ is continuous on } [0, +\infty)\}) = 1, \quad \bar{\mathcal{E}}_D\text{-a.e. } \mathbf{x} \in \check{\mathbb{N}}_f^D \tag{5.1}$$

(see [15] for the definition of a property holding  $\bar{\mathcal{E}}_D$ -a.e.);

(2) the transition semigroup of  $\mathbf{M}_{D,\mathbb{P}_D}$  is given by

$$p_t F(\xi) := \mathbb{E}_\mathbf{x}[F(\mathbf{M}_t)], \quad \mathbf{x} \in \check{\mathbb{N}}_f^D, \quad F : \check{\mathbb{N}}_f^D \rightarrow \mathbb{R} \quad \text{square integrable,}$$

and it is properly associated with the Dirichlet form  $(\bar{\mathcal{E}}_D, \text{Dom}(\overline{\mathcal{H}}_D^{1/2}))$  in the sense that  $p_t F$  is an  $\bar{\mathcal{E}}_D$ -a.c. (see [15] for the notion of  $\bar{\mathcal{E}}_D$ -almost continuity),  $\mathbb{P}_D$ -version of  $\exp(-t\mathcal{H}_D^{gen})F$ , for all square integrable  $F : \check{\mathbb{N}}_f^D \rightarrow \mathbb{R}$  and  $t > 0$  (where  $\mathcal{H}_D^{gen}$  is the generator of  $\bar{\mathcal{E}}_D$ );

(3)  $\mathbf{M}_{D,\mathbb{P}_D}$  is unique up to  $\mathbb{P}_D$ -equivalence (we refer the reader to Definition 6.3 page 140 in [15] for the meaning of this notion);

(4)  $\mathbf{M}_{D,\mathbb{P}_D}$  is  $\mathbb{P}_D$ -symmetric, i.e.

$$\mathbb{E}[G(\mathbf{X}^D) p_t F(\mathbf{X}^D)] = \mathbb{E}[F(\mathbf{X}^D) p_t G(\mathbf{X}^D)],$$

for  $F, G \in L_D^2$ ;

(5)  $\mathbf{M}_{D,\mathbb{P}_D}$  has  $\mathbb{P}_D$  as invariant measure.

*Proof.* We apply Theorem 4.13 of [16, p. 308]. Using the notation of [16], set

$$S(f, g)(x) := \nabla f(x) \cdot \nabla g(x), \quad x \in \mathbb{R}^d, \quad f, g \in C^\infty(D),$$

where  $\nabla$  is the usual gradient on  $\mathbb{R}^d$ , and

$$S^\Gamma(F, G)(\mathbf{x}) := \sum_{i=1}^{\mathbf{x}(D)} \nabla_{x_i}^{N_{if}} F(\mathbf{x}) \cdot \nabla_{x_i}^{N_{if}} G(\mathbf{x}), \quad \mathbf{x} = \{x_1, \dots, x_{\mathbf{x}(D)}\} \in N_f^D, \quad F, G \in \tilde{\mathcal{S}}_D.$$

Then, it is readily seen that  $(S, C^\infty(D))$  satisfies conditions (D.1), (S.1), (S.2) and (S.3) of [16], and  $(S^\Gamma, \tilde{\mathcal{S}}_D)$  satisfies condition  $(S^\Gamma.\mu)$  of [16, p. 282]. Furthermore,  $\mathbb{P}_D$  satisfies condition  $(\mu.1)$  of [16, p. 282] and condition (Q) of [16] holds since  $\mathbb{R}^d$  is complete (see Example 4.5.1 in [16]). The assumptions of Theorem 4.13 are therefore verified and the proof is completed.  $\square$

*Non-collision property of the associated diffusions.* In the following, we will show the non-collision property of the diffusion constructed in the previous theorem which, roughly speaking, means that the diffusion takes values on  $N_f^D$ .

We start by recalling the following lemma, which is borrowed from [20].

**Lemma 5.2.** *Assume (H1) – (H4) and let  $(\mathbf{M}_{D, \mathbb{P}_D}, (\theta_t)_{t \geq 0})$  be the diffusion given by Theorem 5.1. Let  $u_n \in \text{Dom}(\bar{\mathcal{E}}_D)$ ,  $n \geq 1$ , be such that:  $u_n : \ddot{N}_f^D \rightarrow \mathbb{R}$  is continuous,  $u_n \rightarrow u$  point-wise in  $\ddot{N}_f^D$ ,*

$$\sup_{n \geq 1} \bar{\mathcal{E}}_D(u_n, u_n) < \infty. \tag{5.2}$$

*Then  $u$  is  $\bar{\mathcal{E}}_D$ -a.c. and, in particular,*

$$\mathbb{P}_{\mathbf{x}}(\{\omega \in \Omega : t \mapsto u(\mathbf{M}_t)(\omega) \text{ is continuous on } [0, +\infty)\}) = 1, \quad \bar{\mathcal{E}}_D\text{-a.e. } \mathbf{x} \in N_f^D.$$

The next theorem provides the non-collision property.

**Theorem 5.3.** *Assume  $d \geq 2$ , and (H1) – (H4). Then*

$$\mathbb{P}_{\mathbf{x}}(\{\omega \in \Omega : \mathbf{M}_t(\omega) \in N_f^D \quad \forall 0 \leq t < \infty\}) = 1, \quad \bar{\mathcal{E}}_D\text{-a.e. } \mathbf{x} \in N_f^D.$$

*Proof.* Since the proof is similar to the proof of Proposition 1 in [20] we skip some details. For every positive integer  $a$ , define  $u := \mathbf{1}_N$ , where

$$N := \{\mathbf{x} \in \ddot{N}_f^D : \sup_{x \in [-a, a]^d} \mathbf{x}(\{x\}) \geq 2\}.$$

The claim follows if we prove that  $u$  is  $\bar{\mathcal{E}}_D$ -a.c.. For this we are going to apply Lemma 5.2. Define

$$u_n(\mathbf{x}) = \Psi \left( \sup_{i \in A_n} \sum_{x \in \mathbf{x}} \phi_i(x) \right), \quad n \geq 1,$$

where  $\Psi \in C_b^\infty(\mathbb{R})$  and  $\phi_i \in C^\infty(D)$  are chosen as in the proof of Proposition 1 in [20], and  $A_n := \mathbb{Z}^d \cap [-na, na]^d$ . Note that  $u_n \in \text{Dom}(\bar{\mathcal{H}}_D^{1/2})$  by Proposition 4.4. Furthermore,  $u_n : \ddot{N}_f^D \rightarrow \mathbb{R}$  is continuous and  $u_n \rightarrow u$  point-wise by the proof of

Proposition 1 in [20]. It remains to check (5.2). For  $i = (i^{(1)}, \dots, i^{(d)}) \in \mathbb{Z}^d$  and  $n \geq 1$ , we denote by  $I_i^{(n)}$  the function defined by

$$I_i^{(n)}(x) := \prod_{k=1}^d \mathbf{1}_{[-1/2, 3/2]}(nx^{(k)} - i^{(k)}), \quad x = (x^{(1)}, \dots, x^{(d)}) \in D.$$

As proved in Proposition 1 in [20], the following upper bound holds:

$$\bar{\mathcal{E}}_D(u_n, u_n) \leq Cn^2 \sum_{i \in A_n} \mathbb{E} \left[ \mathbf{1}_{\{\sum_{j=1}^{\mathbf{X}(D)} I_i^{(n)}(X_j) \geq 2\}} \sum_{j=1}^{\mathbf{X}(D)} I_i^{(n)}(X_j) \right], \quad (5.3)$$

where  $C > 0$  is a positive constant.

Now, we upper-bound the r.h.s. of (5.3) by stochastic domination. In [11], it is proved that

$$c[D](x, \mathbf{x}) \leq J[D](x, x), \quad x \in D, \mathbf{x} \in \mathbb{N}_f^D,$$

where we denote by  $c[D](x, \mathbf{x})$  the Papangelou conditional intensity of  $\mathbf{X}^D$ , see [5]. Note that a Poisson process  $\mathbf{Y}^D$  of mean measure  $J[D](x, x)\mu(dx)$  has Papangelou conditional intensity  $J[D](x, x)$ . Therefore, by Theorem 1.1 in [10], we have that  $\mathbf{X}^D$  is stochastically dominated by  $\mathbf{Y}^D$ , in the sense that

$$\mathbb{E}[f(\mathbf{X}^D)] \leq \mathbb{E}[f(\mathbf{Y}^D)] \quad (5.4)$$

for any integrable  $f : \mathbb{N}_f \rightarrow \mathbb{R}$  such that  $f(\mathbf{x}) \leq f(\mathbf{y})$  whenever  $\mathbf{x} \subset \mathbf{y} \in \mathbb{N}_f$ . Since for any  $i \in A_n$ , the mapping  $\mathbf{x} \mapsto \mathbf{1}_{\{\sum_{j=1}^{\mathbf{x}(D)} I_i^{(n)}(x_j) \geq 2\}} \sum_{j=1}^{\mathbf{x}(D)} I_i^{(n)}(x_j)$  is increasing, we then have

$$\bar{\mathcal{E}}_D(u_n, u_n) \leq Cn^2 \sum_{i \in A_n} \mathbb{E} \left[ \mathbf{1}_{\{\sum_{j=1}^{\mathbf{Y}(D)} I_j^{(n)}(Y_j) \geq 2\}} \sum_{j=1}^{\mathbf{Y}(D)} I_i^{(n)}(Y_j) \right]. \quad (5.5)$$

By the properties of the Poisson process, the right hand side of (5.5) is equal to

$$Cn^2 \sum_{i \in A_n} \left( 1 - e^{-\int_D I_i^{(n)}(x) J[D](x, x) \mu(dx)} \right) \int_D I_i^{(n)}(x) J[D](x, x) \mu(dx),$$

which is bounded above by

$$Cn^2 \sum_{i \in A_n} \left( \int_D I_i^{(n)}(x) J[D](x, x) \rho(x) dx \right)^2.$$

By using the Cauchy-Schwarz inequality, this term is further bounded by

$$Cn^2 \sum_{i \in A_n} \left( \int_D I_i^{(n)}(x) dx \right) \left( \int_D I_i^{(n)}(x) J[D](x, x)^2 \rho(x)^2 dx \right). \quad (5.6)$$

We have on the one hand that

$$\int_{\mathbb{R}^d} I_i^{(n)}(x) dx = (2/n)^d$$

and on the other hand for some constant  $C' > 0$ ,

$$\int_D I_i^{(n)}(x) J[D](x, x)^2 \rho(x)^2 dx \leq C'n^{-d},$$

since  $J$  and  $\rho$  are bounded on  $D$ . Moreover,  $\sharp(A_n) \leq (2an)^d$ . Consequently, the quantity (5.6) is in turn bounded by  $C'' n^{2-d}$ , for some constant  $C'' > 0$ . The claim follows by the assumption  $d \geq 2$ .  $\square$

### 6. An Illustrating Example

Let  $S := B(0, 1)$  and  $D := B(0, R) \subset \mathbb{R}^2$  be the closed ball centered at the origin with radius  $R \in (0, 1)$  and  $(\varphi_k^{(R)})_{1 \leq k \leq 3}$ , the orthonormal subset of  $L^2(B(0, R), \ell)$  defined by

$$\varphi_k^{(R)}(x) := \frac{1}{R} \sqrt{\frac{k+1}{\pi}} \left( \frac{x^{(1)}}{R} + i \frac{x^{(2)}}{R} \right)^k, \quad x = (x^{(1)}, x^{(2)}) \in B(0, R), \quad k = 1, 2, 3,$$

where  $\ell$  is the Lebesgue measure and  $i := \sqrt{-1}$  denotes the complex unit. In this example, we consider the truncated Bergman kernel with at most 3 points (see e.g. [13]) restricted to  $D = B(0, R)$ :

$$K_D(x, y) := \sum_{k=1}^3 R^{2(k+1)} \varphi_k^{(R)}(x) \overline{\varphi_k^{(R)}(y)}, \quad x, y \in D,$$

and denote by  $\mathcal{K}_D$  the associated integral operator, which is easily seen to be Hermitian and trace class with non-zero eigenvalues  $\kappa_k := R^{2(k+1)}$ ,  $k = 1, 2, 3$ . As a consequence, the spectrum of  $\mathcal{K}_D$  is contained in  $[0, 1)$  and the triplet  $(\mathcal{K}_D, K_D, \ell)$  satisfies **(H1)**. In addition, **(H2)** is trivially satisfied since the reference measure is the Lebesgue measure. The Janossy densities of  $\mathbf{X}^D$  defined in (2.2) are given by

$$j_D^n(x_1, \dots, x_n) = \text{Det}(\mathbf{Id} - \mathcal{K}_D) \det J[D](x_1, \dots, x_n), \quad n = 1, 2, \quad (x_1, \dots, x_n) \in D^n,$$

where the kernel  $J[D]$  is given by

$$J[D](x, y) := \sum_{h=1}^3 \frac{R^{2(h+1)}}{1 - R^{2(h+1)}} \varphi_h^{(R)}(x) \overline{\varphi_h^{(R)}(y)},$$

cf. (2.6). Since  $\mathbf{X}^D$  has at most 3 points, see e.g. [23], we have  $j_D^n = 0$ , for  $n \geq 4$ . To prove condition **(H3)** it suffices to remark that the function

$$(x_1, \dots, x_n) \rightarrow \det(J[D](x_p, x_q))_{1 \leq p, q \leq n}$$

is continuously differentiable on  $D^n$ , for  $n = 1, 2, 3$ . To show that **(H4)** is verified, we first consider the case  $n = 3$ . Note that

$$(J[D](x_p, x_q))_{1 \leq p, q \leq 3} = A(x_1, x_2, x_3) A(x_1, x_2, x_3)^*,$$

where the matrix  $A(x_1, x_2, x_3) := (A_{ph})_{1 \leq p, h \leq 3}$  is given by

$$A_{ph} := \frac{R^{h+1}}{\sqrt{1 - R^{2(h+1)}}} \varphi_h^{(R)}(x_p)$$

and  $A(x_1, x_2, x_3)^*$  denotes the transpose conjugate of  $A(x_1, x_2, x_3)$ . Hence,

$$\det J[D](x_1, x_2, x_3) = |\det A(x_1, x_2, x_3)|^2,$$

and since the previous determinant can be rewritten involving a Vandermonde determinant, we have

$$\begin{aligned} & \det A(x_1, x_2, x_3) \\ &= \prod_{p=1}^3 \sqrt{\frac{1+p}{\pi(1-R^{2(p+1)})}} (x_p^{(1)} + ix_p^{(2)}) \prod_{1 \leq p < q \leq 3} ((x_p^{(1)} - x_q^{(1)}) + i(x_p^{(2)} - x_q^{(2)})). \end{aligned}$$

Note that **(H4)** with  $n = 3$  is exactly

$$\int_{D^3} \left| \frac{\partial_{x_i^{(h)}} |\det A(x_1, x_2, x_3)|^2 \partial_{x_j^{(k)}} |\det A(x_1, x_2, x_3)|^2}{|\det A(x_1, x_2, x_3)|^2} \right| \mathbf{1}_{\{|\det A(x_1, x_2, x_3)| > 0\}} dx_1 dx_2 dx_3,$$

which is finite for all  $1 \leq i, j \leq 3$  and  $1 \leq h, k \leq 2$ , and so it suffices to check

$$\int_{B(0,R)^3} \left| \frac{\partial_{x_1^{(1)}} |\det A(x_1, x_2, x_3)|^2}{|\det A(x_1, x_2, x_3)|^2} \right| \mathbf{1}_{\{|\det A(x_1, x_2, x_3)| > 0\}} dx_1 dx_2 dx_3 < \infty.$$

This latter integral reduces to

$$\int_{B(0,R)^3} \left| \frac{2x_1^{(1)}}{(x_1^{(1)})^2 + (x_1^{(2)})^2} + 2 \sum_{j=2}^3 \frac{x_1^{(1)} - x_j^{(1)}}{(x_1^{(1)} - x_j^{(1)})^2 + (x_1^{(2)} - x_j^{(2)})^2} \right| dx_1 dx_2 dx_3,$$

which is indeed finite. Consequently, we proved that **(H4)** is verified for  $n \geq 3$  (indeed it is trivially satisfied for  $n > 3$ ). Now, consider  $n = 1, 2$ . We have again

$$J[D](x_1, \dots, x_n) = A(x_1, \dots, x_n)A(x_1, \dots, x_n)^*,$$

where this time,  $A(x_1, \dots, x_n)$  is a rectangular  $n \times 3$  matrix given by

$$A(x_1, \dots, x_n) = (A_{ph})_{1 \leq p \leq n, 1 \leq h \leq 3} := \left( \frac{R^{h+1}}{\sqrt{1-R^{2(h+1)}}} \varphi_h^{(R)}(x_p) \right)_{1 \leq p \leq n, 1 \leq h \leq 3}.$$

Recall the Cauchy-Binet formula:

$$\det J[D](x_1, \dots, x_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_n \leq 3} |\det A^{i_1, \dots, i_n}(x_1, \dots, x_n)|^2, \tag{6.1}$$

where

$$A_{ph}^{i_1, \dots, i_n} := \frac{R^{i_h+1}}{\sqrt{1-R^{2(i_h+1)}}} \varphi_{i_h}^{(R)}(x_p), \quad 1 \leq p, h \leq n,$$

defines a square matrix. We now consider fixed  $1 \leq i_1 < i_2 < \dots < i_n \leq 3$  and evaluate  $|\det A^{i_1, \dots, i_n}(x_1, \dots, x_n)|^2$ . We note that

$$|\det A^{i_1, \dots, i_n}(x_1, \dots, x_n)|^2 = \prod_{p=1}^n \frac{1+i_p}{\pi(1-R^{2(i_p+1)})} |V_{i_1, \dots, i_n}(x_1, \dots, x_n)|^2, \tag{6.2}$$

where

$$V_{i_1, \dots, i_n}(x_1, \dots, x_n) := \det \left( \left( x_h^{i_p} \right)_{1 \leq p, h \leq n} \right)$$

is known in the literature as the generalized Vandermonde determinant. By definition, the Schur polynomial  $s_\lambda$  is the ratio between the generalized Vandermonde determinant and the classical Vandermonde determinant. More precisely,

$$\det \left( \left( x_h^{i_p} \right)_{1 \leq p, h \leq n} \right) = \det \left( \left( x_h^{p-1} \right)_{1 \leq p, h \leq n} \right) s_{\lambda(i_1, \dots, i_n)}(x_1, \dots, x_n), \quad (6.3)$$

where  $\lambda(i_1, \dots, i_n) := (i_n - n + 1, \dots, i_2 - 1, i_1)$ . Combining (6.1), (6.2) and (6.3), we have

$$\det J[D](x_1, \dots, x_n) = \left| \prod_{1 \leq p < q \leq n} ((x_p^{(1)} - x_q^{(1)}) + i(x_p^{(2)} - x_q^{(2)})) \right|^2 \sum_{1 \leq i_1 < i_2 < \dots < i_n \leq 3} \left( \prod_{p=1}^n \frac{1 + i_p}{\pi(1 - R^{2(i_p+1)})} \right) |s_{\lambda(i_1, \dots, i_n)}(x_1, \dots, x_n)|^2.$$

For  $n = 1$ , one has  $s_{\lambda(1)}(x) = x$ ,  $s_{\lambda(2)}(x) = x^2$  and  $s_{\lambda(3)}(x) = x^3$ , see e.g. [12], and therefore

$$\det J[D](x) = \frac{2}{\pi(1 - R^4)} |x|^2 + \frac{3}{\pi(1 - R^6)} |x|^4 + \frac{4}{\pi(1 - R^8)} |x|^6.$$

Thus,

$$\begin{aligned} \partial_{x^{(p)}} \ln(\det J[D](x)) &= \frac{2}{\pi(1 - R^4)} \frac{2x^{(p)}}{\frac{2}{\pi(1 - R^4)} |x|^2 + \frac{3}{\pi(1 - R^6)} |x|^4 + \frac{4}{\pi(1 - R^8)} |x|^6} \\ &+ \frac{3}{\pi(1 - R^6)} \frac{4x^{(p)} |x|^2}{\frac{2}{\pi(1 - R^4)} |x|^2 + \frac{3}{\pi(1 - R^6)} |x|^4 + \frac{4}{\pi(1 - R^8)} |x|^6} \\ &+ \frac{4}{\pi(1 - R^8)} \frac{6x^{(p)} |x|^4}{\frac{2}{\pi(1 - R^4)} |x|^2 + \frac{3}{\pi(1 - R^6)} |x|^4 + \frac{4}{\pi(1 - R^8)} |x|^6}, \end{aligned}$$

and therefore

$$|\partial_{x^{(p)}} \ln(\det J[D](x))| \leq \frac{2|x^{(p)}|}{|x|^2} + \frac{4|x^{(p)}|}{|x|^2} + \frac{6|x^{(p)}|}{|x|^2},$$

which is integrable on  $D$ . For  $n = 2$ , one has  $s_{\lambda(1,2)}(x, y) = 1$ ,  $s_{\lambda(1,3)}(x, y) = x + y$  and  $s_{\lambda(2,3)}(x, y) = xy$ , see e.g. [12], and therefore

$$\det J[D](x, y) = |x - y|^2 \left( \frac{2}{\pi(1 - R^4)} \frac{3}{\pi(1 - R^6)} + \frac{2}{\pi(1 - R^4)} \frac{4}{\pi(1 - R^8)} |x + y|^2 + \frac{3}{\pi(1 - R^6)} \frac{4}{\pi(1 - R^8)} |xy|^2 \right).$$

Note that the differential of the logarithm of  $|x - y|^2$  gives rise to a locally integrable term. So it remains to check that the differential of the logarithm of the second term, hereafter denoted by  $\Psi_D(x, y)$ , is integrable on  $D^2$ . By symmetry



of the Schur polynomials, it suffices to check that the derivative of  $\Psi_D(x, y)$  with respect to  $x^{(p)}$  is integrable on  $D^2$ . We have

$$|\partial_{x^{(p)}} \Psi_D(x, y)| \leq \frac{2|x^{(p)} + y^{(p)}|}{|x + y|^2} + \frac{2|x^{(p)}||y|^2}{|xy|^2} = \frac{2|x^{(p)} + y^{(p)}|}{|x + y|^2} + \frac{2x^{(p)}}{|x|^2},$$

and the claim follows by noticing that the r.h.s. is integrable on  $D^2$ .

## References

1. Albeverio, S., Kondratiev, Yu. G, and and Röckner, M.: Analysis and geometry on configuration spaces, *J. Funct. Anal.* **154** (1998), no. 2, 444–500.
2. Albeverio, S., Kondratiev, Yu. G, and and Röckner, M.: Analysis and geometry on configuration spaces: the Gibbsian case, *J. Funct. Anal.* **157** (1998), no. 1, 242–291.
3. Anderson, R. M.: A nonstandard representation for Brownian motion motion and Itô integration, *Israel J. Math.* **25** (1976), 15–46.
4. Camilier, I. and Decreusefond, L.: Quasi-invariance and integration by parts for determinantal and permanental processes, *J. Funct. Anal.* **259** (2010), no. 1, 268–300.
5. Daley, D. J. and Vere-Jones, D.: *An Introduction to the Theory of Point Processes*. Vol. I, Probability and its Applications, Springer-Verlag, New York, 2003.
6. Daley, D. J. and Vere-Jones, D.: *An Introduction to the Theory of Point Processes*. Vol. II, second ed., Probability and its Applications (New York), Springer, New York, 2008.
7. Dunford, N. and Schwartz, J. T.: *Linear Operators*, vol. II, Interscience publishers, New York, 1957.
8. Flint, I., Lu, X., Privault, N., Niyato, D., and Wang, P.: Performance analysis of ambient RF energy harvesting with repulsive point process modeling, to appear in *IEEE Transactions on Wireless Communications*, 30 pages, 2015.
9. Fukushima, M.: *Dirichlet Forms and Markov Processes*, North-Holland, 1980.
10. Georgii, H. O.: *Canonical Gibbs Measures*, Lecture Notes in Mathematics, vol. 760, Springer-Verlag, 1979.
11. Georgii, H. O. and Yoo, H. J.: Conditional intensity and Gibbsianness of determinantal point processes, *J. Stat. Phys.* **118** (2005), no. 1-2, 55–84.
12. Heineman, E. R.: Generalized Vandermonde determinants, *Trans. Amer. Math. Soc.* **31** (1929), 464–476.
13. Hough, J., Krishnapur, M., Peres, Y., and Virág, B.: *Zeros of Gaussian Analytic Functions and Determinantal Point Processes*, University Lecture Series, vol. 51, American Mathematical Society, Providence, RI, 2009.
14. van Lieshout, M. N. M.: *Markov Point Processes and Their Applications*, Imperial College Press, London, 2000.
15. Ma, Z. M. and Röckner, M.: *An Introduction to the Theory of (Non-symmetric) Dirichlet Forms*, Springer-Verlag, 1992.
16. Ma, Z. M. and Röckner, M.: Construction of diffusions on configuration spaces, *Osaka J. Math.* **37** (2000), no. 2, 273–314.
17. Macchi, O.: The coincidence approach to stochastic point processes, *Advances in Appl. Probability* **7** (1975), 83–122.
18. Møller, J. and Waagepetersen, R. P.: *Statistical Inference and Simulation for Spatial Point Processes*, Monographs on Statistics and Applied Probability, vol. 100, Chapman & Hall/CRC, Boca Raton, FL, 2004.
19. Privault, N. and Torrisi, G. L.: Density estimation of functionals of spatial point processes with application to wireless networks, *SIAM J. Math. Anal.* **43** (2011), 1311–1344.
20. Röckner, M. and Schmuland, B.: *A support property for infinite-dimensional interacting diffusion processes*, C. R. Acad. Sci. Paris Sér. I Math. **326** (1998), 359–364.
21. Shirai, T. and Takahashi, Y.: Random point fields associated with certain Fredholm determinants. I. Fermion, Poisson and boson point processes, *J. Funct. Anal.* **205** (2003), no. 2, 414–463.

22. Simon, B.: *Trace Ideals and Their Applications*, second ed., Mathematical Surveys and Monographs, vol. 120, American Mathematical Society, Providence, RI, 2005.
23. Soshnikov, A.: Determinantal random point fields, *Uspekhi Mat. Nauk* **55** (2000), no. 5, 107–160.
24. Surgailis, D.: On multiple Poisson stochastic integrals and associated Markov semi-groups, *Probability and Mathematical Statistics* **3** (1984), 217–239.
25. Tamura, H. and Ito, K. R.: A canonical ensemble approach to the fermion/boson random point processes and its applications, *Comm. Math. Phys.* **263** (2006), no. 2, 353–380.
26. Torrisi, G. L.: Point processes with Papangelou conditional intensity: from the Skorohod integral to the Dirichlet form, *Markov Processes and Related Fields* **19** (2013), 195–248.
27. Torrisi, G. L.: Quasi-invariance of fermion processes with  $J$ -hermitian kernel, *Commun. Stoch. Anal.* **8** (2014), no. 2, 169–190.
28. Torrisi, G. L. and Leonardi, E.: Large deviations of the interference in the Ginibre network model, *Stochastic Systems* **4** (2014), 173–205.
29. Vergne, A., Flint, I., Decreusefond, L., and Martins, P.: *Disaster Recovery in Wireless Networks: A Homology-based Algorithm*, ICT 2014.
30. Yoo, H. J.: Dirichlet forms and diffusion processes for fermion random point fields, *J. Funct. Anal.* **219** (2005), no. 1, 143–160.

LAURENT DECREUSEFOND: TELECOM PARISTECH, 23 AVENUE D'ITALIE, 75013 PARIS, FRANCE.  
*E-mail address:* laurent.decreusefond@mines-telecom.fr

IAN FLINT: DIVISION OF MATHEMATICAL SCIENCES, NANYANG TECHNOLOGICAL UNIVERSITY,  
SPMS-MAS-05-43, 21 NANYANG LINK SINGAPORE 637371  
*E-mail address:* iflint@ntu.edu.sg

NICOLAS PRIVAULT: DIVISION OF MATHEMATICAL SCIENCES, NANYANG TECHNOLOGICAL UNIVERSITY,  
SPMS-MAS-05-43, 21 NANYANG LINK SINGAPORE 637371  
*E-mail address:* nprivault@ntu.edu.sg

GIOVANNI LUCA TORRISI: ISTITUTO PER LE APPLICAZIONI DEL CALCOLO "MAURO PICONE",  
CNR, VIA DEI TAURINI 19, 00185 ROMA, ITALY  
*E-mail address:* torrisi@iac.rm.cnr.it