

1-1-2019

Dosage compensation and gene expression of the X Chromosome in sheep

Jingyue Duan
University of Connecticut

Kaleigh Flock
University of Connecticut

Nathanial Jue
California State University, Monterey Bay

Mingyuan Zhang
University of Connecticut

Amanda Jones
University of Connecticut

See next page for additional authors

Follow this and additional works at: https://digitalcommons.lsu.edu/animalsciences_pubs

Recommended Citation

Duan, J., Flock, K., Jue, N., Zhang, M., Jones, A., Al Seesi, S., Mandoiu, I., Pillai, S., Hoffman, M., O'Neill, R., Zinn, S., Govoni, K., Reed, S., Jiang, H., Jiang, Z., & Tian, X. (2019). Dosage compensation and gene expression of the X Chromosome in sheep. *G3: Genes, Genomes, Genetics*, 9 (1), 305-314. <https://doi.org/10.1534/g3.118.200815>

This Article is brought to you for free and open access by the School of Animal Sciences at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Authors

Jingyue Duan, Kaleigh Flock, Nathaniel Jue, Mingyuan Zhang, Amanda Jones, Sahar Al Seesi, Ion Mandoiu, Sambhu Pillai, Maria Hoffman, Rachel O'Neill, Steven Zinn, Kristen Govoni, Sarah Reed, Hesheng Jiang, Zongliang Jiang, and Xiuchun Tian

Dosage Compensation and Gene Expression of the X Chromosome in Sheep

Jingyue (Ellie) Duan,^{*,1} Kaleigh Flock,^{*,1} Nathaniel Jue,[†] Mingyuan Zhang,^{*,*,#} Amanda Jones,^{*}

Sahar Al Seesi,^{§,**,†} Ion Mandoiu,^{**,†} Sambhu Pillai,^{*} Maria Hoffman,^{*} Rachel O'Neill,^{††} Steven Zinn,^{*}

Kristen Govoni,^{*} Sarah Reed,^{*} Hesheng Jiang,^{††} Zongliang (Carl) Jiang^{*,§§} and Xiuchun (Cindy) Tian^{*,2}

^{*}Department of Animal Science, ^{**}Department of Computer Science, ^{††}Department of Molecular and Cell Biology, and University of Connecticut, Storrs, CT, 06269, [†]School of Natural Sciences, California State University, Monterey Bay, Seaside, CA 93955, [‡]Laboratory Animal Center, Guangxi Medical University, Nanning 530021, China, [§]Smith College Department of Computer Science, Northampton, MA 01063, ^{††}College of Animal Science and Technology, Guangxi University, Nanning 530004, China, and ^{§§}School of Animal Science, Louisiana State University, Baton Rouge, LA 70803

ORCID IDs: 0000-0001-6416-2250 (J.E.D.); 0000-0001-6143-2200 (M.Z.); 0000-0002-1258-8687 (A.J.); 0000-0002-4818-0237 (I.M.); 0000-0002-9961-6280 (S.R.); 0000-0002-6168-1810 (H.J.); 0000-0002-3040-7771 (Z.C.J.); 0000-0002-8441-2154 (X.C.T.)

ABSTRACT Ohno's hypothesis predicts that the expression of the single X chromosome in males needs compensatory upregulation to balance its dosage with that of the diploid autosomes. Additionally, X chromosome inactivation ensures that quadruple expression of the two X chromosomes is avoided in females. These mechanisms have been actively studied in mice and humans but lag behind in domestic species. Using RNA sequencing data, we analyzed the X chromosome upregulation in sheep fetal tissues from day 135 of gestation under control, over or restricted maternal diets (100%, 140% and 60% of National Research Council Total Digestible Nutrients), and in conceptuses, juvenile, and adult somatic tissues. By computing the mean expression ratio of all X-linked genes to all autosomal genes (X:A), we found that all samples displayed some levels of X chromosome upregulation. The degrees of X upregulation were not significant (P -value = 0.74) between ovine females and males in the same somatic tissues. Brain, however, displayed complete X upregulation. Interestingly, the male and female reproduction-related tissues exhibited divergent X dosage upregulation. Moreover, expression upregulation of the X chromosome in fetal tissues was not affected by maternal diets. Maternal nutrition, however, did change expression levels of several X-linked genes, such as sex determination genes *SOX3* and *NROB1*. In summary, our results showed that X chromosome upregulation occurred in nearly all sheep somatic tissues analyzed, thus support Ohno's hypothesis in a new species. However, the levels of upregulation differed by different subgroups of genes such as those that are house-keeping and "dosage-sensitive".

KEYWORDS

Ohno's hypothesis
X chromosome upregulation
Maternal nutrition
Ovine

In mammals, deviation from diploidy may induce detrimental consequences (Birchler *et al.* 2005). For example, gene duplications or

deletions can induce cancer (Giam and Rancati 2015), and chromosome monosomy or trisomy usually causes fetal lethality (Burgess *et al.* 2014). Mammalian males, however, are monosomic for the X chromosome, yet do not suffer from the deleterious effects of X monosomy (Chen *et al.* 2014). This is likely the result of a still debated mechanism of X chromosome dosage compensation (Pessia *et al.* 2014; Mank *et al.* 2014; Veitia *et al.* 2015; Graves 2016). Susumu Ohno hypothesized that upregulation of the X-linked genes in the heterogametic sex (XY) would be necessary to maintain their expression to the levels of the diploid autosomes (Ohno 1966). This solved the dosage imbalance of X-linked genes in males, yet subjected females to quadruple levels of X expression. Another mechanism, X chromosome inactivation (XCI), the inactivation of one of the two X chromosomes in every cell of the female, balances the X chromosome gene dosage between males and females

Copyright © 2019 Duan *et al.*

doi: <https://doi.org/10.1534/g3.118.200815>

Manuscript received October 17, 2018; accepted for publication November 26, 2018; published Early Online November 27, 2018.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Supplemental material available at Figshare: <https://doi.org/10.25387/g3.7221467>.

¹These authors contributed equally to this work.

²Corresponding Author: Department of Animal Science, University of Connecticut, 1390 Storrs Rd, Storrs, CT 06269. E-mail: xiuchun.tian@uconn.edu.

(Lyon 1961). Both X chromosome upregulation and XCI are necessary components of the X chromosome dosage compensation in mammals (Ohno 1966).

Although XCI has been characterized in many species (Goto and Monk 1998; Xue *et al.* 2002; Deakin *et al.* 2009; Lee 2011; Livernois *et al.* 2012; Sahakyan *et al.* 2018), verification of X chromosome upregulation has not been conducted until chromosome-wide expression analysis became possible (Pessia *et al.* 2014). X chromosome upregulation has been studied using data from microarray (Gupta *et al.* 2006; Nguyen and Disteché 2006) and RNA sequencing (RNA-seq) (Xiong *et al.* 2010; Deng *et al.* 2011; Lin *et al.* 2007, 2011, 2012; Pessia *et al.* 2012) by computing the mean or median expression ratio of all X-linked genes to all autosomal genes (X:A). An X:A ratio of 1.0 or greater implies the doubling of X-linked gene transcription from the single active X, indicating complete compensatory upregulation. An X:A ratio of 0.5 indicates that expression levels of X-linked genes is half of those of the autosome pairs, suggesting no compensatory upregulation. When an X:A ratio falls between 0.5 and 1, it is termed partial compensation (Deng *et al.* 2011). A full compensatory upregulation of X-linked genes has been recently observed only in “dosage-sensitive” genes in eutherian mammals (Julien *et al.*, 2012; Lin *et al.*, 2012; Pessia *et al.*, 2012). These genes usually code for proteins complexes with structural, regulatory and housekeeping functions (Kondrashov and Koonin 2004; Birchler 2012; Pessia *et al.* 2012).

XCI and X chromosome upregulation have been studied in mice and humans, but such investigations lag behind in domestic species (Disteché 2012). XCI has been shown in sheep (Luciani *et al.* 1979), but little is known about its onset and no information is available on X dosage compensation. With completion of the ovine genome sequencing (Jiang *et al.* 2014) and the advancement of RNA-seq technology (Wolf and Bryk 2011), a number of RNA-seq datasets in sheep somatic tissues are now available for X chromosome upregulation studies.

The sheep has been frequently used as a model for human pregnancy and fetal development (Barry and Anthony 2008). Poor maternal nutrition, either over- or restricted-feeding, has been shown to alter gene expression in fetal tissues (Du *et al.* 2011; Pillai *et al.* 2016; Duan *et al.* 2018). Changes in DNA methylation is likely involved because restrictedly nourished ewes carried fetuses with altered DNA methyltransferase in the hypothalamus (Begum *et al.* 2012). Similarly, human metastable epialleles, which are variably expressed in genetically identical individuals, have also been persistently changed epigenetically by maternal nutrition in early pregnancy (Dominguez-Salas *et al.* 2014). These findings that maternal diet alters fetal epigenetics are of particular interest because XCI and X chromosome upregulation are epigenetically regulated processes (Goto and Monk 1998). However, the effects of maternal nutrition on X chromosome dosage compensation and X-linked gene expression have yet to be studied in the sheep, an important species for both agriculture and human medicine.

Using data generated by us (GSE111306) (Duan *et al.* 2018) and two additional RNA-seq datasets, PRJEB6169 (Jiang *et al.* 2014) and PRJNA254105 (Brooks *et al.* 2015), we were able to achieve the first comprehensive evaluation of X chromosome upregulation in sheep. Furthermore, we also investigated the effects of different maternal diets on the expression of X-linked genes. Our hypothesis was that X chromosome upregulation in the sheep would be partial, similar to that in the bovine as reported by us (Duan *et al.* 2016) and others (Ka *et al.* 2016). We further hypothesized that different maternal diets would alter the expression levels of X-linked genes in ovine fetal tissues.

MATERIALS AND METHODS

Experimental design and RNA sequencing

Animal protocols, tissues collection, and RNA sequencing library preparation were described in Pillai *et al.* (2017) and Duan *et al.* (2018). Briefly, 12 pregnant ewes were individually housed and randomly assigned to control- (100% NRC requirement, Con, $n = 4$), overfed- (140%, Over, $n = 4$) or restricted- (60%, Res, $n = 4$) diets calculated by the National Research Council requirement for total digestible nutrients for a ewe pregnant with twins (Pillai *et al.* 2016). The ewes remained on their respective diets until day 135 of gestation when they were killed. Fifteen fetuses, control ($n = 7$), overfed ($n = 4$) and restricted ($n = 4$) were included in this study. Full organ of brain, kidney, and lung were collected, flash-frozen in liquid nitrogen and stored at -80° until RNA extraction.

RNA was extracted from fetal brain, kidney, and lung using TRIzol (Invitrogen, Grand Island, NY) according to the manufacturer's instructions. Library preparation was carried out using TruSeq RNA library prep kit (Illumina, RS-122-2001, RS-122-2002) and quantified using real-time PCR. Agilent 2100 Bioanalyzer (Agilent) was used to assess the size distribution and to determine the RNA integrity number (RIN). All RNA samples for sequencing had RIN values greater or equal to 7 (Duan *et al.* 2018). The sequencing was performed on NextSeq 500 System (Illumina) with 75 bp paired-end reads in three sequencing runs. Overall, we obtained 2,149 million raw sequencing reads that passed filtering from three sequencing runs of 45 fetal tissue samples. The raw read dataset has been uploaded to GEO database with the accession number GSE111306.

Additional RNA-seq datasets

In addition to the RNA-seq data described above, two additional RNA-seq datasets were downloaded from Sequence Read Archive (SRA; <http://www.ncbi.nlm.nih.gov/sra>) under the accession numbers PRJNA254105 (Brooks *et al.* 2015) and PRJEB6169 (Jiang *et al.* 2014). PRJNA254105 included whole conceptuses at day 14 of gestation. PRJEB6169 contained data from adult and juvenile (6-10 months) heart, brain, liver, biceps femoris, rumen, female and male specific tissues, including the cervix, ovarian follicles, ovary, uterus, corpus luteum, testes, and placenta and membranes.

RNA-seq data trimming, mapping and assembly

Sequence adapter and quality trimming were conducted using Sickel v1.33 (Joshi and Fass 2011) with the parameters Q score ≥ 30 and length ≥ 20 ($-q30, -l20$). RNA-seq reads were checked using FastQC v0.11.3 (Andrews 2010) for quality control. Filtered RNA-seq reads from fetal tissues of day 135 of gestation were aligned to the sheep reference genome Oar_v4.0 using Hisat2 v2.0.5 (Kim *et al.* 2015). The mapping rates of all datasets are summarized in Table S1. The average mapping rate for our data are 90% with 19,846,496 reads mapped to the genome, whereas the additional datasets had an averaged 75% mapping rate with 12,854,507 reads aligned.

Aligned reads for each tissue from all three datasets were assembled using IsoEM v1.1.4 (Nicolae *et al.* 2011). The mRNA level of each gene was estimated by \log_2 -transformed transcripts per kilobase million (TPM) within each dataset and quantified using IsoEM (version 1.1.4; Nicolae *et al.*, 2011). TPM normalizes for gene length first and then for sequencing depth. This was preferred to RPKM/FPKM because it normalizes among transcriptome sizes of different samples and allows more appropriate comparisons of gene expression across samples (Soneson *et al.* 2015). Gene expression levels in TPM were \log_2 -transformed to minimize variations. Expressed genes were defined as

TPM ≥ 1 (Clark *et al.* 2017). A total of 7,166 genes were expressed among all tissues and defined as “dosage sensitive” genes (Sangrithi *et al.* 2017). Genes in the pseudoautosomal regions (PARs) of the sex chromosomes were obtained from Ruminant PARs annotation by Raudsepp and Chowdhary (2015).

Dosage compensation calculation

A total of 20,519 genes are in the sheep genome and assigned to each chromosome. The X:A ratio was calculated as the Relative X Expression (RXE); the difference between the \log_2 -transformed mean TPM values of the X chromosome and autosomes (A), using the formula below, where X-linked and autosomal genes were expressed as x and a , respectively:

$$RXE = \log_2 \left(\frac{x}{a} \right) = \log_2 x - \log_2 a$$

An RXE ≥ 0 represents a full up-regulation of X. An RXE between 0 and -1 indicates partial X chromosome upregulation. An RXE of -1 indicates a lack of X chromosome upregulation.

We also calculated the relative expression of each autosome pair (RGE) over all other chromosomes (excluding mitochondria and the Y chromosome which are not annotated in sheep). The RGE value was used to determine if the expression of the X chromosome deviated from the normal range of expression by the autosomes and if a particular autosome pair is more/less expressed than the rest of the chromosomes. The RGE was calculated using the following formula:

$$RGE = \log_2 \left(\frac{a_i}{a_{n-i}} \right) = \log_2 a_i - \log_2 a_{n-i}$$

Where i represents a particular pair of autosomes, n represents all autosomes. $n-i$ represents all autosomes excluding the autosome i . If the RGE of an autosome pair was greater than or equal to 0, it represents upregulation of that autosome pair. An RGE between 0 and -1 indicates downregulation.

The boxplots of RXE and RGE were generated in R (R Development Core Team 2008) using ggplot2 package (Wickham 2009). In these plots the lower and upper hinges encompass the 25th and 75th percentile of the data. The distance between the hinges is the interquartile range (IQR). The lower and upper whiskers extending from the hinges represent values no further than 1.5-fold of the IQR or within 95% confidence interval. Outliers were plotted individually beyond the end of whiskers (McGill *et al.* 1978) and labeled with numbers of the corresponding chromosomes.

Differentially expressed X-linked genes across maternal nutrition

Differentially expressed genes (DEGs) between Con and Over or Con and Res were determined using IsoDE version 2 (Al Seesi *et al.* 2014; Mandric *et al.* 2017). IsoDE2 is based on 200 bootstrap replicates where sampling from the original data were performed with replacement and stratified by the group variables (Al Seesi *et al.* 2014). Bootstrapping (Efron and Tibshirani 1994) is advantageous because it offers a reliable solution to the lack/low replicates and allows distinction between biological differences and technical variability or noise. Bootstrapping method is simple to apply and does not require any distribution assumptions. In each comparison, a gene was deemed differentially expressed if it showed \log_2 fold change (FC) > 1 between two treatments and significantly different

(P-value ≤ 0.05). The X-linked DEGs (Table S2) were a sub-group of the total DEGs from all chromosomes.

Gene ontology analysis

A Gene Ontology (GO) classification was conducted using DAVID 6.8 (Huang *et al.* 2009b, 2009a). GO categories with P-value ≤ 0.05 were considered significantly overrepresented. The pie plot of “dosage sensitive” genes categorized by protein functions was made in PANTHER classification system (Mi *et al.* 2013).

Data Availability Statement

The datasets analyzed during the current study are available in the Gene Expression Omnibus and BioProject:

GSE111306: <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE111306>

PRJNA254105: <https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA254105>

PRJEB6169: <https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJEB6169>

Supplemental material available at Figshare: <https://doi.org/10.25387/g3.7221467>.

RESULTS

X chromosome upregulation in ovine major organs and reproductive tissues

We found that the liver, muscle, rumen, heart of juveniles and adults, conceptuses and placenta and membrane at day 14 of gestation displayed partial X chromosome upregulation with RXE values ranging from -0.19 to -0.05, and an overall average RXE of -0.12 (Figure 1A). Interestingly, all RXE were much closer to 0 than to -1, indicating a substantial amount of dosage compensation across the entire X chromosome. The upregulation, however, appeared to be more pronounced in the brain. The RXE ranged from -0.12 to 0.16 in the cerebrum, cerebellum, hypothalamus, and pituitary (Figure 1B), suggesting complete X chromosome upregulation with the exception of the cerebellum. No significant difference (P-value = 0.74, by student *t*-test) was observed between males and females in the same tissue, demonstrating that X chromosome upregulation occurred similarly in both sexes, despite of the difference in the number of X chromosomes. Moreover, the RXE values fell within 1.5 times of the interquartile ranges (25–75% of the data) of RGEs of autosome pairs for all examined tissues (Figure 1A and 1B), suggesting the single active X chromosome in somatic tissues balanced its gene transcription outputs with those of the autosome pairs.

Partial X chromosome upregulation was also observed in juvenile and adult female reproduction-related tissues. These included the cervix, ovarian follicle, ovary, uterus, and corpus luteum. The overall averaged RXE was -0.19 and -0.15 for juvenile and adult female tissues, respectively (Figure 2). Among these, juvenile follicles, adult ovaries and corpora lutea had the greatest RXE values. However, a different pattern was observed in the male specific reproductive tissues studied. The testes exhibited an average RXE of -0.84, or a near lack of X chromosome upregulation (*i.e.*, RXE = -1; Figure 2).

We also observed that a number of autosome pairs had either greater than or less than the overall averaged gene expression. Chromosome 14, for example, was very “quiet” in gene expression at the chromosomal level, falling outside of the 1.5 interquartile ranges of RGEs of the other autosome pairs in many tissues (Figure 1A). Conversely, Chromosomes 20 and 25 were “active” in many tissues with high RGE values (greater than 0.5). Interestingly, the expression activity was negatively




Figure 1 Boxplots of \log_2 -transformed relative expression of the X chromosome (RXE) and each autosome pair (RGE) in major tissues (A) and brain (B) of juvenile and adult sheep. Red dots: mean RXEs for all replicates within a tissue type. Black dots: mean RGEs for each autosome pair. Numbers by black dots: autosomes whose RGEs fell outside of the expression quartiles for the tissue. Red dotted line: the border for complete (above line) and incomplete (below line) dosage compensation. The X:A ratio was calculated as the Relative X expression, $RXE = \log_2(X) - \log_2(A)$, the difference between the \log_2 -transformed mean TPM values of X and A. An RXE value of 0 means the expression of X and autosome is equal, suggesting X dosage compensation. Positive and negative RXE values indicate complete and incomplete dosage compensation, respectively. An RXE of -1, however, represents the lack of X dosage compensation. RGE of each autosome pair over all other chromosomes was used to evaluate the deviation of X expression to autosomes.

correlated ($r = -0.94$) with the numbers of expressed genes ($TPM \geq 1$) on these chromosomes. With an averaged 751 expressed genes, Chromosome 14 was less active (average $RGE = -0.44$) than Chromosomes 20 (average $RGE = 0.22$) and 25 (average $RGE = 0.25$) with 416 and 202 expressed genes, respectively. Rather, expression activity of these chromosomes may be related to the functions of genes that they contain. We therefore analyzed the gene ontology (GO) terms of lowly ($1 \leq TPM < 50$) expressed genes on Chromosomes 14, 20, and 25. The GO terms for Chromosome 14 included regulation of DNA-templated transcription, which corresponds to the reduced expression of transcription factors in most of tissues (Vaquerizas *et al.* 2009). On the other hand, the major GO

categories of highly expressed genes ($TPM > 100$) on Chromosomes 20 and 25 were enriched in nucleosome assembly and sarcomere organization (Table S3). These terms corresponded to greater activities of Chromosome 20 in early conceptuses and Chromosome 25 in biceps muscles (Figure 1A).

X chromosome upregulation in ovine fetuses under different maternal nutrition

X chromosome upregulation in ovine fetuses at day 135 of gestation was not affected by maternal diet, either over- or restricted-nutrition (P -value = 0.59, 0.70, respectively, by Wilcoxon Rank Sums tests). Tissues (brain, kidney, and lung) of both male and female fetuses from




Figure 2 Boxplots of \log_2 -transformed relative expression of the X chromosome (RXE) and each autosome pair (RGE) in female- and male-specific tissues. Red dots: RXEs for all replicates within a tissue type. Black dots: RGEs for each autosome pair. Numbers by black dots: autosomes whose RGEs fell outside of the expression quartiles for the tissue. Red dotted line: the border for complete (above line) and incomplete (below line) dosage compensation. The X:A ratio was calculated as the Relative X expression, $RXE = \log_2(X) - \log_2(A)$, the difference between the \log_2 -transformed mean TPM values of X and A. An RXE value of 0 means the expression of X and autosome is equal, suggesting X dosage compensation. Positive and negative RXE values indicate complete and incomplete dosage compensation, respectively. An RXE of -1, however, represents the lack of X dosage compensation. RGE of each autosome pair over all other chromosomes was used to evaluate the deviation of X expression to autosomes.

complete dosage compensation, respectively. An RXE of -1, however, represents the lack of X dosage compensation over all other chromosomes was used to evaluate the deviation of X expression to autosomes.

mothers of all three treatment groups displayed partial dosage compensation. The RXEs of the three tissues ranged from -0.13 to -0.05, (Figure 3A), from -0.14 to -0.05 (Figure 3B), and from -0.16 to -0.08 (Figure 3C) for the Con-, Over- and Res-fed group, respectively. All RXE values fell within the RGE ranges, suggesting the single X upregulated on its expression to levels close to the autosome pairs. These observations indicate that different maternal nutrition did not affect the upregulation of the X chromosome in fetal tissues.

X chromosome upregulation in different gene subgroups

We calculated the RXE values in the gene categories of “All genes”, “Expressed genes”, “Genes subject to XCI (removal of PAR genes)” and “Dosage sensitive genes” (Figure 4). “All genes” included low- and non-expressed genes ($TPM < 1$), and had the lowest RXE values among the four subgroups of genes. The median of RXE in the “All genes” category was close to -0.5, indicating that when all X-linked genes were




Figure 3 Boxplots of \log_2 -transformed expression of the X chromosome (RXE) and each autosome pair (RGE) by fetal tissues from mothers under different nutritional treatments: Control (A), Overfed (B) and Restricted (C). Red dots: RXEs for all replicates within a treatment group. Black dots: RGEs for each autosome pair. Numbers by black dots: autosomes whose RGEs fell outside of the expression quartiles for the tissue. Red dotted line: the border for complete (above line) and incomplete (below line) dosage compensation. The X:A ratio was calculated as the Relative X expression, $RXE = \log_2(X) - \log_2(A)$, the difference between the \log_2 -transformed mean TPM values of X and A. An RXE value of 0 means the expression of X and autosome is equal, suggesting X dosage compensation. Positive and negative RXE values indicate complete and incomplete dosage compensation, respectively. An RXE of -1, however, represents the lack of X dosage compensation. RGE of each autosome pair over all other chromosomes was used to evaluate the deviation of X expression to autosomes.

considered (including those with leaky expression), there was nearly no upregulation of the X chromosome. “Expressed genes” gave a partial X chromosome upregulation with a median RXE value close to 0, suggesting that this group contained genes of the X chromosome that were not subjected to X dosage compensation. Therefore, we removed the 14 genes located in ovine PAR. These genes have a homologous copy on the Y chromosome, and are not subjected to XCI. The RXE values without PAR were slightly increased, indicating that the PAR genes had lower expression. Moreover, we characterized another category - “Dosage sensitive genes”, which were ubiquitously expressed across all samples in our study and were mostly housing-keeping genes such as those involved in nucleic acid binding, cytoskeletal proteins and transferase (Figure S1 and Table S4). These “Dosage sensitive genes” had the highest median RXEs of greater than 0, corresponding to a full X upregulation. Taken together, our analysis of the four gene categories suggests that dosage regulation is highly related to gene functions.

Effects of maternal nutrition on the expression of X-linked genes in ovine fetal tissues

A total of 1,228 X-linked genes were annotated in the current ovine genome (Jiang *et al.* 2014). Among these, 625 genes were expressed (TPM ≥ 1) by the three fetal tissues combined (Table S2). The mean number of expressed X-linked genes by each fetal tissue was calculated for each maternal nutrition group (Table 1). On average, 518.2 ± 14.7 out of 625 X-linked genes were expressed in the three organs. Specifically, the brain expressed the most genes (536.1 ± 14.3 ; control group), followed by the kidney (517.3 ± 9.2) and the lung expressed the fewest (506 ± 3.7). The numbers of the expressed X-linked genes were not significantly different (P-value > 0.05 , by one-way ANOVA) across the maternal nutrition treatments (Table 1). However, the levels of expression of the X-linked genes were affected by maternal nutrition. A total of 57 X-linked genes were differentially expressed among treatment groups. For example, two genes related to sex determination- *SOX3* and *NROB1*-were down-regulated in fetal brains of the Over group (Table S5). The changes in sex-linked genes may provide a mechanism for the highly debated observation that skewed sex ratio was related to maternal nutrition (Mathews *et al.* 2008). The top eight X-linked DEGs (*PAGE4*, *S100G*, *SOX3*, *KCNE5*, *CLDN2*, *DUSP21*, *LOC105610402*, and *SLC6A14*) were summarized in Table 2 and were all expressed 8X ($> 3 \log_2$ -Fold Change) more than that of the controls. Taken together, these expression data clearly demonstrated an effect of poor maternal nutrition on gene expression during fetal development.

DISCUSSION

To our knowledge, this is the first study of X chromosome compensatory expression upregulation in sheep. We conclude that X chromosome upregulation was present, but largely partial. Additionally, X chromosome upregulation in fetal organs was not affected by the different maternal diets. While a number of species, both invertebrates and vertebrates, have been examined for their X:A ratios, whether X expression is globally upregulated is still highly debated [reviewed in (Gu and Walters 2017)]. Recent studies in therian mammals, including the human, mouse, bovine, and non-human primates mostly support the partial X chromosome upregulation conclusion with X:A ratio being close to 1 (Gu and Walters 2017; Duan *et al.*, 2016; Ka *et al.*, 2016). Our findings here contribute to the consensus of partial X chromosome upregulation in a new species.




Figure 4 Boxplot of RXE values in the categories of “All genes”, “Expressed genes”, “Genes subject to XCI (removal of genes in PAR)” and “Dosage sensitive genes”. Red dotted line: the border for complete (above line) and incomplete (below line) dosage compensation.

The estimation of X:A ratios differs when different gene subgroups and different tissues are analyzed, thus resulting in completely different conclusions over Ohno’s hypothesis (Sangrithi and Turner 2018). Some of the low- and non-expressed genes in somatic tissues were found to be highly expressed in testis. These genes are more enriched on the X chromosome than on autosomes (Rice 1996; Deng *et al.* 2011; Disteche 2016). Therefore, when the analysis included low- and non-expressed genes, the estimation of X chromosome upregulation is biased. Our result showed that RXE was closer to -0.5 when “all genes” were included, while RXE was close to 0 when only expressed genes were used. These two different types of gene categorization and inclusion corresponded to the opposite findings by Xiong *et al.* (2010) and Deng *et al.* (2011). Additionally, dosage compensation requires both X chromosome upregulation and XCI. Not all genes on the inactive X, however, are silenced. A group of X-linked genes escape inactivation (Disteche *et al.* 2002; Berletch *et al.* 2011; Al Nadaf *et al.* 2012; Balaton and Brown 2016). These include all genes in PARs (Helena Mangs and Morris 2007) and a few in non-PAR regions of X (Tukiainen *et al.* 2017). As the homologous region of the mammalian sex chromosomes, genes on PARs are expressed from both the X and Y chromosomes (Vermeesch *et al.* 1997). However, we are not able to exclude any

■ **Table 1** Mean numbers of expressed (TPM ≥ 1) X-linked genes in tissues of day 135 fetuses from control (n = 7), overfed (n = 4) and restricted (n = 4) mothers

	Treatments			P-value
	Control	Overfed	Restricted	
Brain	536.1 \pm 15.4	531.8 \pm 8.7	530.3 \pm 9.2	0.73
Kidney	517.3 \pm 9.9	514.5 \pm 9.1	518.8 \pm 2.2	0.77
Lung	506.0 \pm 4.0	503.3 \pm 7.4	502.0 \pm 4.2	0.44

non-PAR genes from the group “subject to XCI” due to the lack of information on non-PAR genes that escape XCI from these regions in the ovine, we were only able to exclude PAR genes in the group of “Subject to XCI”. Of the 20 annotated genes in ovine PAR (Figure S2) (Raudsepp and Chowdhary 2015), 14 were expressed in our study. They were *P2RY8*, *DHRX*, *ZBED1*, *CD99*, *XG*, *GYG2*, *ARSE*, *MXRA5*, *PRKX*, *NLGN4X*, *STS*, *PNPLA4*, *TBL1X*, and *GPR143*. They had relatively low expression levels, ranging in TPM from 1–50 while the average expression level of X-linked genes was 78.5 in TPM. Not much change in RXE values was found when the PAR genes were removed from the expressed group; possibly due their small number. Furthermore, our analysis showed a full compensatory upregulation of “dosage-sensitive” X-linked genes in sheep. This is in agreement with previous findings in the mouse and human by Ramsköld *et al.* (2009) and Sangrithi *et al.* (2017) who suggested that ubiquitous gene expression corresponded to the housekeeping function of dosage sensitive genes. It is likely that in order not to create limiting effects, gene products of this subgroup must be generated at comparable levels to those of the same pathways yet encoded by autosome pairs. Therefore, ubiquitously expressed genes are much more upregulated compared to other genes on the X chromosome.

There are a few exceptions to the general finding that ovine tissues underwent partial X chromosome upregulation. One exception is the brain, which had the greatest overall RXE values among all somatic tissues (RXE ranged from -0.12 to 0.16). This greater degree of X chromosome upregulation has also been observed in other species, including the human, mouse (Nguyen and Disteche 2006), old world monkeys, opossum, platypus, and chicken (Julien *et al.* 2012). The higher X chromosome upregulation is likely the result of both greater levels as well as numbers of expression of X-linked genes in the brain (Table 1). During evolution, the X chromosome accumulated an excess of sex- and reproduction-related genes (Saifi and

Chandra 1999). Greater expression of X-linked genes in the brain has been described as “the large X-chromosome effect” (Wu and Davis 1993), which was hypothesized to influence general cognitive ability, female mating choices and contribute to species diversification (Zechner *et al.* 2001). Therefore, it is expected that the brain would have a higher RXE.

Another exception to the overall X chromosome expression upregulation was seen in the sheep male reproduction-related tissues. The RXE was extremely low in sheep testes, corresponding to the observation of low X:A ratio in both the testes and spermatids in mice, indicating an X-specific partial repression in these cells (Nguyen and Disteche 2006). It was reported that the X:A ratio remained low in spermatogonia (Nguyen and Disteche 2006; Sangrithi and Turner 2018). Subsequently both the X and Y chromosomes become inactivated by meiotic sex chromosome inactivation during spermatogenesis (Manterola *et al.* 2009). This suppression of the X chromosome is likely the cause for the low X:A ratio.

Day 14 whole embryos, on the other hand, had an RXE value of -0.05 which was very close to full dosage compensation (RXE = 0). High X chromosome upregulation in early embryos could be a rebound after the release of repression of sex chromosomes in sperm. In the early embryos this release is necessary for X upregulation initiation (Wang *et al.* 2016). The expression of X chromosome was reported to be upregulated after the blastocyst stage which continued during 6.5 to 10.5 days post coitum development in mice (Nguyen and Disteche 2006). Mouse embryonic stem cells from both XX and XY embryos were also found to undergo X upregulation (Lin *et al.* 2011). Although XCI is known to operate in sheep fetuses (Luciani *et al.* 1979), little is known about its onset and regulation. In the bovine conceptuses, the onset of random XCI is found to have been established before day 14 (Bermejo-Alvarez *et al.* 2011). In the ovine, it is very likely that XCI has occurred by day 14 due to its shorter gestation (King *et al.* 1985; Stevens *et al.* 1990). It is therefore highly possible that the Day 14 ovine conceptuses had only one active X chromosome. The greater RXE value in Day 14 conceptuses thus may imply that the single active X chromosome just started its compensation process. This is consistent with the greater RXE values observed in Day 10–19 conceptuses in the bovine (Duan *et al.* 2017).

In summary, our comprehensive analyses of X chromosome dosage compensation suggest upregulation of gene expression from the single active X chromosome in most ovine tissues of both sexes.

■ **Table 2** Differentially expressed X-linked genes by tissues of ovine fetuses from mothers under control, overfed and restricted nutrition treatments

Comparison	Tissue	Gene	Expression in controls (TPM)	Expression in treated (TPM)	Log ₂ FC*
Con vs. Over	Brain	<i>PAGE4</i>	99.80	0.18	-∞
	Brain	<i>S100G</i>	33.62	0.46	-∞
	Brain	<i>SOX3</i>	1.23	0.15	-∞
	Kidney	<i>KCNE5</i>	1.71	0.27	-∞
	Kidney	<i>PAGE4</i>	1.46	0.86	-3.23
	Lung	<i>CLDN2</i>	0.26	6.36	4.71
Con vs. Res	Brain	<i>DUSP21</i>	0.08	1.04	5.21
	Brain	<i>LOC105610402</i>	1.04	0.00	-∞
	Brain	<i>S100G</i>	33.62	1.27	-5.20
	Lung	<i>SLC6A14</i>	1.57	0.10	-5.43

*Log₂ FC: calculated by using bootstrapping; FC: fold change.

∞: infinity;

PAGE4: PAGE family member 4; *S100G*: S100 calcium binding protein G; *SOX3*: SRY-box 3; *KCNE5*: potassium voltage-gated channel subfamily E regulatory subunit 5; *CLDN2*: claudin 2; *DUSP21*: dual specificity phosphatase 21; *LOC105610402*: 60S ribosomal protein L17; *SLC6A14*: solute carrier family 6 member 14.

ACKNOWLEDGMENTS

The authors thank Zoetis for donating the controlled intravaginal drug release devices (CIDRs) used for estrus synchronization and the UConn Livestock staff, Dr. Thomas Hoagland, Victor Delaire and the animal science undergraduate students for the animal care during this experiment. This work was supported by the USDA-ARS grant: 1265-31000-091-02S, the USDA Multi-state regional grant: W3171, National Institute of Food and Agriculture, U.S. Department of Agriculture, under award number 2013-01919, Department of Education of Xinjiang Uygur Autonomous Region Scholarship: 2016-3-0036 and Studying Abroad program for Excellent Ph.D. Students of Guangxi Zhuang Autonomous Region: 2014-2. XCT, JD, KF, MZ and ZJ designed the study; AJ, SP, MH, SR, KG and SZ performed animal breeding, feeding and care, necropsy, and tissue sample collection; KF, MZ and HJ performed the RNA-seq experiment; JD, NJ, SA, IM and RO analyzed the data; KF, JD and XCT wrote the manuscript, NJ, AJ, RO, SZ, and KG edited the manuscript. All authors read and approved the final manuscript.

LITERATURE CITED

- Al Nadaf, S., J. E. Deakin, C. Gilbert, T. J. Robinson, J. A. M. Graves *et al.*, 2012 A cross-species comparison of escape from X inactivation in Eutheria: implications for evolution of X chromosome inactivation. *Chromosoma* 121: 71–78. <https://doi.org/10.1007/s00412-011-0343-8>
- Al Seesi, S., Y. T. Tiague, A. Zelikovsky, and I. I. Mandoiu, 2014 Bootstrap-based differential gene expression analysis for RNA-Seq data with and without replicates. *BMC Genomics* 15: S2. <https://doi.org/10.1186/1471-2164-15-S8-S2>
- Andrews, S., 2010 FastQC A Quality Control tool for High Throughput Sequence Data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- Balaton, B. P., and C. J. Brown, 2016 Escape Artists of the X Chromosome. *Trends Genet.* TIG 32: 348–359. <https://doi.org/10.1016/j.tig.2016.03.007>
- Barry, J. S., and R. V. Anthony, 2008 The Pregnant Sheep as a Model for Human Pregnancy. *Theriogenology* 69: 55–67. <https://doi.org/10.1016/j.theriogenology.2007.09.021>
- Begum, G., A. Stevens, E. B. Smith, K. Connor, J. R. G. Challis *et al.*, 2012 Epigenetic changes in fetal hypothalamic energy regulating pathways are associated with maternal undernutrition and twinning. *FASEB J.* 26: 1694–1703. <https://doi.org/10.1096/fj.11-198762>
- Berlitch, J. B., F. Yang, J. Xu, L. Carrel, and C. M. Disteche, 2011 Genes that escape from X inactivation. *Hum. Genet.* 130: 237–245. <https://doi.org/10.1007/s00439-011-1011-z>
- Bermejo-Alvarez, P., D. Rizo, P. Lonergan, and A. Gutierrez-Adan, 2011 Transcriptional sexual dimorphism in elongating bovine embryos: implications for XCI and sex determination genes. *Reproduction* 141: 801–808. <https://doi.org/10.1530/REP-11-0006>
- Birchler, J. A., 2012 Claims and counterclaims of X-chromosome compensation. *Nat. Struct. Mol. Biol.* 19: 3–5. <https://doi.org/10.1038/nsmb.2218>
- Birchler, J. A., N. C. Riddle, D. L. Auger, and R. A. Veitia, 2005 Dosage balance in gene regulation: biological implications. *Trends Genet.* 21: 219–226. <https://doi.org/10.1016/j.tig.2005.02.010>
- Brooks, K. E., G. W. Burns, and T. E. Spencer, 2015 Peroxisome proliferator activator receptor gamma (PPARG) regulates conceptus elongation in sheep. *Biol. Reprod.* 92: 42. <https://doi.org/10.1095/biolreprod.114.123877>
- Burgess, T., L. Downie, M. D. Pertile, D. Francis, M. Glass *et al.*, 2014 Monosomy 21 Seen in Live Born Is Unlikely to Represent True Monosomy 21: A Case Report and Review of the Literature. *Case Rep. Genet.* 2014: 965401.
- Chen, Z.-X., K. Golovkina, H. Sultana, S. Kumar, and B. Oliver, 2014 Transcriptional effects of gene dose reduction. *Biol. Sex Differ.* 5: <https://doi.org/10.1186/2042-6410-5-5>
- Clark, E. L., S. J. Bush, M. E. B. McCulloch, I. L. Farquhar, R. Young *et al.*, 2017 A high resolution atlas of gene expression in the domestic sheep (*Ovis aries*). *PLoS Genet.* 13: e1006997. <https://doi.org/10.1371/journal.pgen.1006997>
- Deakin, J. E., J. Chaumeil, T. A. Hore, and J. A. M. Graves, 2009 Unravelling the evolutionary origins of X chromosome inactivation in mammals: insights from marsupials and monotremes. *Chromosome Res.* 17: 671–685. <https://doi.org/10.1007/s10577-009-9058-6>
- Deng, X., J. B. Hiatt, D. K. Nguyen, S. Ercan, D. Sturgill *et al.*, 2011 Evidence for compensatory upregulation of expressed X-linked genes in mammals, *Caenorhabditis elegans* and *Drosophila melanogaster*. *Nat. Genet.* 43: 1179–1185. <https://doi.org/10.1038/ng.948>
- Disteche, C. M., 2012 Dosage compensation of the sex chromosomes. *Annu. Rev. Genet.* 46: 537–560. <https://doi.org/10.1146/annurev-genet-110711-155454>
- Disteche, C. M., 2016 Dosage compensation of the sex chromosomes and autosomes. *Semin. Cell Dev. Biol.* 56: 9–18. <https://doi.org/10.1016/j.semcdb.2016.04.013>
- Disteche, C. M., G. N. Filippova, and K. D. Tsuchiya, 2002 Escape from X inactivation. *Cytogenet. Genome Res.* 99: 36–43. <https://doi.org/10.1159/000071572>
- Dominguez-Salas, P., S. E. Moore, M. S. Baker, A. W. Bergen, S. E. Cox *et al.*, 2014 Maternal nutrition at conception modulates DNA methylation of human metastable epialleles. *Nat. Commun.* 5: 3746. <https://doi.org/10.1038/ncomms4746>
- Du, M., J. X. Zhao, X. Yan, Y. Huang, L. V. Nicodemus *et al.*, 2011 Fetal muscle development, mesenchymal multipotent cell differentiation, and associated signaling pathways. *J. Anim. Sci.* 89: 583–590. <https://doi.org/10.2527/jas.2010-3386>
- Duan, J., N. K. Jue, Z. Jiang, R. O'Neill, E. Wolf *et al.*, 2017 125 incomplete compensatory up-regulation of x-linked genes in bovine germline, early embryos, and somatic tissues. *Reprod. Fertil. Dev.* 29: 171. <https://doi.org/10.1071/RDv29n1Ab125>
- Duan, J. E., N. K. Jue, Z. Jiang, R. O'Neill, E. Wolf *et al.*, 2016 144 dosage compensation and x-linked gene expression in bovine in vivo and in vitro embryos. *Reprod. Fertil. Dev.* 28: 202. <https://doi.org/10.1071/RDv28n2Ab144>
- Duan, J., M. Zhang, K. Flock, S. A. Seesi, and I. Mandoiu *et al.*, 2018 Effects of Maternal Nutrition on the Expression of Genomic Imprinted Genes in Ovine Fetuses. *Epigenetics* 13: 793–807. <https://doi.org/10.1080/15592294.2018.1503489>
- Efron, B., and R. J. Tibshirani, 1994 *An Introduction to the Bootstrap*, CRC Press, Boca Raton, FL.
- Giam, M., and G. Rancati, 2015 Aneuploidy and chromosomal instability in cancer: a jackpot to chaos. *Cell Div.* 10: 3. <https://doi.org/10.1186/s13008-015-0009-7>
- Goto, T., and M. Monk, 1998 Regulation of X-Chromosome Inactivation in Development in Mice and Humans. *Microbiol. Mol. Biol. Rev.* 62: 362–378.
- Graves, J. A. M., 2016 Did sex chromosome turnover promote divergence of the major mammal groups? *BioEssays* 38: 734–743. <https://doi.org/10.1002/bies.201600019>
- Gu, L., and J. R. Walters, 2017 Evolution of Sex Chromosome Dosage Compensation in Animals: A Beautiful Theory, Undermined by Facts and Bedeviled by Details. *Genome Biol. Evol.* 9: 2461–2476. <https://doi.org/10.1093/gbe/evx154>
- Gupta, V., M. Parisi, D. Sturgill, R. Nuttall, M. Doctolero *et al.*, 2006 Global analysis of X-chromosome dosage compensation. *J. Biol.* 5: 3. <https://doi.org/10.1186/jbiol30>
- Helena Mangs, A., and B. J. Morris, 2007 The Human Pseudoautosomal Region (PAR): Origin, Function and Future. *Curr. Genomics* 8: 129–136. <https://doi.org/10.2174/138920207780368141>
- Huang, D. W., B. T. Sherman, and R. A. Lempicki, 2009a Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* 37: 1–13. <https://doi.org/10.1093/nar/gkn923>
- Huang, D. W., B. T. Sherman, and R. A. Lempicki, 2009b Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.* 4: 44–57. <https://doi.org/10.1038/nprot.2008.211>

- Jiang, Y., M. Xie, W. Chen, R. Talbot, J. F. Maddox *et al.*, 2014 The Sheep Genome Illuminates Biology of the Rumen and Lipid Metabolism. *Science* 344: 1168–1173. <https://doi.org/10.1126/science.1252806>
- Joshi, N., and J. Fass, 2011 Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files.
- Julien, P., D. Brawand, M. Soumillon, A. Necsulea, A. Liechti *et al.*, 2012 Mechanisms and evolutionary patterns of mammalian and avian dosage compensation. *PLoS Biol.* 10: e1001328. <https://doi.org/10.1371/journal.pbio.1001328>
- Ka, S., H. Ahn, M. Seo, H. Kim, J. N. Kim *et al.*, 2016 Status of dosage compensation of X chromosome in bovine genome. *Genetica* 144: 435–444. <https://doi.org/10.1007/s10709-016-9912-3>
- Kim, D., B. Langmead, and S. L. Salzberg, 2015 HISAT: a fast spliced aligner with low memory requirements. *Nat. Methods* 12: 357–360. <https://doi.org/10.1038/nmeth.3317>
- King, K. K., G. E. Seidel, and R. P. Elsdon, 1985 Bovine Embryo Transfer Pregnancies. II. Lengths of Gestation. *J. Anim. Sci.* 61: 758–762. <https://doi.org/10.2527/jas1985.614758x>
- Kondrashov, F. A., and E. V. Koonin, 2004 A common framework for understanding the origin of genetic dominance and evolutionary fates of gene duplications. *Trends Genet.* 20: 287–290. <https://doi.org/10.1016/j.tig.2004.05.001>
- Lee, J. T., 2011 Gracefully ageing at 50, X-chromosome inactivation becomes a paradigm for RNA and chromatin control. *Nat. Rev. Mol. Cell Biol.* 12: 815–826. <https://doi.org/10.1038/nrm3231>
- Lin, H., V. Gupta, M. D. Vermilyea, F. Falciani, J. T. Lee *et al.*, 2007 Dosage compensation in the mouse balances up-regulation and silencing of X-linked genes. *PLoS Biol.* 5: e326. <https://doi.org/10.1371/journal.pbio.0050326>
- Lin, H., J. A. Halsall, P. Antczak, L. P. O'Neill, F. Falciani *et al.*, 2011 Relative overexpression of X-linked genes in mouse embryonic stem cells is consistent with Ohno's hypothesis. *Nat. Genet.* 43: 1169–1170, author reply 1171–1172. <https://doi.org/10.1038/ng.992>
- Lin, F., K. Xing, J. Zhang, and X. He, 2012 Expression reduction in mammalian X chromosome evolution refutes Ohno's hypothesis of dosage compensation. *Proc. Natl. Acad. Sci. USA* 109: 11752–11757. <https://doi.org/10.1073/pnas.1201816109>
- Livernois, A. M., J. M. Graves, and P. D. Waters, 2012 The origin and evolution of vertebrate sex chromosomes and dosage compensation. *Heredity* 108: 50–58. <https://doi.org/10.1038/hdy.2011.106>
- Luciani, J., J. Bézard, M. Devictor-Vuillet, and P. Mauléon, 1979 ³H-thymidine labelling pattern of preleptotene chromosome condensation stages in the fetal sheep ovary. *Ann. Biol. Anim. Biochim. Biophys.* 19: 1241–1250.
- Lyon, M. F., 1961 Gene Action in the X-chromosome of the Mouse (*Mus musculus* L.). *Nature* 190: 372–373. <https://doi.org/10.1038/190372a0>
- Mandric, I., Y. Temate-Tiagueu, T. Shcheglova, S. Al Seesi, A. Zelikovsky *et al.*, 2017 Fast bootstrapping-based estimation of confidence intervals of expression levels and differential expression from RNA-Seq data. *Bioinformatics* 33: 3302–3304. <https://doi.org/10.1093/bioinformatics/btx365>
- Mank, J. E., D. J. Hosken, and N. Wedell, 2014 Conflict on the Sex Chromosomes: Cause, Effect, and Complexity. *Cold Spring Harb. Perspect. Biol.* 6: a017715. <https://doi.org/10.1101/cshperspect.a017715>
- Manterola, M., J. Page, C. Vasco, S. Berrios, M. T. Parra *et al.*, 2009 A High Incidence of Meiotic Silencing of Unsynapsed Chromatin Is Not Associated with Substantial Pachytene Loss in Heterozygous Male Mice Carrying Multiple Simple Robertsonian Translocations. *PLoS Genet.* 5: e1000625. <https://doi.org/10.1371/journal.pgen.1000625>
- Mathews, F., P. J. Johnson, and A. Neil, 2008 You are what your mother eats: evidence for maternal preconception diet influencing foetal sex in humans. *Proc. Biol. Sci.* 275: 1661–1668. <https://doi.org/10.1098/rspb.2008.0105>
- McGill, R., J. W. Tukey, and W. A. Larsen, 1978 Variations of Box Plots. *Am. Stat.* 32: 12–16.
- Mi, H., A. Muruganujan, J. T. Casagrande, and P. D. Thomas, 2013 Large-scale gene function analysis with the PANTHER classification system. *Nat. Protoc.* 8: 1551–1566. <https://doi.org/10.1038/nprot.2013.092>
- Nguyen, D. K., and C. M. Disteché, 2006 Dosage compensation of the active X chromosome in mammals. *Nat. Genet.* 38: 47–53. <https://doi.org/10.1038/ng1705>
- Nicolae, M., S. Mangul, I. I. Măndoiu, and A. Zelikovsky, 2011 Estimation of alternative splicing isoform frequencies from RNA-Seq data. *Algorithms Mol. Biol.* 6: 9. <https://doi.org/10.1186/1748-7188-6-9>
- Ohno, S., 1966 *Sex Chromosomes and Sex-Linked Genes*, Springer-Verlag, Berlin, Heidelberg. <https://doi.org/10.1007/978-3-662-35113-0>
- Pessia, E., J. Engelstädter, and G. A. B. Marais, 2014 The evolution of X chromosome inactivation in mammals: the demise of Ohno's hypothesis? *Cell. Mol. Life Sci. CMLS* 71: 1383–1394. <https://doi.org/10.1007/s00018-013-1499-6>
- Pessia, E., T. Makino, M. Bailly-Bechet, A. McLysaght, and G. A. B. Marais, 2012 Mammalian X chromosome inactivation evolved as a dosage-compensation mechanism for dosage-sensitive genes on the X chromosome. *Proc. Natl. Acad. Sci. USA* 109: 5346–5351. <https://doi.org/10.1073/pnas.1116763109>
- Pillai, S. M., N. H. Sereda, M. L. Hoffman, E. V. Valley, T. D. Crenshaw *et al.*, 2016 Effects of Poor Maternal Nutrition during Gestation on Bone Development and Mesenchymal Stem Cell Activity in Offspring. *PLoS One* 11: e0168382. <https://doi.org/10.1371/journal.pone.0168382>
- Pillai, S. M., A. K. Jones, M. L. Hoffman, K. K. McFadden, S. A. Reed *et al.*, 2017 Fetal and organ development at gestational days 45, 90, 135 and at birth of lambs exposed to under- or over-nutrition during gestation. *Trans. Anim. Sci.* 1: 16–25.
- R Development Core Team, 2008 R: A language and environment for statistical computing.
- Ramsköld, D., E. T. Wang, C. B. Burge, and R. Sandberg, 2009 An Abundance of Ubiquitously Expressed Genes Revealed by Tissue Transcriptome Sequence Data. *PLOS Comput. Biol.* 5: e1000598. <https://doi.org/10.1371/journal.pcbi.1000598>
- Raudsepp, T., and B. P. Chowdhary, 2015 The Eutherian Pseudoautosomal Region. *Cytogenet. Genome Res.* 147: 81–94. <https://doi.org/10.1159/000443157>
- Rice, W. R., 1996 Evolution of the Y Sex Chromosome in Animals. *BioScience* 46: 331–343. <https://doi.org/10.2307/1312947>
- Sahakyan, A., Y. Yang, and K. Plath, 2018 The Role of Xist in X-Chromosome Dosage Compensation. *Trends Cell Biol.* S0962–8924: 30100–4.
- Saifi, G. M., and H. S. Chandra, 1999 An apparent excess of sex- and reproduction-related genes on the human X chromosome. *Proc. Biol. Sci.* 266: 203–209. <https://doi.org/10.1098/rspb.1999.0623>
- Sangrithi, M. N., H. Royo, S. K. Mahadevaiah, O. Ojarikre, L. Bhaw *et al.*, 2017 Non-Canonical and Sexually Dimorphic X Dosage Compensation States in the Mouse and Human Germline. *Dev. Cell* 40: 289–301.e3. <https://doi.org/10.1016/j.devcel.2016.12.023>
- Sangrithi, M. N., and J. M. A. Turner, 2018 Mammalian X Chromosome Dosage Compensation: Perspectives From the Germ Line. *BioEssays News Rev. Mol. Cell. Dev. Biol.* 40: e1800024.
- Soneson, C., M. I. Love, and M. D. Robinson, 2015 Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. *F1000 Res.* 4: 1521. <https://doi.org/10.12688/f1000research.7563.1>
- Stevens, D., G. Alexander, and A. W. Bell, 1990 Effect of prolonged glucose infusion into fetal sheep on body growth, fat deposition and gestation length. *J. Dev. Physiol.* 13: 277–281.
- Tukiainen, T., A.-C. Villani, A. Yen, M. A. Rivas, J. L. Marshall *et al.*, 2017 Landscape of X chromosome inactivation across human tissues. *Nature* 550: 244–248. <https://doi.org/10.1038/nature24265>
- Vaquerizas, J. M., S. K. Kummerfeld, S. A. Teichmann, and N. M. Luscombe, 2009 A census of human transcription factors: function, expression and evolution. *Nat. Rev. Genet.* 10: 252–263. <https://doi.org/10.1038/nrg2538>
- Veitia, R. A., F. Veyrunes, S. Bottani, and J. A. Birchler, 2015 X chromosome inactivation and active X upregulation in therian mammals: facts, questions, and hypotheses. *J. Mol. Cell Biol.* 7: 2–11. <https://doi.org/10.1093/jmcb/mjv001>
- Vermeesch, J. R., P. Petit, A. Kermouni, J.-C. Renaud, H. Van Den Berghe *et al.*, 1997 The IL-9 Receptor Gene, Located in the Xq/Yq Pseudoautosomal Region, Has an Autosomal Origin, Escapes X Inactivation and Is Expressed from the Y. *Hum. Mol. Genet.* 6: 1–8. <https://doi.org/10.1093/hmg/6.1.1>

- Wang, F., J. Shin, J. M. Shea, J. Yu, A. Bošković *et al.*, 2016 Regulation of X-linked gene expression during early mouse development by Rlim. *eLife* 5. <https://doi.org/10.7554/eLife.19127>
- Wickham, H., 2009 Ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York.
- Wolf, J. B., and J. Bryk, 2011 General lack of global dosage compensation in ZZ/ZW systems? Broadening the perspective with RNA-seq. *BMC Genomics* 12: 91. <https://doi.org/10.1186/1471-2164-12-91>
- Wu, C. I., and A. W. Davis, 1993 Evolution of postmating reproductive isolation: the composite nature of Haldane's rule and its genetic bases. *Am. Nat.* 142: 187–212. <https://doi.org/10.1086/285534>
- Xiong, Y., X. Chen, Z. Chen, X. Wang, S. Shi *et al.*, 2010 RNA sequencing shows no dosage compensation of the active X-chromosome. *Nat. Genet.* 42: 1043–1047. <https://doi.org/10.1038/ng.711>
- Xue, F., X. C. Tian, F. Du, C. Kubota, M. Taneja *et al.*, 2002 Aberrant patterns of X chromosome inactivation in bovine clones. *Nat. Genet.* 31: 216–220. <https://doi.org/10.1038/ng900>
- Zechner, U., M. Wilda, H. Kehrer-Sawatzki, W. Vogel, R. Fundele *et al.*, 2001 A high density of X-linked genes for general cognitive ability: a run-away process shaping human evolution? *Trends Genet. TIG* 17: 697–701. [https://doi.org/10.1016/S0168-9525\(01\)02446-5](https://doi.org/10.1016/S0168-9525(01)02446-5)

Communicating editor: J. Birchler