5-1-2017

Mitochondrial dysfunction and ovarian aging

Tianren Wang
Yale School of Medicine

Man Zhang
Yale School of Medicine

Zongliang Jiang
Yale School of Medicine

Emre Seli
Yale School of Medicine

Follow this and additional works at: https://digitalcommons.lsu.edu/animalsciences_pubs

Recommended Citation

Wang, T., Zhang, M., Jiang, Z., & Seli, E. (2017). Mitochondrial dysfunction and ovarian aging. *American Journal of Reproductive Immunology*, 77 (5) <https://doi.org/10.1111/aji.12651>

This Conference Proceeding is brought to you for free and open access by the School of Animal Sciences at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Mitochondrial dysfunction and ovarian aging

Tianren Wang | Man Zhang | Zongliang Jiang | Emre Seli

Department of Obstetrics, Gynecology and Reproductive Sciences, Yale School of Medicine, New Haven, CT, USA

Correspondence

Emre Seli, Department of Obstetrics, Gynecology and Reproductive Sciences, Yale University School of Medicine, New Haven, CT, USA.
Email: emre.seli@yale.edu

Funding information

National Institute of Health, Grant/Award Number: R01HD059909.

Mitochondria are double-membrane-bound organelles that are responsible for the generation of most of the cell's energy. Mitochondrial dysfunction has been implicated in cellular senescence in general and ovarian aging in particular. Recent studies exploited this association by studying mitochondrial DNA (mtDNA) copy number as a potential biomarker of embryo viability and the use of mitochondrial nutrients and autologous mitochondrial transfer as a potential treatment for poor ovarian function and response.

KEYWORDS

aging, mitochondria, ovary, reproduction

1 | INTRODUCTION

Twentieth century witnessed significant changes in the socioeconomic parameters that define the role of women in the society. In 1900, <20% of women participated in the labor force in the United States. This number increased to 33% by 1950 and 60% by 2000.

In parallel with women's increasing participation in the workforce and realization that college education increases employment and earning potential, college graduation rate that was less than 4% for women in the United States in 1940 reached 32.5% in 2015, equaling that of men. These changes made a significant impact on women's decision regarding the timing of childbearing, and the number of women delivering their first child after the age of 35 increased from 1/100 in 1970 to 1/12 in 2006.¹

However, delaying childbearing was not consequences as reflected by the data provided by the Society of Assisted Reproductive Technology (SART) regarding women undergoing infertility treatment in the United States. Indeed, despite a steady increase in human longevity over the past centuries, culminating in a life expectancy at birth that is longer than 80 years for women in the United States, age of menopause remained essentially unchanged at approximately 50, and scientific research failed to impact upon ovarian aging.^{2,3}

Within the context of assisted reproduction, ovarian aging commonly refers to the declining potential of ovaries to produce oocytes in adequate number or quality in response to controlled ovarian stimulation.⁴ Patients who present with ovarian aging are commonly diagnosed with diminished ovarian reserve (DOR) and offered in vitro fertilization (IVF) using either their own eggs or donated eggs or embryos. The number of women undergoing IVF for DOR has steadily

increased over the past decade in the United States, where DOR accounted for 8% of assisted reproductive technology (ART) cycles in 2004 and 17% in 2014, and the number of fresh IVF cycles using non-donor oocytes performed for women with DOR increased from 6947 in 2004 to 17 115 in 2014. Similarly, the number of total oocyte donation cycles (fresh and frozen-thawed oocyte or embryo) in the United States increased from 12 454 in 2004 to 19 375 in 2014.⁵

There are several potential molecular mechanisms contributing to ovarian aging, including impaired DNA repair,⁶ telomere shortening,⁷ metabolic/energetic disorders, and mitochondrial dysfunction. Understanding the mechanisms of reproductive senescence and developing strategies to prevent or delay ovarian aging would have significant medical and social implications. In this review, we will summarize data related to the association of mitochondrial dysfunction with ovarian aging.

2 | MITOCHONDRIA: STRUCTURE AND FUNCTION

Mitochondria are double-membrane-bound organelles that are ubiquitously present in the cytoplasm of eukaryotic cells and generate most of the energy of the cell in the form of adenosine triphosphate (ATP). In addition to supplying cellular energy, mitochondria are involved in other cellular activities, such as calcium homeostasis and substrate oxidation, and play a key role in the regulation of cell cycle, differentiation, senescence, and death.⁸

Mitochondrial oxidative phosphorylation (OXPHOS) system is the metabolic pathway in which cells use enzymes to oxidize nutrients,

FIGURE 1 Electron transport chain

thereby releasing energy, which is used to produce ATP. OXPHOS depends on the activity of five multi-enzymatic complexes (CI-CV) embedded in the inner mitochondrial membrane and of two mobile electron carriers, coenzyme Q10 (CoQ10) and cytochrome c (Cyt c), collectively referred to as the electron transport chain (ETC.) (Figure 1). The OXPHOS system couples the electron flow with proton translocation across the inner mitochondrial membrane generating an electrochemical gradient between the mitochondrial matrix and the intermembrane space, which then fuels ATP synthesis (Figure 1).

Mitochondria generate much of the endogenous reactive oxygen species (ROS), a potentially toxic by-product of OXPHOS. These organelles also integrate the mitochondrial permeability transition pores (mtPTPs) that initiate cell death when the mitochondrial energy function declines. Therefore, inefficiency in the OXPHOS system may result in the generation of high levels of ROS, which may then lead to cellular dysfunction and apoptosis. In fact, mitochondrial dysfunction is associated with heterogeneous group of pathologies and metabolic syndromes such as atherosclerosis, type 2 diabetes, and neurodegenerative disorders.⁹

Mitochondria are unique organelles as they contain DNA. Mitochondrial DNA (mtDNA) in mammals is a double-stranded circular molecule of about 16.6 kb and contains 37 genes. An estimated 2-10 mtDNA copies are found in each mitochondrion and 100-10 000 in each somatic cell. Mitochondrial DNA encodes two ribosomal RNAs, 22 transport RNAs, and 13 proteins, all of which are part of the ETC.¹⁰ The 13 proteins encoded by mtDNA account for only a small portion of approximately 80 proteins required for the formation of ETC. and 1500 proteins that function within mitochondria. Therefore, appropriate function of the ETC. and the mitochondria requires coordinated transcription and translation from both nuclear and mitochondrial DNA.¹¹

3 | PROPOSED MECHANISMS ON HOW MITOCHONDRIAL DYSFUNCTION MAY CONTRIBUTE TO OVARIAN AGING

In 1956, Dr. Denham Harman proposed the mitochondrial free radical theory of aging (MFRTA), which suggests that mitochondrial free radicals, produced as by-products during normal metabolism, cause

oxidative damage.¹² Free radicals (such as superoxide anion $[O_2^-]$ or hydrogen peroxide $[H_2O_2]$) are compounds that have one or more unpaired (excess) electrons. Free radicals can readily react with surrounding molecules and cause damage to DNA, proteins, and lipids by stealing their electrons through a process called oxidation. Another term that is often used within this context is reactive oxygen species (ROS), which denotes chemically reactive species containing oxygen, and includes the free radical superoxide (O_2^-) and also hydroxyl radical, singlet oxygen, and peroxides. Mitochondrion is the main source of free radicals and reactive oxygen species (ROS) in the cell.

The mitochondrial free radical theory of aging proposes that the high levels of free radicals and ROS within mitochondria may result in mtDNA mutations leading to loss of synchrony in the production on functional ETC. proteins by mtDNA and nuclear DNA, which results in partial uncoupling of the respiratory chain.^{13,14} Indeed, due to the close proximity of mtDNA to ETC., its lack of protective histones, and limited repair mechanisms, the mutation rate of mtDNA is almost 25 times higher than that of nuclear DNA.¹⁵ The proposed model further suggests that increased mtDNA mutations would cause further accumulation of ROS and increased mtDNA mutations. This cycle could result in an exponential increase in mtDNA mutation burden, which could range from ATP reduction up to cell cycle arrest or even apoptosis. Indeed, increased ROS levels, accumulated mtDNA mutations, and a decline in mitochondrial function have been reported in aging. Also consistent with this theory, mouse models that are prone to mtDNA mutations due to a PolgA (the nucleus-encoded catalytic subunit of mtDNA polymerase) deficient in proof-reading function, are associated with accumulated mtDNA mutations and a premature aging phenotype, leading to both male and female infertility.¹⁶ Recent studies showed that maternally inherited mtDNA mutations alone or in combination with somatic mtDNA mutation may induce the aging phenotype and reduce the life span in mice, and that the reduced fertility caused by mtDNA mutations in the maternal germline could be reversed by the introduction of normal mtDNA.^{17,18} These findings demonstrate that experimental introduction of mtDNA mutations in animal models results in aging in general and ovarian aging in particular. However, recent studies show that mtDNA mutation burden associated with human aging is very low,¹⁹ and biopsies of embryos for women over 40 years old do not show an increase in mtDNA mutations.²⁰ These observations strongly suggest that a model based on

MFRTA is an oversimplification and fails to explain the mechanism of ovarian senescence and overall human aging.

Multiple mitochondrial dysfunctional signaling pathways, besides mitochondrial ROS, are implicated in establishing cellular senescence (defined as an irreversible state of G1 cell cycle arrest in which cells are refractory to growth factor stimulation). These include impaired mitochondrial fusion and fission (mitochondrial dynamics),^{21,22} defective mitochondrial electron transport chain (ETC.),^{23,24} bioenergetic imbalance,^{23,25,26} altered mitochondrial metabolism,^{27,28} and altered mitochondrial membrane potential.²⁹ Above-mentioned pathways highlight the perspective for understanding the effects of mitochondrial stress on aging, linking mitochondrial dysfunction, and aging through the process of cellular senescence. However, the mechanism(s) by which these factors promote cellular senescence, and whether these pathways are conserved in all senescent cells, for example, ovarian somatic and germ cells, is poorly understood. Investigating the mechanisms of mitochondrial dysfunction in the ovary may help us understand the biology of ovarian senescence, providing new potential targets to treat age-related diseases.

4 | GENES IMPORTANT FOR MITOCHONDRIAL FUNCTION

Mitochondrial diseases are mostly caused by mutations in nuclear DNA and in some cases (about 15%) in mtDNA, which affect mitochondrial function.³⁰ Several reports based on the use of transgenic methods have revealed a key role for specific genes in mitochondrial diseases.^{31–36} Regulation of mtDNA gene expression is extremely important for modulating the OXPHOS capacity in response to metabolic requirements and in stress conditions.³⁷

To ensure proper mitochondrial function, mtDNA genome must be replicated, transcribed, and translated accurately. All the components involved in mtDNA replication and transcription are encoded in the nuclear genome, and mutations of genes essential for mtDNA replication (POL γ and TWINKLE) and for transcription/translation (TFAM, TFB1M, mTERFs, POLRMT, LRPPRC, and DARS2) are all embryonic lethal except mTERF1.^{34–36,38–44}

Mitochondria are dynamic organelles that are constantly undergoing fusion and fission to adapt to cellular energetic demands and maintain the mitochondrial function. A number of proteins belonging to the dynamin GTPase protein family are involved in mitochondrial fission and fusion.⁴⁵ To explore the implication of dynamic genes in mitochondrial function, several laboratories have investigated the impact of global genetic disruption of genes in mouse. The work from Chen et al.³¹ revealed that knockout of *Mfn1* or *Mfn2*, which are responsible for mitochondrial fusion in mouse, results in embryonic lethality due to a severe developmental delay and placental defects. OPA1, another gene involved in mitochondrial fusion, is targeted to the inner mitochondrial membrane. Homozygous deletion of OPA1 is also embryonic lethal with a block in development.^{46,47} DRP1, a large GTPase, mediates mitochondrial fission. The complete loss of *Drp1* in mice is embryonic lethal, revealing the essential role of this

gene during development.^{32,48} Moreover, Udagawa et al.⁴⁹ established oocyte-specific *Drp1*-deficient mice and found that mitochondrial fission is essential for follicular maturation and ovulation in an age-dependent manner.

Removal of unfolded or misfolded mitochondrial proteins to maintain protein hemostasis (proteostasis) is a critical function for mitochondria which houses several proteases that are in different mitochondrial compartments including CLPP, AFG3L2, PARAPLEGIN, OMA1, PHBs, OMI, and PARL. Defects in the mitochondrial protein surveillance and quality control system are associated with numerous disorders associated with impaired mitochondrial function.^{33,50–59} In addition, genes responsible for maintenance of mitochondrial dNTP pool and related to mitochondria OXPHOS also play important roles in mitochondrial function and their absence results in severe phenotypes.^{9,30,60–74} Knockout mouse models of genes involved in mtDNA stability, transcription, translation, mitochondrial dynamics, mitochondrial protein quality control, and associated phenotypes are summarized in Table 1.

5 | CLINICAL IMPLICATIONS FOR TESTING AND TREATMENT

5.1 | mtDNA copy number as a predictor of embryo viability

There seems to be a close relationship between mitochondrial dysfunction and the decline of oocyte quality and embryo developmental potential in human and in mammalian aging models.⁷⁵

The number of mitochondria in mammalian cells ranges from a few hundred to thousands, with the highest number observed in mature oocytes at approximately 100 000.⁷⁶ As the mitochondria are inherited exclusively from the oocyte, and the replication of mtDNA only resumes at blastocyst stage after the differentiation of the inner cell mass and the trophoctoderm,^{77,78} the mtDNA copy number remains unchanged during cleavage stages and reflects the mtDNA content of the oocyte.^{76,79–82}

Recent studies suggest that elevated mtDNA copy number is associated with decreased implantation potential in human IVF embryos. Fragouli et al.⁷⁵ assessed day 3 and day 5 embryos by using a combination of array-comparative genomic hybridization (aCGH), real-time quantitative PCR (RT-qPCR), and next-generation sequencing (NGS) and found that a significantly higher mtDNA copy number was observed in embryos from older women. Additionally, mtDNA levels were elevated in aneuploid embryos (independent of age) compared to euploid ones and euploid blastocysts that failed to implant compared to those that implanted. Authors established a mtDNA quantity threshold, above which pregnancy did not occur. In a subsequent study, Diez-Juan et al.⁸³ found that a higher mtDNA copy number in euploid embryos is indicative of lower embryo viability and implantation for both blastocysts and cleavage-stage embryos. The above-mentioned studies proposed that mtDNA copy number is an exciting new paradigm for embryo quality assessment. The importance of mitochondrial threshold in oocyte competence is also shown on many animal models including mouse,⁸⁴ pig,⁸⁵ and cattle.^{86,87}

TABLE 1 Knockout mouse models of genes involved in mitochondrial function

Gene	Gene function	Global/conditional knockout phenotype	Global knockout reproductive phenotype	Global knockout reference
Mitochondrial protein quality control				
CLPP	Mitochondrial matrix ATP-dependent protease	Impaired survival and hearing, resistance to ulcerative dermatitis, growth retardation, motor activity reduction	Both males and females were infertile (reduced macroscopic size of testes and ovaries already at age 3 months)	33
SPG7	Metalloprotease, a member of the AAA protein family.	Axonal degeneration, cerebellar, and hippocampal degeneration accompanied by gliosis		50, 51
AFG3L2	m-AAA protease, assembly of CI, CIII, and CIV, mitochondrial protein synthesis, mitochondrial morphology, and network maintenance	Severe progressive paralysis with premature death by P15, impaired axonal development, and delayed myelination		45
OMA1	Metalloprotease, degrade OPA1 to prevent mitochondrial fusion	Viable, marked obesity with metabolic alterations, reduced energy expenditure, and altered thermogenic response	Both males and females were fertile	53
PHB1	Mitochondrial chaperone, mtDNA maintenance, mitochondrial dynamics	Embryonic lethal		54
PHB2	Mitochondrial chaperone, mtDNA maintenance, mitochondrial dynamics	Embryonic lethal		55
HTTRA2/OMI	Serine protease, clearance of unfolded proteins	Parkinson like phenotype: neurodegeneration with loss of neurons in the striatum. Premature death by P30.		8, 10
PARL	Mitochondrial inner membrane protease, proteolytic cleavage OPA1 to inhibit apoptosis	Severe growth retardation, cachexia, atrophy of thymus, and spleen. Premature death by 8-12 wks of age. Neuronal death in thalamus and striatum.		11
PINK1	Mitochondrial serine/threonine protein kinase, involved in mitochondrial fission and motility	Parkinson's disease, evoked release of striatal dopamine, and catecholamine are reduced		52
Mitochondrial dynamics				
MFN1	Mitochondrial fusion	Embryonic lethal (E10.5-E11.5)		31
MFN2	Mitochondrial fusion	Embryonic lethal (E10.5-E11.5)		31
OPA1	Mitochondrial fusion and cristae structure	Embryonic lethal (E13.5)		46, 47
DRP1	Mitochondrial fission, mitophagy, apoptosis regulation	Embryonic lethal		32, 48
DRP1	Mitochondrial fusion	Drp1 oocyte specific knockout mice: female infertility, defective for follicular maturation and ovulation, reduced ovulation activity.		49
mtDNA stability, replication, transcription, and translation				
POLGA	mtDNA replication (catalytic subunit of Pol γ)	Embryonic lethal (E7.5-8.5)		34
POLG2	mtDNA replication (accessory subunit of Pol γ)	Embryonic lethal (E8.0-8.5)		36

(Continues)

TABLE 1 (Continued)

Gene	Gene function	Global/conditional knockout phenotype	Global knockout reproductive phenotype	Global knockout reference
TWINKLE	mtDNA replication (helicase)	Embryonic lethal		38
TFAM	mtDNA maintenance and mtDNA transcription	Embryonic lethal (E8.5-10.5)		39
TFB1M	Mitochondrial transcription and translation	Embryonic lethal (E8.5)		40
MTERF1	Transcription initiation at the light-strand promoter of the mtDNA	Viable, with normal oxidative phosphorylation capacity		41
MTERF3	Transcription terminator of the mtDNA	Embryonic lethal (E8.5-E10.5)		42
MTERF4	Mitochondrial ribosomal biogenesis	Embryonic lethal (before E8.5)		43
LRPPRC	Translation/stability of the mitochondrial mRNAs	Embryonic lethal (before E12.5)		35
DARS2	Mitochondrial aspartyl-tRNA synthetase	Embryonic lethal (E8.5)		44
Maintenance of mitochondrial dNTP pool				
ANT1	Adenine nucleotide translocase	Mitochondrial myopathy, progressive cardiac hypertrophy, metabolic abnormalities, severe exercise intolerance		60
ANT2	Adenine nucleotide translocase	Postnatal growth retardation with lactic acidosis, hematopoietic failure		61
ANT4	Adenine nucleotide translocase	Viable and exhibited apparently normal development	Male mice were infertile Female mice are fertile but produce a smaller number of progeny	62,63
TK2	Synthesis of deoxypyrimidine nucleoside triphosphates	Growth retardation, hypothermia and early mortality, neurodegeneration		64-66
TP/UP	Metabolism of nucleotides	Leukoencephalopathy		67-69
RRM2B	Synthesis of dNTPs	Severe renal failure, growth retardation, and early mortality		70
MPV17	Unknown function	Renal degeneration, hearing loss, mild hepatopathy, and myopathy		71,72
AIF	Involved in initiating a caspase-independent pathway of apoptosis	Neuronal degeneration, reduction in CI activity		73,74
FXN	Biosynthesis of the iron-sulfur clusters	Embryonic lethal (E6.5)		9
ETHE1	Mitochondrial sulfur dioxygenase (SDO)	Ethylmalonic encephalopathy, growth arrest at postnatal day 15, lethal at 5-6 wks after birth. Reduced motor activity		30

It is noteworthy that accurate quantification of the mtDNA copy number requires robust techniques. In experimental settings, qPCR is commonly used to quantitate the mtDNA copy number, with a challenging normalization to nuclear DNA when single cells (or 5–10 cells as in the case of blastocyst biopsies) are used. Another platform called droplet digital PCR is shown to be a superior method to qPCR for assessment of gene copy number variation,^{87–89} the accuracy of which is essential for studies requiring mtDNA copy number correlation to phenotypic outcomes of oocytes and embryos. More recently, a probe-based, single-tube multiplex method providing high specificity while eliminating well-to-well variability has also been described.⁹⁰ It is likely that some of these more advanced approaches and/or NGS will be utilized in clinical settings in the future, where the impact of mtDNA copy number analysis on clinical outcomes will need to be established.

5.2 | Mitochondrial nutrients

As mitochondrial dysfunction is associated with ovarian aging and is believed to negatively affect reproduction, an improvement in mitochondrial performance could potentially slow or reverse the decline in reproductive function associated with aging. Mitochondrial nutrients are naturally occurring agents that have been used to boost energy-producing capacity of mitochondria.⁹¹ These include α -lipoic acid (ALA), coenzyme Q10 (CoQ10), resveratrol, sirtuin 3 (Sirt3), rapamycin, and L-carnitine among others.¹⁷ ALA is a coenzyme that plays an important role in mitochondrial metabolism. The reduced form of ALA (dihydrolipoic acid) is a powerful mitochondrial antioxidant formed by NADH-dependent mitochondrial dihydrolipoamide dehydrogenase,⁹² and ALA is required for normal embryonic development in mice.⁹³ CoQ10 is a lipid-soluble electron transporter, which transports electrons from complexes I and II to complex III in ETC. (Figure 1) and is critical for the stabilization of complex III.⁹⁴ CoQ10 is also involved in the transport of protons into the mitochondria to maintain the membrane potential and ATP production.⁹⁵ Several studies reported decreased levels of CoQ10 associated with tissue-specific senescence.^{96,97} In addition, supplementation of CoQ10 seems to restore mitochondrial function in the oocyte and delay ovarian aging in mice.^{98–100} Dietary supplementation of CoQ10 could also possibly decrease aneuploidy rate in human oocytes.¹⁰¹ Resveratrol is a chemical reported to have the ability to delay aging and prevent age-related diseases by increasing the number of mitochondria and improving mitochondrial function.^{102,103} Resveratrol was found to improve mitochondrial performance and the number and quality of oocytes, protecting against the infertility associated with ovarian aging in mice.¹⁰⁴ Sirtuin 3 (SIRT3) is a molecule strongly implicated in aging, which can be activated by resveratrol and enhance mitochondrial biogenesis.^{105,106} It has recently been shown that inefficient SIRT3 expression may impair the developmental competence of human IVM oocytes via decreased mitochondrial biogenesis.¹⁰⁷ Rapamycin was recently found to improve oocyte competence by increasing mitochondrial autophagy and renewal and to result in prolongation of reproductive life.^{108–110} L-carnitine, a small molecule which has a critical role in fatty acid β -oxidation, could improve meiotic spindle assembly and mitochondrial distribution in metaphase II (MII) oocytes.¹¹¹ While these

initial findings are very encouraging and suggest a potential benefit from the use of mitochondrial nutrients, well-designed clinical studies are required to test the clinical efficacy and safety.

5.3 | Mitochondrial transfer

Two decades ago, transfer of cytoplasm from an oocyte obtained from a younger woman into the oocyte of an older woman was reported to improve oocyte competence and embryo development.¹¹² Subsequently, more than 50 successful live births have been achieved with this method.¹¹³ However, due to lack of adequate experimental data and the potential risk associated with the transmission third-party genetic material, the US Food and Drug Administration (FDA) suggested (in 2002) that donor cytoplasmic transfer should be suspended pending the outcome of further studies.

As oocyte cytoplasm has numerous components with yet unknown functions, mitochondrial replacement has been proposed as an alternative. Indeed, transfer of isolated mitochondria is sufficient to improve mitochondrial function and reproduction in the mouse.¹¹⁴ Spindle chromosome transfer or pronuclear transfer (PNT) of oocytes has also been proposed as potential treatments for human diseases caused by mtDNA mutations.^{115,116} However, these approaches received a significant amount of criticism worldwide due to the risk of heteroplasmy and potential long-term health consequences in the offspring.^{117,118} Indeed, studies in mice showed that heteroplasmy is associated with cognitive abnormalities¹¹⁹ and a phenotype resembling early-onset metabolic syndrome.¹²⁰ Nevertheless, mitochondrial transfer from a donor has recently been approved for use in humans in the UK for the treatment of carriers of severe mitochondrial disorders (February 3, 2015).

In the setting of age-related infertility, autologous transfer of mitochondria from somatic cells has been proposed to avoid risks associated with heteroplasmy. This method utilizes mitochondria obtained from the ovarian germline cells to eliminate the tissue-specific differences that could potentially affect embryonic development.^{121,122} Autologous germline mitochondrial energy transfer (AUGMENT) has recently been commercialized. Using this technology, a prospective cohort study with historical controls by Fakhri et al. reported an approximately eightfold to 10-fold increase in clinical pregnancy rates in a selected group of women with very poor IVF outcomes in the past.¹²³ A subsequent study using the same technology also reported a significant improvement in fertilization rates.¹²⁴ While the initial studies showed significant improvement compared to the outcomes of the same women in the past, they failed to achieve pregnancy in older women, and the true efficacy of this approach remains to be established in well-designed clinical studies without selection bias.

6 | SUMMARY AND CONCLUSIONS

Mitochondria play a number of key roles in cellular physiology and have been strongly implicated in aging through a number of mechanisms. Within the context of reproduction, mtDNA copy number may constitute a test for the prediction of embryo viability, increasing the

predictive efficacy of preimplantation genetic screening (PGS). In addition, nutrients improving mitochondrial function and replacement of functional mitochondria into the germ cells may be used in the clinical setting as a potential treatment for ovarian aging. All these exciting clinical applications will need to be validated for safety and efficacy.

DISCLOSURE STATEMENT

The authors have nothing to disclose.

REFERENCES

- Cil AP, Turkogeldi L, Seli E. Oocyte cryopreservation as a preventive measure for age-related fertility loss. *Semin Reprod Med*. 2015;33:429–435.
- Weismiller DG. Menopause. *Prim Care*. 2009;36:199–226, x.
- Bengtsson C, Lindquist O, Redvall L. Is the menopausal age rapidly changing? *Maturitas*. 1979;1:159–164.
- Seli E. Ovarian aging. *Semin Reprod Med*. 2015;33:375–376.
- Kochanek KD, Murphy SL, Xu J, Tejada-Vera B. Deaths: final data for 2014. *Natl Vital Stat Rep*. 2016;65:1–122.
- Titus S, Stobezki R, Oktay K. Impaired DNA repair as a mechanism for oocyte aging: Is it epigenetically determined? *Semin Reprod Med*. 2015;33:384–388.
- Kalmbach KH, Antunes DM, Kohlrausch F, Keefe DL. Telomeres and female reproductive aging. *Semin Reprod Med*. 2015;33:389–395.
- McBride HM, Neuspiel M, Wasiak S. Mitochondria: more than just a powerhouse. *Curr Biol*. 2006;16:R551–R560.
- Tilly JL, Sinclair DA. Germline energetics, aging, and female infertility. *Cell Metab*. 2013;17:838–850.
- Anderson S, Bankier AT, Barrell BG, et al. Sequence and organization of the human mitochondrial genome. *Nature*. 1981;290:457–465.
- Johnston IG, Williams BP. Evolutionary inference across eukaryotes identifies specific pressures favoring mitochondrial gene retention. *Cell Syst*. 2016;2:101–111.
- Harman D. Aging: a theory based on free radical and radiation chemistry. *J Gerontol*. 1956;11:298–300.
- Richter C. Oxidative damage to mitochondrial DNA and its relationship to ageing. *Int J Biochem Cell Biol*. 1995;27:647–653.
- Greaves LC, Beadle NE, Taylor GA, et al. Quantification of mitochondrial DNA mutation load. *Aging Cell*. 2009;8:566–572.
- Lynch M, Koskella B, Schaack S. Mutation pressure and the evolution of organelle genomic architecture. *Science*. 2006;311:1727–1730.
- Trifunovic A, Wredenberg A, Falkenberg M, et al. Premature ageing in mice expressing defective mitochondrial DNA polymerase. *Nature*. 2004;429:417–423.
- Ross JM, Coppotelli G, Hoffer BJ, Olson L. Maternally transmitted mitochondrial DNA mutations can reduce lifespan. *Sci Rep*. 2014;4:6569.
- Ross JM, Stewart JB, Hagstrom E, et al. Germline mitochondrial DNA mutations aggravate ageing and can impair brain development. *Nature*. 2013;501:412–415.
- Brenner CA, Wolny YM, Barritt JA, Matt DW, Munne S, Cohen J. Mitochondrial DNA deletion in human oocytes and embryos. *Mol Hum Reprod*. 1998;4:887–892.
- Alfarawati S, Fragouli E, Colls P, Wells D. Embryos of robertsonian translocation carriers exhibit a mitotic interchromosomal effect that enhances genetic instability during early development. *PLoS Genet*. 2012;8:e1003025. doi:10.1371/journal.pgen.1003025.
- Lee S, Jeong SY, Lim WC, et al. Mitochondrial fission and fusion mediators, hFis1 and OPA1, modulate cellular senescence. *J Biol Chem*. 2007;282:22977–22983.
- Seo AY, Joseph AM, Dutta D, Hwang JC, Aris JP, Leeuwenburgh C. New insights into the role of mitochondria in aging: mitochondrial dynamics and more. *J Cell Sci*. 2010;123:2533–2542.
- Stockl P, Zankl C, Hutter E, et al. Partial uncoupling of oxidative phosphorylation induces premature senescence in human fibroblasts and yeast mother cells. *Free Radic Biol Med*. 2007;43:947–958.
- Moiseeva O, Bourdeau V, Roux A, Deschenes-Simard X, Ferbeyre G. Mitochondrial dysfunction contributes to oncogene-induced senescence. *Mol Cell Biol*. 2009;29:4495–4507.
- Wang W, Yang X, Lopez de Silanes I, Carling D, Gorospe M. Increased AMP:ATP ratio and AMP-activated protein kinase activity during cellular senescence linked to reduced HuR function. *J Biol Chem*. 2003;278:27016–27023.
- Zwerschke W, Mazurek S, Stockl P, Hutter E, Eigenbrodt E, Jansen-Durr P. Metabolic analysis of senescent human fibroblasts reveals a role for AMP in cellular senescence. *Biochem J*. 2003;376:403–411.
- Jiang P, Du W, Mancuso A, Wellen KE, Yang X. Reciprocal regulation of p53 and malic enzymes modulates metabolism and senescence. *Nature*. 2013;493:689–693.
- Salminen A, Kaarniranta K, Hiltunen M, Kauppinen A. Krebs cycle dysfunction shapes epigenetic landscape of chromatin: novel insights into mitochondrial regulation of aging process. *Cell Signal*. 2014;26:1598–1603.
- Wiel C, Lallet-Daher H, Gitenay D, et al. Endoplasmic reticulum calcium release through ITPR2 channels leads to mitochondrial calcium accumulation and senescence. *Nat Commun*. 2014;5:3792.
- Dimauro S, Davidzon G. Mitochondrial DNA and disease. *Ann Med*. 2005;37:222–232.
- Chen H, Detmer SA, Ewald AJ, Griffin EE, Fraser SE, Chan DC. Mitofusins Mfn1 and Mfn2 coordinately regulate mitochondrial fusion and are essential for embryonic development. *J Cell Biol*. 2003;160:189–200.
- Ishihara N, Nomura M, Jofuku A, et al. Mitochondrial fission factor Drp1 is essential for embryonic development and synapse formation in mice. *Nat Cell Biol*. 2009;11:958–966.
- Gispert S, Parganlija D, Klinkenberg M, et al. Loss of mitochondrial peptidase Clpp leads to infertility, hearing loss plus growth retardation via accumulation of CLPX, mtDNA and inflammatory factors. *Hum Mol Genet*. 2013;22:4871–4887.
- Hance N, Ekstrand MI, Trifunovic A. Mitochondrial DNA polymerase gamma is essential for mammalian embryogenesis. *Hum Mol Genet*. 2005;14:1775–1783.
- Xu F, Addis JB, Cameron JM, Robinson BH. LRPPRC mutation suppresses cytochrome oxidase activity by altering mitochondrial RNA transcript stability in a mouse model. *Biochem J*. 2012;441:275–283.
- Humble MM, Young MJ, Foley JF, Pandiri AR, Travlos GS, Copeland WC. Polg2 is essential for mammalian embryogenesis and is required for mtDNA maintenance. *Hum Mol Genet*. 2013;22:1017–1025.
- Scarpulla RC. Transcriptional paradigms in mammalian mitochondrial biogenesis and function. *Physiol Rev*. 2008;88:611–638.
- Milenkovic D, Matic S, Kuhl I, et al. TWINKLE is an essential mitochondrial helicase required for synthesis of nascent D-loop strands and complete mtDNA replication. *Hum Mol Genet*. 2013;22:1983–1993.
- Larsson NG, Wang J, Wilhelmsson H, et al. Mitochondrial transcription factor A is necessary for mtDNA maintenance and embryogenesis in mice. *Nat Genet*. 1998;18:231–236.
- Metodiev MD, Lesko N, Park CB, et al. Methylation of 12S rRNA is necessary for in vivo stability of the small subunit of the mammalian mitochondrial ribosome. *Cell Metab*. 2009;9:386–397.
- Terzioglu M, Ruzzenente B, Harmel J, et al. MTERF1 binds mtDNA to prevent transcriptional interference at the light-strand promoter but is dispensable for rRNA gene transcription regulation. *Cell Metab*. 2013;17:618–626.
- Park CB, Asin-Cayuela J, Camara Y, et al. MTERF3 is a negative regulator of mammalian mtDNA transcription. *Cell*. 2007;130:273–285.
- Camara Y, Asin-Cayuela J, Park CB, et al. MTERF4 regulates translation by targeting the methyltransferase NSUN4 to the mammalian mitochondrial ribosome. *Cell Metab*. 2011;13:527–539.

44. Dogan SA, Pujol C, Maiti P, et al. Tissue-specific loss of DARS2 activates stress responses independently of respiratory chain deficiency in the heart. *Cell Metab*. 2014;19:458–469.
45. Hoppins S. The regulation of mitochondrial dynamics. *Curr Opin Cell Biol*. 2014;29:46–52.
46. Davies VJ, Hollins AJ, Piechota MJ, et al. Opa1 deficiency in a mouse model of autosomal dominant optic atrophy impairs mitochondrial morphology, optic nerve structure and visual function. *Hum Mol Genet*. 2007;16:1307–1318.
47. Alavi MV, Bette S, Schimpf S, et al. A splice site mutation in the murine Opa1 gene features pathology of autosomal dominant optic atrophy. *Brain*. 2007;130:1029–1042.
48. Wakabayashi J, Zhang Z, Wakabayashi N, et al. The dynamin-related GTPase Drp1 is required for embryonic and brain development in mice. *J Cell Biol*. 2009;186:805–816.
49. Udagawa O, Ishihara T, Maeda M, et al. Mitochondrial fission factor Drp1 maintains oocyte quality via dynamic rearrangement of multiple organelles. *Curr Biol*. 2014;24:2451–2458.
50. Ferreira F, Quattrini A, Pirozzi M, et al. Axonal degeneration in paraplegin-deficient mice is associated with abnormal mitochondria and impairment of axonal transport. *J Clin Invest*. 2004;113:231–242.
51. Martinelli P, La Mattina V, Bernacchia A, et al. Genetic interaction between the m-AAA protease isoenzymes reveals novel roles in cerebellar degeneration. *Hum Mol Genet*. 2009;18:2001–2013.
52. Maltecca F, Aghaie A, Schroeder DG, et al. The mitochondrial protease AFG3L2 is essential for axonal development. *J Neurosci*. 2008;28:2827–2836.
53. Quiros PM, Ramsay AJ, Sala D, et al. Loss of mitochondrial protease OMA1 alters processing of the GTPase OPA1 and causes obesity and defective thermogenesis in mice. *EMBO J*. 2012;31:2117–2133.
54. He B, Feng Q, Mukherjee A, et al. A repressive role for prohibitin in estrogen signaling. *Mol Endocrinol*. 2008;22:344–360.
55. Merkwirth C, Dargazanli S, Tatsuta T, et al. Prohibitins control cell proliferation and apoptosis by regulating OPA1-dependent cristae morphogenesis in mitochondria. *Genes Dev*. 2008;22:476–488.
56. Martins LM, Morrison A, Klusch K, et al. Neuroprotective role of the Reaper-related serine protease HtrA2/Omi revealed by targeted deletion in mice. *Mol Cell Biol*. 2004;24:9848–9862.
57. Plun-Favreau H, Burchell VS, Holmstrom KM, et al. HtrA2 deficiency causes mitochondrial uncoupling through the F(1)F(0)-ATP synthase and consequent ATP depletion. *Cell Death Dis*. 2012;3:e335. doi:10.1038/cddis.2012.77.
58. Cipolat S, Rudka T, Hartmann D, et al. Mitochondrial rhomboid PARL regulates cytochrome c release during apoptosis via OPA1-dependent cristae remodeling. *Cell*. 2006;126:163–175.
59. Kitada T, Pisani A, Porter DR, et al. Impaired dopamine release and synaptic plasticity in the striatum of PINK1-deficient mice. *Proc Natl Acad Sci USA*. 2007;104:11441–11446.
60. Graham BH, Waymire KG, Cottrell B, Trounce IA, MacGregor GR, Wallace DC. A mouse model for mitochondrial myopathy and cardiomyopathy resulting from a deficiency in the heart/muscle isoform of the adenine nucleotide translocator. *Nat Genet*. 1997;16:226–234.
61. Cho J, Seo J, Lim CH, et al. Mitochondrial ATP transporter Ant2 depletion impairs erythropoiesis and B lymphopoiesis. *Cell Death Differ*. 2015;22:1437–1450.
62. Brower JV, Rodic N, Seki T, et al. Evolutionarily conserved mammalian adenine nucleotide translocase 4 is essential for spermatogenesis. *J Biol Chem*. 2007;282:29658–29666.
63. Lim CH, Brower JV, Resnick JL, Oh SP, Terada N. Adenine nucleotide translocase 4 is expressed within embryonic ovaries and dispensable during oogenesis. *Reprod Sci*. 2015;22:250–257.
64. Zhou X, Solaroli N, Bjerke M, et al. Progressive loss of mitochondrial DNA in thymidine kinase 2-deficient mice. *Hum Mol Genet*. 2008;17:2329–2335.
65. Bartesaghi S, Betts-Henderson J, Cain K, et al. Loss of thymidine kinase 2 alters neuronal bioenergetics and leads to neurodegeneration. *Hum Mol Genet*. 2010;19:1669–1677.
66. Akman HO, Dorado B, Lopez LC, et al. Thymidine kinase 2 (H126N) knockin mice show the essential role of balanced deoxynucleotide pools for mitochondrial DNA maintenance. *Hum Mol Genet*. 2008;17:2433–2440.
67. Haraguchi M, Tsujimoto H, Fukushima M, et al. Targeted deletion of both thymidine phosphorylase and uridine phosphorylase and consequent disorders in mice. *Mol Cell Biol*. 2002;22:5212–5221.
68. Lopez LC, Akman HO, Garcia-Cazorla A, et al. Unbalanced deoxynucleotide pools cause mitochondrial DNA instability in thymidine phosphorylase-deficient mice. *Hum Mol Genet*. 2009;18:714–722.
69. Garcia-Diaz B, Garone C, Barca E, et al. Deoxynucleoside stress exacerbates the phenotype of a mouse model of mitochondrial neurogastrointestinal encephalopathy. *Brain*. 2014;137:1337–1349.
70. Kimura T, Takeda S, Sagiya Y, Gotoh M, Nakamura Y, Arakawa H. Impaired function of p53R2 in Rrm2b-null mice causes severe renal failure through attenuation of dNTP pools. *Nat Genet*. 2003;34:440–445.
71. Clozel M, Hess P, Fischli W, et al. Age-dependent hypertension in Mpv17-deficient mice, a transgenic model of glomerulosclerosis and inner ear disease. *Exp Gerontol*. 1999;34:1007–1015.
72. Viscomi C, Spinazzola A, Maggioni M, et al. Early-onset liver mtDNA depletion and late-onset proteinuric nephropathy in Mpv17 knock-out mice. *Hum Mol Genet*. 2009;18:12–26.
73. Satoh M, Kuroiwa T. Organization of multiple nucleoids and DNA molecules in mitochondria of a human cell. *Exp Cell Res*. 1991;196:137–140.
74. Scheibye-Knudsen M, Fang EF, Croteau DL, Wilson DM 3rd, Bohr VA. Protecting the mitochondrial powerhouse. *Trends Cell Biol*. 2015;25:158–170.
75. Fragouli E, Spath K, Alfarawati S, et al. Altered levels of mitochondrial DNA are associated with female age, aneuploidy, and provide an independent measure of embryonic implantation potential. *PLoS Genet*. 2015;11:e1005241. doi:10.1371/journal.pgen.1005241.
76. May-Panloup P, Chretien MF, Jacques C, Vasseur C, Malthiery Y, Reynier P. Low oocyte mitochondrial DNA content in ovarian insufficiency. *Hum Reprod*. 2005;20:593–597.
77. Eichenlaub-Ritter U, Wieczorek M, Luke S, Seidel T. Age related changes in mitochondrial function and new approaches to study redox regulation in mammalian oocytes in response to age or maturation conditions. *Mitochondrion*. 2011;11:783–796.
78. St John JC, Facucho-Oliveira J, Jiang Y, Kelly R, Salah R. Mitochondrial DNA transmission, replication and inheritance: a journey from the gamete through the embryo and into offspring and embryonic stem cells. *Hum Reprod Update*. 2010;16:488–509.
79. Chan CC, Liu VW, Lau EY, Yeung WS, Ng EH, Ho PC. Mitochondrial DNA content and 4977 bp deletion in unfertilized oocytes. *Mol Hum Reprod*. 2005;11:843–846.
80. Lin DP, Huang CC, Wu HM, Cheng TC, Chen CI, Lee MS. Comparison of mitochondrial DNA contents in human embryos with good or poor morphology at the 8-cell stage. *Fertil Steril*. 2004;81:73–79.
81. Reynier P, May-Panloup P, Chretien MF, et al. Mitochondrial DNA content affects the fertilizability of human oocytes. *Mol Hum Reprod*. 2001;7:425–429.
82. Steuerwald N, Barritt JA, Adler R, et al. Quantification of mtDNA in single oocytes, polar bodies and subcellular components by real-time rapid cycle fluorescence monitored PCR. *Zygote*. 2000;8:209–215.
83. Diez-Juan A, Rubio C, Marin C, et al. Mitochondrial DNA content as a viability score in human euploid embryos: less is better. *Fertil Steril*. 2015;104:534–541.e531.
84. Cheng Y, Wang K, Kellam LD, et al. Effects of ooplasm manipulation on DNA methylation and growth of progeny in mice. *Biol Reprod*. 2009;80:464–472.

85. El Shourbagy SH, Spikings EC, Freitas M, St John JC. Mitochondria directly influence fertilisation outcome in the pig. *Reproduction*. 2006;131:233–245.
86. Chiaratti MR, Bressan FF, Ferreira CR, et al. Embryo mitochondrial DNA depletion is reversed during early embryogenesis in cattle. *Biol Reprod*. 2010;82:76–85.
87. Chiaratti MR, Ferreira CR, Perecin F, et al. Ooplast-mediated developmental rescue of bovine oocytes exposed to ethidium bromide. *Reprod Biomed Online*. 2011;22:172–183.
88. Hindson BJ, Ness KD, Masquelier DA, et al. High-throughput droplet digital PCR system for absolute quantitation of DNA copy number. *Anal Chem*. 2011;83:8604–8610.
89. Pinheiro LB, Coleman VA, Hindson CM, et al. Evaluation of a droplet digital polymerase chain reaction format for DNA copy number quantification. *Anal Chem*. 2012;84:1003–1011.
90. Phillips NR, Sprouse ML, Roby RK. Simultaneous quantification of mitochondrial DNA copy number and deletion ratio: a multiplex real-time PCR assay. *Sci Rep*. 2014;4:3887. doi:10.1038/srep03887.
91. Bentov Y, Yavorska T, Esfandiari N, Jurisicova A, Casper RF. The contribution of mitochondrial function to reproductive aging. *J Assist Reprod Genet*. 2011;28:773–783.
92. Moini H, Packer L, Saris NE. Antioxidant and prooxidant activities of alpha-lipoic acid and dihydrolipoic acid. *Toxicol Appl Pharmacol*. 2002;182:84–90.
93. Yi X, Maeda N. Endogenous production of lipoic acid is essential for mouse development. *Mol Cell Biol*. 2005;25:8387–8392.
94. Santos-Ocana C, Do TQ, Padilla S, Navas P, Clarke CF. Uptake of exogenous coenzyme Q and transport to mitochondria is required for bc1 complex stability in yeast coq mutants. *J Biol Chem*. 2002;277:10973–10981.
95. Mitchell P. The protonmotive Q cycle: a general formulation. *FEBS Lett*. 1975;59:137–139.
96. Kalen A, Appelkvist EL, Dallner G. Age-related changes in the lipid compositions of rat and human tissues. *Lipids*. 1989;24:579–584.
97. Miles MV, Horn PS, Tang PH, et al. Age-related changes in plasma coenzyme Q10 concentrations and redox state in apparently healthy children and adults. *Clin Chim Acta*. 2004;347:139–144.
98. Ben-Meir A, Burstein E, Borrego-Alvarez A, et al. Coenzyme Q10 restores oocyte mitochondrial function and fertility during reproductive aging. *Aging Cell*. 2015;14:887–895.
99. Bentov Y, Casper RF. The aging oocyte—can mitochondrial function be improved? *Fertil Steril*. 2013;99:18–22.
100. Bentov Y, Esfandiari N, Burstein E, Casper RF. The use of mitochondrial nutrients to improve the outcome of infertility treatment in older patients. *Fertil Steril*. 2010;93:272–275.
101. Bentov Y, Hannam T, Jurisicova A, Esfandiari N, Casper RF. Coenzyme Q10 supplementation and oocyte aneuploidy in women undergoing IVF-ICSI treatment. *Clin Med Insights Reprod Health*. 2014;8:31–36.
102. Lagouge M, Argmann C, Gerhart-Hines Z, et al. Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1alpha. *Cell*. 2006;127:1109–1122.
103. Pearson KJ, Baur JA, Lewis KN, et al. Resveratrol delays age-related deterioration and mimics transcriptional aspects of dietary restriction without extending life span. *Cell Metab*. 2008;8:157–168.
104. Liu M, Yin Y, Ye X, et al. Resveratrol protects against age-associated infertility in mice. *Hum Reprod*. 2013;28:707–717.
105. Kincaid B, Bossy-Wetzel E. Forever young: SIRT3 a shield against mitochondrial meltdown, aging, and neurodegeneration. *Front Aging Neurosci*. 2013;5:48.
106. Pacella-Ince L, Zander-Fox DL, Lan M. Mitochondrial SIRT3 and its target glutamate dehydrogenase are altered in follicular cells of women with reduced ovarian reserve or advanced maternal age. *Hum Reprod*. 2014;29:1490–1499.
107. Zhao HC, Ding T, Ren Y, et al. Role of Sirt3 in mitochondrial biogenesis and developmental competence of human in vitro matured oocytes. *Hum Reprod*. 2016;31:607–622.
108. Lee SK, Zhao MH, Kwon JW, et al. The association of mitochondrial potential and copy number with pig oocyte maturation and developmental potential. *J Reprod Dev*. 2014;60:128–135.
109. Lee SE, Kim EY, Choi HY, et al. Rapamycin rescues the poor developmental capacity of aged porcine oocytes. *Asian-Australas J Anim Sci*. 2014;27:635–647.
110. Zhang XM, Li L, Xu JJ, et al. Rapamycin preserves the follicle pool reserve and prolongs the ovarian lifespan of female rats via modulating mTOR activation and sirtuin expression. *Gene*. 2013;523:82–87.
111. Moawad AR, Xu B, Tan SL, Taketo T. L-carnitine supplementation during vitrification of mouse germinal vesicle stage-oocytes and their subsequent in vitro maturation improves meiotic spindle configuration and mitochondrial distribution in metaphase II oocytes. *Hum Reprod*. 2014;29:2256–2268.
112. Cohen J, Scott R, Schimmel T, Levron J, Willadsen S. Birth of infant after transfer of anucleate donor oocyte cytoplasm into recipient eggs. *Lancet*. 1997;350:186–187.
113. May-Panloup P, Boucret L, Chao de la Barca JM, et al. Ovarian ageing: the role of mitochondria in oocytes and follicles. *Hum Reprod Update*. 2016;22:725–743.
114. Van Blerkom J, Sinclair J, Davis P. Mitochondrial transfer between oocytes: potential applications of mitochondrial donation and the issue of heteroplasmy. *Hum Reprod*. 1998;13:2857–2868.
115. Craven L, Tuppen HA, Greggains GD, et al. Pronuclear transfer in human embryos to prevent transmission of mitochondrial DNA disease. *Nature*. 2010;465:82–85.
116. Paull D, Emmanuele V, Weiss KA, et al. Nuclear genome transfer in human oocytes eliminates mitochondrial DNA variants. *Nature*. 2013;493:632–637.
117. Reinhardt K, Dowling DK, Morrow EH. Mitochondrial replacement, evolution, and the clinic. *Science*. 2013;341:1345–1346.
118. Isasi R, Kleiderman E, Knoppers BM. Genetic technology regulation. Editing policy to fit the genome? *Science*. 2016;351:337–339.
119. Sharpley MS, Marciniak C, Eckel-Mahan K, et al. Heteroplasmy of mouse mtDNA is genetically unstable and results in altered behavior and cognition. *Cell*. 2012;151:333–343.
120. Acton BM, Lai I, Shang X, Jurisicova A, Casper RF. Neutral mitochondrial heteroplasmy alters physiological function in mice. *Biol Reprod*. 2007;77:569–576.
121. Takeda K, Tasai M, Akagi S, et al. Microinjection of serum-starved mitochondria derived from somatic cells affects parthenogenetic development of bovine and murine oocytes. *Mitochondrion*. 2010;10:137–142.
122. Schatten H, Sun QY, Prather R. The impact of mitochondrial function/dysfunction on IVF and new treatment possibilities for infertility. *Reprod Biol Endocrinol*. 2014;12:111. doi:10.1186/1477-7827-12-111.
123. Woods DC, Tilly JL. Autologous Germline Mitochondrial Energy Transfer (AUGMENT) in human assisted reproduction. *Semin Reprod Med*. 2015;33:410–421.
124. Oktay K, Baltaci V, Sonmezer M, et al. Oogonial precursor cell-derived autologous mitochondria injection to improve outcomes in women with multiple IVF failures due to low oocyte quality: a clinical translation. *Reprod Sci*. 2015;22:1612–1617.

How to cite this article: Wang T, Zhang M, Jiang Z, Seli E. Mitochondrial dysfunction and ovarian aging. *Am J Reprod Immunol*. 2017;77:e12651. <https://doi.org/10.1111/aji.12651>