

1-1-2015

The ubiquitin ligase Siah2 regulates obesity-induced adipose tissue inflammation

Gail Kilroy
Pennington Biomedical Research Center

Lauren E. Carter
Pennington Biomedical Research Center

Susan Newman
Pennington Biomedical Research Center

David H. Burk
Pennington Biomedical Research Center

Justin Manuel
Pennington Biomedical Research Center

See next page for additional authors

Follow this and additional works at: https://digitalcommons.lsu.edu/biosci_pubs

Recommended Citation

Kilroy, G., Carter, L., Newman, S., Burk, D., Manuel, J., Möller, A., Bowtell, D., Mynatt, R., Ghosh, S., & Floyd, Z. (2015). The ubiquitin ligase Siah2 regulates obesity-induced adipose tissue inflammation. *Obesity*, 23 (11), 2223-2232. <https://doi.org/10.1002/oby.21220>

This Article is brought to you for free and open access by the Department of Biological Sciences at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Authors

Gail Kilroy, Lauren E. Carter, Susan Newman, David H. Burk, Justin Manuel, Andreas Möller, David D. Bowtell, Randall L. Mynatt, Sujoy Ghosh, and Z. Elizabeth Floyd

The Ubiquitin Ligase Siah2 Regulates Obesity-Induced Adipose Tissue Inflammation

Gail Kilroy¹, Lauren E. Carter¹, Susan Newman¹, David H. Burk¹, Justin Manuel¹, Andreas Möller², David D. Bowtell³, Randall L. Mynatt¹, Sujoy Ghosh^{1,4}, and Z. Elizabeth Floyd¹

Objective: Chronic, low-grade adipose tissue inflammation associated with adipocyte hypertrophy is an important link in the relationship between obesity and insulin resistance. Although ubiquitin ligases regulate inflammatory processes, the role of these enzymes in metabolically driven adipose tissue inflammation is relatively unexplored. Herein, the effect of the ubiquitin ligase Siah2 on obesity-related adipose tissue inflammation was examined.

Methods: Wild-type and Siah2KO mice were fed a low- or high-fat diet for 16 weeks. Indirect calorimetry, body composition, and glucose and insulin tolerance were assayed along with glucose and insulin levels. Gene and protein expression, immunohistochemistry, adipocyte size distribution, and lipolysis were also analyzed.

Results: Enlarged adipocytes in obese Siah2KO mice were not associated with obesity-induced insulin resistance. Proinflammatory gene expression, stress kinase signaling, fibrosis, and crown-like structures were reduced in the Siah2KO adipose tissue, and Siah2KO adipocytes were more responsive to insulin-dependent inhibition of lipolysis. Loss of Siah2 increased expression of PPAR γ target genes involved in lipid metabolism and decreased expression of proinflammatory adipokines regulated by PPAR γ .

Conclusions: Siah2 links adipocyte hypertrophy with adipocyte dysfunction and recruitment of proinflammatory immune cells to adipose tissue. Selective regulation of PPAR γ activity is a Siah2-mediated mechanism contributing to obesity-induced adipose tissue inflammation.

Obesity (2015) **23**, 2223–2232. doi:10.1002/oby.21220

Introduction

Obesity-associated insulin resistance is linked to dysregulation of lipid storage and chronic, low-grade inflammation of adipose tissue (1). As adipose tissue expands to accommodate the lipid storage demands of excess energy intake, adipocyte hypertrophy becomes a defining characteristic. When the capacity to expand is exceeded, inflammatory signaling in adipose tissue is activated (2). Adipocyte secretion of proinflammatory adipokines correlates with infiltration of M1-like macrophages and proinflammatory T lymphocytes (1), setting up an inflammatory state focused on removing necrotic adipocytes (3). Accompanied by increased release of fatty acids from adipose tissue, this leads to impaired insulin signaling in skeletal muscle and liver and systemic insulin resistance (4).

Adipose tissue inflammation is sustained by a positive feedback loop in which cytokines secreted by infiltrating macrophages activate stress kinase signaling pathways in adipocytes and macrophages that up-regulate proinflammatory genes via activator protein-1 (AP-1) and nuclear factor- κ B (NF- κ B) transcriptional activity (5). Although signaling events controlling NF- κ B activation (6) are regulated by enzymes of the ubiquitin–proteasome system, involvement of the ubiquitin–proteasome system in obesity-induced adipose tissue inflammation is relatively unexplored.

Post-translational modification of proteins by ubiquitin, the major pathway controlling non-lysosomal intracellular protein degradation, begins with ubiquitin binding to the ubiquitin activating enzyme (E1), followed by transfer of ubiquitin from E1 to the targeted protein via ubiquitin conjugating enzymes (E2s) and ubiquitin ligases

¹ Pennington Biomedical Research Center, Baton Rouge, Louisiana, USA. Correspondence: Z. Elizabeth Floyd (elizabeth.floyd@pbrc.edu) ² QIMR Berghofer Medical Research Institute, Herston, Queensland, Australia ³ Peter MacCallum Cancer Centre, East Melbourne, Victoria, Australia ⁴ Cardiovascular and Metabolic Disease Program and Center for Computational Biology, Duke-NUS Graduate Medical School, Singapore, Singapore.

Funding agencies: This work is supported by the National Institutes of Health (NIDDK, 5R56DK89020, ZEF and NIDDK, 1R01DK099625, ZEF). This work used the Cell Biology and Bioimaging Core and the Genomics Core facilities at Pennington Biomedical Research Center that are supported in part by COBRE (NIH 8P20-GM103528) and NORC (NIH 2P30-DK072476) center grants from the National Institutes of Health. This work also used Core Services supported by grant DK097153 of the National Institutes of Health to the University of Michigan.

Disclosure: The authors declare no conflict of interest.

Additional Supporting Information may be found in the online version of this article.

Received: 28 April 2015; **Accepted:** 16 June 2015; **Published online** 18 September 2015. doi:10.1002/oby.21220

(E3s), which determine the specificity of ubiquitylation (7). Deletion of E3 ligases c-Cbl (8) or ITCH (9), an E3 ligase involved in T-cell differentiation, increases energy expenditure and prevents high fat-induced obesity. Our studies in 3T3-L1 adipocytes found the ubiquitin ligase mammalian homologue of seven-in-absentia-2 (Siah2) alters peroxisome proliferator-activated receptor γ (PPAR γ) protein levels and selectively regulates PPAR γ activity (10). Given the central role of PPAR γ in forming and maintaining adipocytes, regulating insulin sensitivity, and inflammatory gene expression in adipocytes and macrophages (11), we hypothesized that Siah2 regulates obesity-induced changes in adipose tissue. In this study, we examined the adipose tissue phenotype in global Siah2-null mice challenged with chronic excess energy intake.

Methods

Experimental animals

Siah2KO mice were generated and maintained as described (12,13). All animal experiments were approved by the Pennington Biomedical Research Center Animal Care and Use Committee. The animals were housed with a 12-h light-dark cycle at 24°C. At four weeks of age, wild-type (WT) and Siah2KO male mice were randomly assigned ($n = 10/\text{group}$) to a defined 10% low-fat diet (LFD) or 45% high-fat diet (HFD) and were fed *ad libitum* for 4 months thereafter. Body weight was measured weekly and body composition was measured bi-weekly by NMR. Food intake, activity, and indirect calorimetry were measured at 12 weeks on each diet (TSE PhenoMaster). At the end of the study, the mice were euthanized between 8 and 11 am. Adipose tissue was harvested for analysis of adipocyte size distribution and lipolysis from a separate cohort of WT and Siah2KO male mice maintained on LFD or HFD for 2 months.

Glucose and insulin tolerance tests

For the glucose (GTT) and insulin (ITT) tolerance tests, the amount of glucose or insulin administered was normalized to fat-free mass (FFM) (14), which did not vary significantly among groups ($20.1 \pm 0.13 \text{ g}$) at 12 weeks on each diet. Mice were fasted 4 h prior to administering 2 g/kg FFM of glucose/mouse (GTT) or 1 U/kg lean mass insulin/mouse (HumulinR) (ITT) by intraperitoneal injection.

Blood chemistry and lipolysis

Fasting serum glucose levels were measured by hexokinase activity assay (Sigma). Fasting insulin levels were assayed via ELISA (Cystal Chem). Serum nonesterified fatty acids (Abcam) and triglycerides (Eagle Diagnostics) levels were assayed according to manufacturers' instructions. For lipolysis assays, adipocytes were isolated by collagenase treatment from epididymal adipose tissue of HFD-fed mice (15). Insulin-mediated inhibition of lipolysis in the presence of isoproterenol (0.1 μM) was assessed by glycerol release (Zen-Bio).

Microarray analysis

Epididymal adipose tissue RNA (RNA integrity number ≥ 8) was analyzed for gene expression on Illumina MouseRef-8v2.0 expression arrays. RNA from 8 to 10 animals/group was combined into three pooled samples/group. Raw gene expression signals were background adjusted and quantile normalized using GenomeStudio

(V2011.1.Illumina Inc.). For each sample, probes were considered "expressed" if their detection P -value was <0.05 . Probes that failed to reach a detection P -value <0.05 in any sample across the four treatments (WT high fat, low fat; Siah2KO high fat, low fat) were removed from analysis. Remaining probes were log transformed (base 2) and treatment specific fold-changes computed as log ratios. The statistical significance of differential expression was ascertained by regularized t -test, based on a Bayesian probabilistic framework (16). Genes with an absolute fold-change ≥ 1.5 -fold and $P \leq 0.001$ were considered significantly differentially expressed, unless otherwise noted. Biological pathway analysis was conducted on the gene expression signals using gene-set enrichment (GSEA) or overrepresentation based approaches (Ingenuity Pathway Analysis, Ingenuity). Statistical analyses were controlled for multiple testing via the false discovery rate (FDR) (17). The microarray dataset was submitted to Gene Expression Omnibus (GEO) data repository (GSE61839).

Quantitative PCR

Total RNA was purified from epididymal adipose tissue, (200 ng) reverse transcribed and real-time PCR performed with TaqMan chemistry as described (10). The results were normalized to *ubiquitin B* mRNA levels and analyzed by the $2^{-\Delta\Delta\text{CT}}$ method with WT values used as the calibrator. The gene list is provided in Supporting Information Table S1.

Preparation of whole cell extracts and immunoblotting

Adipose tissue was homogenized in a denaturing buffer and processed for immunoblotting or immunoprecipitation as described (10). Nitrocellulose membranes were incubated with antibodies (Supporting Information Table S2) for 1-2 h at room temperature or overnight at 4°C.

Immunohistochemistry and immunostaining

Epididymal adipose tissue was fixed in 10% formalin, then embedded in paraffin, sectioned onto slides and hematoxylin and eosin (H&E) stained. Adipose tissue collagen content was determined by trichrome staining and macrophage content by immunodetection using anti-Iba1 antibody.

Adipocyte area and size distribution

H&E stained inguinal and epididymal adipose tissue were analyzed using Image J software programmed to measure the area of each adipocyte based on size and shape exclusion limits. The number of adipocytes counted/experimental condition ranged from 2,639-14,819. Epididymal adipocyte size distribution was determined as described (18) and analyzed on a Multisizer-3 (Beckman Coulter) using a 400- μm aperture with a dynamic linear range 12-320 μm and reported as the mean at 20 μm intervals from 22 to 240 μm .

Statistical analysis

Normal distribution of glucose and insulin levels, food intake, and body weight was assessed using the D'Agostino-Pearson omnibus normality test. Statistical significance for body weight, GTT, and ITT was determined using ANOVA. Statistical significance for all other data was determined using an unpaired two-tailed t -test. JMP Pro




Figure 1 Siah2KO mice fed HFD are obese with reduced energy expenditure compared to WT. (A) Body weight, (B) percent FM, and (C) percent FFM were measured in the WT and Siah2KO mice fed a defined LFD or HFD over 16 weeks. (D) Food intake and (E) energy expenditure were measured at 12 weeks on each diet. Dark bars indicate 7P-7A. (F) Regression analysis of energy expenditure (kcal/h) related to body weight or FFM (lean mass) was carried out using the daily energy expenditure of each animal over three consecutive days. Activity and RER were also measured (Supporting Information Figure S1). Statistical significance compared to WT, * $P < 0.05$.

10.0 (SAS Institute) and GraphPad Prism 5 software were used for statistical analyses. Variability was expressed as the mean \pm SEM.

Results

Siah2 regulates energy expenditure in diet-induced obesity

To characterize the role of Siah2 in obesity, we conducted a feeding study with a 10% LFD or 45% HFD diet for 4 months. Based on our studies in 3T3-L1 adipocytes showing Siah2 promotes adipogenesis (10), we anticipated loss of Siah2 *in vivo* would lead to reduced body weight and fat mass (FM), and that Siah2KO mice would develop insulin resistance more readily than WT mice when fed HFD. However, HFD Siah2KO mice body weight trended higher than WT and body weight for the LFD Siah2KO mice was significantly higher than WT mice until late in the study (Figure 1A). After 8 weeks, there was no significant difference in the per-

cent FM within each diet group (Figure 1B), but percent FFM was lower in the Siah2KO LFD mice at 4 and 12 weeks (Figure 1C). Food intake was unaffected by genotype (Figure 1D), although body weight positively correlated with food intake for both groups on the HFD (Supporting Information Figure S1A). Energy expenditure was measured after 12 weeks on each diet. Genotype had no effect on energy expenditure in the LFD-fed mice. However, unlike WT, energy expenditure of the Siah2KO mice did not increase with HFD (Figure 1E) and loss of Siah2 had a significant effect on the relationship between energy expenditure and body weight or lean mass in the HFD-fed mice (Figure 1F). HFD Siah2KO mice activity was significantly lower than WT (Supporting Information Figure S1C). When the reduced activity of the HFD Siah2KO mice was considered, lower energy expenditure in the Siah2KO mice was largely accounted for by genotype, FFM, and activity using analysis of covariance (19). However, the effect of genotype on energy expenditure remained significant, suggesting Siah2 also affects energy expenditure independent of body composition or activity (Figure 1F).




Figure 2 Siah2KO adipocytes are enlarged but more insulin-sensitive than WT. (A) H&E staining of HFD WT and Siah2KO epididymal fat. (B) Adipocyte size (area) quantitated using Image J. iWAT, inguinal adipose tissue, eWAT, epididymal adipose tissue. (C) Adipocyte size distribution of epididymal adipocytes at 2 months on LFD or HFD analyzed using Coulter counting. (D) WT and Siah2KO (KO) serum-free fatty acids and triglyceride levels. (E) Insulin-dependent inhibition of lipolysis in isolated HFD WT and Siah2KO epididymal adipocytes. Statistical significance compared to WT, * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

Obese Siah2KO mice have improved insulin sensitivity

To investigate the adipose tissue of the Siah2KO mice, we carried out H&E staining of the epididymal fat pad. Adipocytes from obese Siah2KO mice were hypertrophied but more uniform in size and shape than obese WT adipocytes (Figure 2A). Adipocyte area measurements indicated Siah2KO adipocytes trended larger than WT adipocytes independent of diet or fat depot (Figure 2B). Analysis of adipocyte size distribution after 2 months on LFD or HFD shows the shift toward larger adipocytes began earlier in the Siah2KO mice than WT (Figure 2C), consistent with Siah2 affecting adipocyte size independent of diet, body weight, or FM. However, circulating free fatty acid and triglyceride levels were significantly lower in the LFD- and HFD-fed Siah2KO mice, and triglyceride levels were lower in the obese

Siah2KO mice compared to WT (Figure 2D). This suggests the hypertrophied Siah2KO adipocytes were less lipolytic. To test this, adipocytes were isolated from obese WT and Siah2KO mice and assayed for insulin-mediated inhibition of adrenergic-stimulated lipolysis. Siah2KO adipocytes were significantly more sensitive to insulin-mediated inhibition of lipolysis (Figure 2E).

The lower free fatty acid levels corresponded to lower blood glucose levels in the Siah2KO mice, and the glucose levels did not increase with obesity in the absence of Siah2 (Figure 3A). Fasting insulin levels trended lower in the lean Siah2KO mice compared to lean WT mice and increased with diet-induced obesity, but they remained significantly lower than WT (Figure 3B). Although insulin levels in the Siah2KO mice increased with obesity, glucose (Figure 3C,D) and insulin tolerance (Figure 3E,F) tests showed loss of Siah2




Figure 3 Carbohydrate metabolism is improved in the lean or obese Siah2KO mice. (A) Fasting blood glucose and (B) insulin levels assayed at 16 weeks on the LFD or HFD. (C) GTT and (D) GTT AUC and (E) ITT and (F) ITT AUC at 12 weeks on each diet. Statistical significance compared to WT, [#] $P < 0.05$ for LFD and $^*P < 0.05$; $^{**}P < 0.01$ for HFD.

improved glucose tolerance and insulin sensitivity in the lean and obese mice. Notably, glucose and insulin tolerance in the obese Siah2KO mice was comparable to lean WT, indicating Siah2 contributes to the negative effects of obesity-induced adipocyte hypertrophy on glucose metabolism.

Siah2KO mice have reduced adipose tissue inflammation

To begin understanding the mechanism underlying the effect of Siah2 on insulin sensitivity and adipocyte size, we conducted micro-

array analysis of Siah2KO and WT epididymal adipose tissue. The most significant differences in gene expression were related to inflammatory responses, hematological function, and immune cell trafficking ($P < 0.02$) (Figure 4A). Specifically, pathways related to Toll-like receptor signaling, B- and T-cell receptor signaling, and natural killer cell-mediated cytotoxicity were significantly down-regulated ($P < 0.01$) in HFD-fed Siah2KO adipose, compared to WT samples (Supporting Information Table S3). Hierarchical clustering of the top 50 differentially expressed transcripts showed down-regulation of a broad range of genes involved in proinflammatory responses (Figure 4B).




Figure 4 Siah2 controls adipose tissue inflammation. WT and Siah2KO epididymal adipose tissue was evaluated using microarray analysis of gene expression. (A) Most significant biological functions affected by Siah2 determined by Ingenuity software analysis. (B) Hierarchical clustering of the top 50 genes up- or down-regulated by Siah2 in the LFD- or HFD-fed mice. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

Adipose tissue expression of macrophage markers, including *F4/80*, *cd68*, *cd11c*, *cd11d*, *trem2*, and *Tyrobp*, was reduced in the visceral adipose tissue of the obese Siah2KO mice (Figure 5A). Although proinflammatory markers were decreased (*F4/80*, *cd11c*, *cd11d*), expression of *Arg-1*, an indicator of anti-inflammatory macrophage, was increased. Expression of proinflammatory cytokines, including *tnfa*, *ccl2*, *ccr2* (CCL2 receptor), *gdf3*, *saa3*, and *pai-1*, were also significantly reduced in the Siah2KO adipose tissue (Figure 5B, full gene name given in Supporting Information Table S1). Two of these genes encode proteins strongly associated with inflammation and insulin resistance. Serum amyloid A3 (SAA3) is a proinflammatory and lipolytic cytokine secreted by macrophages and adipocytes that promotes macrophage recruitment (20). Plasminogen activator inhibitor-1 (serpine-1/PAI-1) is a serine protease inhibitor that is associated with obesity, inflammation, and insulin resistance in rodents and humans (21). In contrast, adipsin, a complement factor secreted by adipocytes whose expression is increased by PPAR γ agonists (22), but decreased in obesity (23), was expressed at higher levels in Siah2KO mice. Siah2-mediated changes in inflammatory

markers also include down-regulation of *tcirg1*, a T lymphocyte membrane ATPase that promotes proinflammatory T cell activation (24), and up-regulation of *fox3p*, a marker of the anti-inflammatory regulated T (T_{reg}) cells (Figure 5C).

Blunted proinflammatory responses to obesity in Siah2KO adipose tissue were further demonstrated by comparing inflammatory gene expression in the LFD- and HFD-fed mice (Figure 5D). Although inflammatory marker gene expression was consistently reduced in the HFD-fed Siah2KO adipose tissue, gene expression in the LFD Siah2KO adipose tissue was either unchanged or increased, including *adam 8*, an extracellular matrix metallopeptidase up-regulated with adipose tissue inflammation (25), and *col6a2*, a collagen VI subunit.

Reduced fibrosis and stress kinase activation in Siah2KO mice

The extracellular matrix surrounding adipocytes contributes to the relationship between adipocyte expandability and developing adipose




Figure 5 Siah2 regulates inflammatory gene expression in adipose tissue of HFD-fed mice. Gene expression of markers of (A) inflammation, (B) cytokines and chemokines, and (C) T lymphocytes was assayed in the HFD-fed mice or (D) compared between the LFD- and HFD-fed mice using real-time qRT-PCR. Statistical significance compared to WT, * $P < 0.01$; ** $P < 0.001$.

tissue inflammation with obesity (2). The inflammatory response to adipocyte hypertrophy leads to excess collagen deposition and adipose tissue fibrosis, restricting the ability of the adipocytes to expand further (26). Khan et al. showed collagens IV and VI are generally expressed at high levels in the visceral fat depot of db/db mice and are down-regulated by $PPAR\gamma$ activation or overexpression of adiponectin (2). Furthermore, we previously demonstrated that Siah2 has a promoting role for tissue fibrosis through collagens in wound healing (27). While the collagen IV subunit *col4a6* was up-regulated in Siah2KO adipose tissue, collagen VI subunit *col6a2* was reduced and *col6a3* was unchanged (Figure 6A). *Adam8* was twofold lower in the Siah2KO visceral fat. Lumican, a

proteoglycan promoting fibrosis that is negatively regulated by $PPAR\gamma$, was reduced although decorin, another proteoglycan that regulates fibrosis (2), was unchanged (Figure 6A).

To determine whether changes in expression of inflammatory markers and extracellular matrix components correlate with reduced fibrosis, trichrome staining of the Siah2KO epididymal adipose tissue was compared to WT (Figure 6B). The widespread collagen deposition present in WT was absent in the Siah2KO visceral fat. Consistent with reduced fibrosis, there were significantly fewer macrophages configured as crown-like structures surrounding adipocytes in the Siah2KO adipose tissue (Figure 6B, Iba-1).




Figure 6 Siah2 deficiency decreases markers of adipocyte dysfunction in obesity. Indicators of adipocyte dysfunction were assayed in epididymal adipose tissue from HFD-fed mice. **(A)** Extracellular matrix components were determined using real-time qRT-PCR. Statistical significance compared to WT, * $P < 0.05$; ** $P < 0.01$. **(B)** Fibrosis was determined using trichrome staining of collagen, and crown-like structures were visualized using anti-Iba-1 immunostaining of macrophages. **(C)** Stress kinase activation was determined via Western blot analysis. β -actin is included as a loading control. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

Fibrotic changes with adipocyte expansion in obesity cause mechanical stress that activates stress kinase signaling pathways, including the Jun N-terminal kinase (JNK) and extracellular signal-regulated (ERK p42/44) MAPK pathways (5). Activation of JNK and NF κ B by TNF α elicits proinflammatory changes in obese adipose tissue (1) and *in vitro* TNF α -mediated insulin resistance depends on activation of the ERK p42/44-MAPK pathway in adipocytes (28). Figure 6C demonstrates that activation of these key signaling pathways was attenuated in the obese Siah2KO animals.

Partial activation of PPAR γ in the Siah2KO adipose tissue

PPAR γ transcriptional activity plays a pivotal role in regulating adipose tissue inflammation in response to chronic energy excess (11,29). Our studies in 3T3-L1 adipocytes indicate Siah2 regulates expression of a subset of PPAR γ target genes and affects ligand-dependent degradation of PPAR γ proteins (10). Therefore, we assayed the expression of a subset of PPAR γ target genes that were differentially regulated in our microarray analysis. Although classic markers of adipogenesis such as *pck-1* and *fabp4* were down-regulated in the Siah2KO epididymal fat, several genes encoding proteins involved in fatty acid transport and storage (*lpl*, *cd36*, *soat1*) and lipid synthesis (*fasn*, *lipin1*) were up-regulated (Figure 7A). While the increase in *lpl* expression was modest, expression of *fasn*, *cd36*, *lipin1*, and *soat1* was increased two- to fivefold. Activation of PPAR γ was not associated with regulation of PPAR γ 1 or PPAR γ 2 gene expression in the Siah2KO adipose tissue (Figure 7B) but did correlate with increased PPAR γ protein levels (Figure 7C-E) and decreased PPAR γ ubiquitylation (Figure 7F,G). Although identified as a target of Siah2 *in vitro* (30), nuclear corepressor-1 (NCoR) levels were decreased, and the levels of Siah2 target proteins, histone deacetylase-3 (HDAC3) (31), and prolyl-3-hydroxylase (PHD3) (32) were unaltered.

Discussion

The ubiquitin-proteasome system is well described as a critical regulator of immune responses (6). For example, TNF α -mediated

activation of NF κ B is regulated by multiple ubiquitin ligases, including several TNF receptor-associated factors (TRAFs 2, 5, and 6) as well as A20/TNF α -induced protein-3 (6) and ITCH, a JNK phosphorylation-dependent ligase that modulates TNF α signaling (33). Much less is known about how ubiquitin ligases affect the chronic inflammation that occurs in response to obesity. In this study, we report the relationship between obesity-related adipocyte hypertrophy, adipose tissue inflammation, and insulin sensitivity is regulated by the ubiquitin ligase Siah2. Although Siah2KO adipocytes trended larger than WT, deletion of Siah2 protected mice with diet-induced obesity from adipose tissue inflammation, and insulin sensitivity in the obese Siah2KO animals was comparable to lean WT. The striking decrease in proinflammatory cytokine and chemokine expression along with reduced fibrosis and crown-like structures and elevated levels of regulatory T cell markers indicates Siah2 is necessary for adipocyte hypertrophy to elicit the inflammatory changes in obese adipose tissue that lead to insulin resistance. Several mechanisms may underlie the effect of Siah2 on the relationship between adipocyte hypertrophy and insulin resistance.

Although a sustained influx of M1-like macrophages in adipose tissue restricts adipocyte expansion (2,34), macrophage infiltration is essential for healthy adipose tissue remodeling in response to excess energy intake. As adipocytes expand, dysfunctional enlarged adipocytes are removed by macrophages and replaced by newly formed adipocytes (35). However, the early distribution toward larger adipocytes in the lean and obese Siah2KO mice and our *in vitro* studies showing loss of Siah2 impairs adipogenesis (10) argue against accelerated adipose tissue remodeling in the Siah2KO animals. Instead, the broad transcriptional response in markers of inflammation suggests Siah2 regulates the influx, production, or activation of proinflammatory immune cells. Initial studies of Siah2 found Siah2 deletion enhances expansion of bone marrow-derived myeloid progenitor cells *in vitro* without altering the levels of myeloid cells *in vivo* (12). Our study found that Siah2 deficiency substantially reduces expression of *trem2* in adipose tissue, which encodes a protein found on monocyte-derived dendritic cells that promotes myeloid cell activation (36). The possibility that Siah2 is affecting inflammation via




Figure 7 Loss of Siah2 selectively regulates PPAR γ target genes involved in lipid metabolism and alters PPAR γ protein levels. All measurements were carried out in epididymal adipose tissue from HFD-fed mice. (A) A subset of PPAR γ target genes selected based on microarray analysis was analyzed via real-time qRT-PCR. (B) Gene expression of PPAR γ 1, PPAR γ 2, or PPAR γ 1 and PPAR γ 2 combined. (C) Steady-state level of PPAR γ proteins and selected Siah2 target proteins. (D) PPAR γ protein levels from expanded number of mice. β -actin is included as a loading control in panels C and D. (E) Quantification of PPAR γ protein levels in panel D. (F) Ubiquitin modification of PPAR γ determined by Western blot analysis of PPAR γ (IP: PPAR γ , IB: PPAR γ) or ubiquitin (IP: PPAR γ , IB: ubiquitin) after immunoprecipitation of PPAR γ from WT or Siah2KO epididymal adipose tissue. (G) Quantification of high-molecular-weight PPAR γ ubiquitin conjugates. Statistical significance compared to WT, * $P < 0.05$; ** $P < 0.01$.

TREM-2 is supported by evidence that dendritic cells promote macrophage infiltration of adipose tissue (37).

Our data also show the absence of proinflammatory changes in the obese Siah2KO visceral adipose tissue corresponds to increased lipid storage capacity in the enlarged adipocytes. Isolated Siah2KO adipocytes were more sensitive to the anti-lipolytic effect of insulin, and this was reflected *in vivo* by reduced circulating fatty acids and triglyceride levels. The “lipid steal” hypothesis proposes that PPAR γ activation in adipose tissue enhances the ability of adipocytes to store excess lipids in obesity (11). This protects other insulin-responsive tissues from reactive lipids, resulting in improved insulin sensitivity. Although markers of adipogenesis such as *fabp4*, *pck1*, and *adiponectin* are not increased, PPAR γ target genes involved in

lipid uptake, lipid synthesis, and lipid storage are up-regulated in the adipose tissue. Increased lipid storage in the hypertrophied adipocytes is consistent with our observation that genotype significantly contributed to decreased energy expenditure in the Siah2KO obese mice independent of body composition. Several adipokines influenced by PPAR γ activation that impact adipose tissue inflammation are also regulated by Siah2. Higher adiponectin levels and lower *mcp-1*, *saa3*, and *pai-1* levels are associated with ligand activation of PPAR γ (11), but unaltered expression of adiponectin and other PPAR γ targets such as *pck-1* suggests that a role for PPAR γ transcriptional activity in Siah2-mediated regulation of inflammation does not occur via classical PPAR γ ligand binding. This is consistent with elevated PPAR γ protein levels accompanied by only modest reductions in PPAR γ ubiquitylation. PPAR γ -mediated changes in

lipid storage capacity can be regulated by TNF α (1,29). TNF α inhibits PPAR γ activity via multiple mechanisms (29), including caspase-dependent degradation of PPAR γ proteins (38,39). Moreover, Siah proteins are structurally similar to the TRAF ubiquitin ligases that regulate TNF α signaling (40). Therefore, reduced caspase cleavage of PPAR γ proteins may also regulate PPAR γ protein levels and activity in the Siah2KO animals.

The well-described role of Siah2 in regulating cellular responses to hypoxia (32) suggests Siah2 deletion inhibits hypoxia-related adipose tissue fibrosis and the subsequent influx of immune cells (26). This possibility is supported by our evidence that fibrosis was reduced in the adipose tissue of obese Siah2KO mice. A Siah2-mediated effect on hypoxia is unlikely to occur via a HIF-1 α -dependent pathway since Siah2 deletion does not affect the protein levels of PHD3, the Siah2 target protein responsible for targeting HIF-1 α for degradation. However, Siah2 also affects hypoxia responses via HIF-1 α -independent pathways that involve MAPK signaling (32), which was reduced in the Siah2KO adipose tissue. A Siah2-dependent mechanism influencing the relationship between adipocyte hypertrophy and insulin sensitivity that indirectly affects inflammation is an attractive possibility given our results showing glucose tolerance and insulin sensitivity were also improved in the lean Siah2KO mice without consistent changes in inflammatory markers.

Future studies that take advantage of adipocyte-specific deletion of Siah2 are needed to fully address the mechanisms by which Siah2 is regulating adipose tissue inflammation, lipid metabolism, and insulin sensitivity in adipose tissue. Nevertheless, our data clearly establishes a role for Siah2 in determining the inflammatory response to adipocyte hypertrophy and regulation of insulin sensitivity in obesity. **O**

© 2015 The Obesity Society

References

- Gregor MF, Hotamisligil GS. Inflammatory mechanisms in obesity. *Annu Rev Immunol* 2011;29:415-445.
- Khan T, Muise ES, Iyengar P, et al. Metabolic dysregulation and adipose tissue fibrosis: role of collagen VI. *Mol Cell Biol* 2009;29:1575-1591.
- Cinti S, Mitchell G, Barbatelli G, et al. Adipocyte death defines macrophage localization and function in adipose tissue of obese mice and humans. *J Lipid Res* 2005;46:2347-2355.
- Qatanani M, Lazar MA. Mechanisms of obesity-associated insulin resistance: many choices on the menu. *Genes Dev* 2007;21:1443-1455.
- Rudich A, Kanety H, Bashan N. Adipose stress-sensing kinases: linking obesity to malfunction. *Trends Endocrinol Metab* 2007;18:291-299.
- Malynn BA, Ma A. Ubiquitin Makes Its Mark on Immune Regulation. *Immunity* 2010;33:843-852.
- Kerscher O, Felberbaum R, Hochstrasser M. Modification of proteins by ubiquitin and ubiquitin-like proteins. *Annu Rev Cell Dev Biol* 2006;22:159-180.
- Molero JC, Jensen TE, Withers PC, et al. c-Cbl-deficient mice have reduced adiposity, higher energy expenditure, and improved peripheral insulin action. *J Clin Invest* 2004;114:1326-1333.
- Marino A, Menghini R, Fabrizi M, et al. ITC deficiency protects from diet-induced obesity. *Diabetes* 2014;63:550-561.
- Kilroy G, Kirk-Ballard H, Carter LE, Floyd ZE. The ubiquitin ligase Siah2 regulates PPAR γ activity in adipocytes. *Endocrinology* 2012;153:1206-1218.
- Tontonoz P, Spiegelman BM. Fat and beyond: the diverse biology of PPAR γ . *Annu Rev Biochem* 2008;77:289-312.
- Frew IJ, Hammond VE, Dickins RA, et al. Generation and analysis of Siah2 mutant mice. *Mol Cell Biol* 2003;23:9150-9161.
- Wong CS, Sceneay J, House CM, et al. Vascular normalization by loss of Siah2 results in increased chemotherapeutic efficacy. *Cancer Res* 2012;72:1694-1704.
- Ayala JE, Samuel VT, Morton GJ, et al. Standard operating procedures for describing and performing metabolic tests of glucose homeostasis in mice. *Dis Model Mech* 2010;3:525-534.
- Rodbell M. Metabolism of isolated fat cells. I. Effects of hormones on glucose metabolism and lipolysis. *J Biol Chem* 1964;239:375-380.
- Baldi P, Long AD. A Bayesian framework for the analysis of microarray expression data: regularized *t*-test and statistical inferences of gene changes. *Bioinformatics* 2001;17:509-519.
- Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Ser B* 1995;57:289-300.
- Donkor J, Sparks LM, Xie H, Smith SR, Reue K. Adipose tissue lipin-1 expression is correlated with peroxisome proliferator-activated receptor alpha gene expression and insulin sensitivity in healthy young men. *J Clin Endocrinol Metab* 2008;93:233-239.
- Tschop MH, Speakman JR, Arch JR, et al. A guide to analysis of mouse energy metabolism. *Nat Methods* 2012;9:57-63.
- Han CY, Subramanian S, Chan CK, et al. Adipocyte-derived serum amyloid A3 and hyaluronan play a role in monocyte recruitment and adhesion. *Diabetes* 2007;56:2260-2273.
- Ma LJ, Mao SL, Taylor KL, et al. Prevention of obesity and insulin resistance in mice lacking plasminogen activator inhibitor 1. *Diabetes* 2004;53:336-346.
- Asterholm IW, McDonald J, Blanchard PG, et al. Lack of "immunological fitness" during fasting in metabolically challenged animals. *J Lipid Res* 2012;53:1254-1267.
- Flier JS, Cook KS, Usher P, Spiegelman BM. Severely impaired adipon expression in genetic and acquired obesity. *Science* 1987;237:405-408.
- Utku N, Heinemann T, Tullius SG, et al. Prevention of acute allograft rejection by antibody targeting of TIRC7, a novel T cell membrane protein. *Immunity* 1998;9:509-518.
- Xu H, Barnes GT, Yang Q, et al. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest* 2003;112:1821-1830.
- Sun K, Tordjman J, Clément K, Scherer PE. Fibrosis and adipose tissue dysfunction. *Cell Metab* 2013;18:470-477.
- Musyoka JN, Liu MC, Pouniotis DS, et al. Siah2-deficient mice show impaired skin wound repair. *Wound Repair Regen* 2013;21:437-447.
- Engelman JA, Berg AH, Lewis RY, Lisanti MP, Scherer PE. Tumor necrosis factor alpha-mediated insulin resistance, but not dedifferentiation, is abrogated by MEK1/2 inhibitors in 3T3-L1 adipocytes. *Mol Endocrinol* 2000;14:1557-1569.
- Guilherme A, Virbasius JV, Puri V, Czech MP. Adipocyte dysfunctions linking obesity to insulin resistance and type 2 diabetes. *Nat Rev Mol Cell Biol* 2008;9:367-377.
- Zhang J, Guenther MG, Carthew RW, Lazar MA. Proteasomal regulation of nuclear receptor corepressor-mediated repression. *Genes Dev* 1998;12:1775-1780.
- Zhao HL, Ueki N, Hayman MJ. The Ski protein negatively regulates Siah2-mediated HDAC3 degradation. *Biochem Biophys Res Commun* 2010;399:623-628.
- Nakayama K, Qi J, Ronai Z. The ubiquitin ligase Siah2 and the hypoxia response. *Mol Cancer Res* 2009;7:443-451.
- Chang L, Kamata H, Solinas G, et al. The E3 ubiquitin ligase itch couples JNK activation to TNF α -induced cell death by inducing c-FLIP(L) turnover. *Cell* 2006;124:601-613.
- Sun K, Kusminski CM, Scherer PE. Adipose tissue remodeling and obesity. *J Clin Invest* 2011;121:2094-2101.
- Strissel KJ, Stancheva Z, Miyoshi H, et al. Adipocyte death, adipose tissue remodeling, and obesity complications. *Diabetes* 2007;56:2910-2918.
- Colonna M. TREMs in the immune system and beyond. *Nat Rev Immunol* 2003;3:445-453.
- Stefanovic-Racic M, Yang X, Turner MS, et al. Dendritic cells promote macrophage infiltration and comprise a substantial proportion of obesity-associated increases in CD11c+ cells in adipose tissue and liver. *Diabetes* 2012;61:2330-2339.
- Guilherme A, Tesz GJ, Guntur KV, Czech MP. Tumor necrosis factor-alpha induces caspase-mediated cleavage of peroxisome proliferator-activated receptor gamma in adipocytes. *J Biol Chem* 2009;284:17082-17091.
- He F, Doucet JA, Stephens JM. Caspase-mediated degradation of PPAR γ proteins in adipocytes. *Obesity* 2008;16:1735-1741.
- Polekhina G, House CM, Traficante N, et al. Siah ubiquitin ligase is structurally related to TRAF and modulates TNF-alpha signaling. *Nat Struct Biol* 2002;9:68-75.