

6-1-2020

Miocene C₄ Grassland Expansion as Recorded by the Indus Fan

Sarah J. Feakins
University of Southern California

Hannah M. Liddy
University of Southern California

Lisa Tauxe
Scripps Institution of Oceanography

Valier Galy
Woods Hole Oceanographic Institution

Xiaojuan Feng
Chinese Academy of Sciences

See next page for additional authors

Follow this and additional works at: https://digitalcommons.lsu.edu/geo_pubs

Recommended Citation

Feakins, S., Liddy, H., Tauxe, L., Galy, V., Feng, X., Tierney, J., Miao, Y., & Warny, S. (2020). Miocene C₄ Grassland Expansion as Recorded by the Indus Fan. *Paleoceanography and Paleoclimatology*, 35(6)
<https://doi.org/10.1029/2020PA003856>

This Article is brought to you for free and open access by the Department of Geology and Geophysics at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Authors

Sarah J. Feakins, Hannah M. Liddy, Lisa Tauxe, Valier Galy, Xiaojuan Feng, Jessica E. Tierney, Yunfa Miao, and Sophie Warny

Miocene C₄ grassland expansion as recorded by the Indus Fan

Sarah J. Feakins^{1*}, Hannah M. Liddy¹, Lisa Tauxe², Valier Galy³, Xiaojuan Feng⁴, Jessica E. Tierney⁵, Yunfa Miao^{6,7}, Sophie Warny⁸

¹Department of Earth Sciences, University of Southern California, Los Angeles, CA, USA,

²Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA,

³Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution,

Woods Hole, MA, USA, ⁴State Key Laboratory of Vegetation and Environmental Change,

Institute of Botany, Chinese Academy of Sciences, Beijing, China, ⁵Department of Geosciences,

University of Arizona, Tucson, AZ, USA, ⁶Key Laboratory of Desert and Desertification,

Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences,

Lanzhou, CN, ⁷CAS Center for Excellence in Tibetan Plateau Earth Sciences and Key

Laboratory of Continental Collision and Plateau Uplift, Institute of Tibetan Plateau Research,

Chinese Academy of Sciences, Beijing, CN, ⁸Department of Geology and Geophysics &

Museum of Natural Science, Louisiana State University, Baton Rouge, LA, USA.

*Corresponding author: Sarah J. Feakins (feakins@usc.edu)

Key Points:

- Multi-proxy study of the Indus Fan organic matter reveals terrestrial sources differ between turbidite and hemipelagic facies.
- Carbon isotopes, grass pollen and charcoal indicate C₄ grasslands expanded between 7.4-7.2 Ma in hemipelagic facies.
- Hydrogen isotopes in plant waxes differ between source regions but do not detect monsoon rainfall changes across the C₄ grassland expansion.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1029/2020PA003856](https://doi.org/10.1029/2020PA003856)

Abstract

In the late Miocene, grasslands spread across the forested floodplains of the Himalayan foreland, but the causes of the ecological transition are still debated. Recent seafloor drilling by the International Ocean Discovery Program (IODP) provides an opportunity to study the transition across a larger region as archived in the Indus sub-marine fan. We present a multi-proxy study of past vegetation change based on analyses of the carbon isotopic composition ($\delta^{13}\text{C}$) of bulk organic carbon, plant wax *n*-alkanes and *n*-alkanoic acids, and quantification of lignin phenols, charcoal and pollen. We analyze the hydrogen isotopic composition (δD) of plant wax to reconstruct precipitation δD . We use the Branched and Isoprenoid Tetraether (BIT) index to diagnose shifts between terrestrial versus marine lipid inputs between turbidite and hemipelagic sediments. We reconstruct ocean temperatures using the TEX₈₆ index only where marine lipids dominate. We find evidence for the late Miocene grassland expansion in both facies, confirming this was a regional ecosystem transformation. Turbidites contain dominantly terrestrial matter from the Indus catchment (D-depleted plant wax), delivered via fluvial transport as shown by the presence of lignin. In contrast, hemipelagic sediments lack lignin and bear D-enriched plant wax consistent with wind-blown inputs from the Indian peninsula; these show a 7.4–7.2 Ma expansion of C₄ grasslands on the Indian sub-continent. Within each facies, we find no clear change in δD values across the late Miocene C₄ expansion, implying consistent distillation of rainfall by monsoon dynamics. Yet, a cooling in the Arabian Sea is coincident with the C₄ expansion.

Plain Language Summary

This project studied the mud and sand on the seafloor in the Indian Ocean, west of India and south of Pakistan. We drilled a core through the mud and focused on a section corresponding to 5.5 to 10 million years ago. Much of the sand came from the Indus River, but when the river sediments went elsewhere, layers of chalky sediments formed from the shells of marine organisms. Those chinks contain the history of longer spans of time. In the sediments, we found molecular fossils from the waxy coating on plant leaves and the woody parts of plants as well as pollen and charcoal; these point to the types of plants that were growing on land whose remains were washed or blown out to sea. We also found molecular fossils of microbes that lived in the oceans, whose structures indicate ocean temperatures – although when soils washed out in large amounts, ocean temperature estimates are not available. The main finding is that grasses replaced forests across much of the Indian subcontinent and Indus catchment, accompanied by more grass fire and cooler ocean temperatures.

1 Introduction

The Siwalik Group in the Himalayan foothills famously documented a one-step late Miocene expansion of C₄ vegetation, beginning at 7.3 Ma and with almost complete C₄ coverage by 6 Ma (Quade and Cerling, 1995; Quade et al., 1989). This >10‰ shift in the carbon isotopic composition ($\delta^{13}\text{C}$) of soil carbonates, has now been replicated in outcrops from Pakistan to Nepal in both soil carbonates and plant wax biomarkers (Freeman and Colarusso, 2001; Ghosh et al., 2017; Vögeli et al., 2017). Further west, in NE Africa, a 6.5‰ rise in the median $\delta^{13}\text{C}$ in fossil teeth enamel of equids reveals a dietary switch, with horses rapidly incorporating C₄ grasses in their diets in the late Miocene (Uno et al., 2011). The timing of the vegetation shift was documented by plant wax records in offshore marine sedimentary archives that indicate a regional expansion of C₄ plants (+4.4‰) soon after 10 Ma (Feakins et al., 2013), later

corroborated across N Africa (Polissar et al., 2019; Uno et al., 2016), with reversals and re-expansions of C₄ plants during the Pliocene (Feakins et al., 2005; Levin, 2015; Liddy et al., 2016). Thus, C₄ vegetation expansion around the Indian Ocean does not generally appear as an abrupt, unidirectional transition.

Early studies suggested that the C₄ grassland expansion in Pakistan required a major climatic shift that would have necessarily included the development of a winter dry season and (perhaps) an increase in summer rainfall (Quade et al., 1989). Subsequently, an increase in the relative abundance of the planktic foraminifer *G. bulloides* in the Arabian Sea in the late Miocene was interpreted to indicate an amplification of wind-driven upwelling and thus the summer monsoon (Kroon et al., 1991). Recently however, that increase in *G. bulloides* has been argued to be an artifact of enhanced carbonate preservation caused by the coincident uplift of the Owen Ridge (Rodriguez et al., 2014) and efforts at replication of *G. bulloides* counts from different locations revealed problematic sensitivity to shell size and bathymetry (Gupta et al., 2015). Furthermore, an increase in cold-tolerant *G. bulloides* could simply reflect a cooling of upwelled waters. This latter interpretation is consistent with evidence for the late Miocene global deep ocean cooling (Holbourn et al., 2013; Holbourn et al., 2018; Lear et al., 2000; Lear et al., 2015). In terms of the surface ocean temperature effects, in the northern Arabian Sea, the late Miocene variability of SSTs is as yet uncertain, as alkenone U^k₃₇ values at Site 722 (Herbert et al., 2016; Huang et al., 2007) hit the upper limit of the paleothermometer and are thus invariant, leaving any change in proximal SSTs associated with the C₄ grassland expansion an open question.

The nature of the C₄ vegetation transition is not well-known south of the Siwalik Group sediments where the grassland expansion was first documented. Expedition 355 of the International Ocean Discovery Program provided a first opportunity to sample Miocene sediments eroded mostly from the lowland Indus Catchment. Although the Indus River now barely flows to the ocean with damming and water use for agriculture, in the past, high sediment loads (Milliman and Syvitski, 1992) built the second largest megafan (Métivier et al., 1999).

In this study, we reconstruct vegetation change as recorded by the late Miocene sediments from IODP Site U1457 (**Figure 1**), collected in 2015 during IODP Expedition 355 to the Indus Fan (Pandey et al., 2015). Late Miocene age sediments comprise voluminous and rapidly emplaced turbidic deposits, and slower accumulation rate hemipelagic sediments. We characterize the provenance of terrestrial vegetation inputs using a combination of inorganic and organic geochemical evidence to differentiate fluvial transport from the Indus River catchment and wind-blown inputs from peninsular India. We seek evidence for vegetation change within each source region based on carbon isotopic composition of bulk organic carbon, plant waxes, and compositional analysis of lignin phenols and pollen assemblages. Drivers of vegetation change that we investigate here include fire (based on measurements of charcoal), hydroclimate (based on hydrogen isotopic analyses of plant waxes) and ocean temperature (based on GDGTs) in order to provide new insights into the regional nature and drivers of the ecological transition.

2 Materials and Methods

2.1 Marine sediments and age model

IODP Expedition 355 drilled Site U1457 (17°9.95'N, 67°55.81'E, 3523 m water depth) in the eastern, distal region of the Indus Fan, ~760 km S of the Indus River mouth (Pakistan) and ~490 km west of India (**Figure 1**). U1457 is located in the western Laxmi Basin, at the foot of the Laxmi Ridge (Pandey et al., 2015). Although the Indus Fan was largely built from N-S

erosion along routes similar to the present Indus River (Bourget et al., 2013; Métivier et al., 1999), drilling at U1457 sampled 190 m of a mass transport deposit (MTD), the Nataraja submarine slide. This event is estimated to have moved 19,000 km³ of sediment, from the continental shelf of peninsular India to the Laxmi Basin (Calvès et al., 2015). Although the triggering mechanism is unknown, the event has been tightly dated to between 9.83–9.69 Ma (Routledge et al., 2019). The present study begins above the MTD (above 880 m) as the MTD lacks the stratigraphic integrity needed for paleoclimate reconstruction.

The late Miocene accumulations above the MTD are mostly turbiditic: silty sands with sharp erosive bases grading upward to clay (Pandey et al., 2015). Recovery is discontinuous (52%) as is common in coarse sediments. Mineralogical analyses indicate that the turbidites, delivered episodically from the shelf, consist mostly of sediments from the southward flowing paleo-Indus River (Pandey et al., 2015). Sediments originating from the Indus River, have been transported through canyons bisecting the continental shelf, delivering turbidity currents into the Indus Fan along channels and building associated levees (Bourget et al., 2013). The locus of deposition has shifted laterally over hundreds of kilometers, and one of these transport channels supplied material to the Laxmi Basin, located in the easternmost region of the fan (Pandey et al., 2015). Such turbidites are known to be efficient mechanisms to deliver (fluvially exported) terrestrial carbon to the deep sea (France-Lanord and Derry, 1997). Additional inputs may derive from westward flowing rivers draining the Indian subcontinent, such as the Narmada and Tapi Rivers (Kessarkar et al., 2003; Yu et al., 2019).

Interspersed between the turbidites is nannofossil ooze, characterized by a mixture of marine carbonates and terrestrial clays. During periods of hemipelagic deposition, eolian influx is likely the dominant conveyor of terrestrial material (Prins et al., 2000). During these periods, Eastern Africa, Arabian Peninsula, Pakistan and peninsular India could all contribute terrestrial material to the Arabian Sea. However modern studies of the eastern marginal seas indicate plant wax from the proximal Indian subcontinent, blown in by winter easterlies (Dahl et al., 2005).

The age model for U1457 is based on nannofossil and foraminiferal datums (Routledge et al., 2019) with refined paleomagnetic datums and age model calculation with error estimation (**Figure 2**) (Tauxe and Feakins, this issue). Ages are on the Geological Time Scale 2012 (GTS 2012) (Gradstein et al., 2012). Depths in this core refer to the core composite depth below seafloor (CCSF) depth scale in meters throughout. In two mostly hemipelagic sections (880–850 m CCSF and 680–620 m CCSF) sufficient age control exists to constrain an age model and associated uncertainty (Tauxe and Feakins, this issue). The limitations of dating turbidic sediments is that there are few available age control points and a highly episodic depositional system. In those sediments age estimation is thus limited to linear interpolation between the constraints in the adjacent hemipelagic units.

There are drastic changes in sedimentation rate associated with the mode of sediment delivery: turbidites have episodic and sometimes voluminous accumulation while hemipelagic sediments have lower and more stable sediment accumulation rates. Between 9.83 and 8 Ma, sediment accumulation rates average ~2 cm kyr⁻¹ including hemipelagic sedimentation with age control from the available magnetostratigraphic constraint of the top of Chron C5n at 859 m (9.79 Ma) and from a nannofossil first appearance datum (FAD) at 832 m (8.12 Ma, FAD *Discoaster quinquerramus*). This hemipelagic interval has five age control datums at an average spacing of 0.3 Ma and appears to be well represented by the linear sedimentation rate, although the dating precision can be no better than 0.3 Ma. In this interval consistent sediment accumulation rates provide a sequence suitable for building a paleoenvironmental archive.

Above this hemipelagic unit, there is a massive accumulation of turbidites. The erosive base (unconformity) above the 8.12 Ma datum (at 832 m, FAD *Discoaster quinqueramus*) is topped by 158 m of sediment (silty sands), with average sediment accumulation rates of $\sim 34 \text{ cm kyr}^{-1}$, up to the bounding constraint at 674 m (7.64 Ma, C4n.1r chron). However, there is no available age control within 158 m of turbidites or 0.5 Ma, and accumulation was via episodic delivery of numerous voluminous transport events whose timing cannot be constrained. Between 7.64 and 5.94 Ma, the latter being the nannofossil last appearance datum (LAD) of *Nicklithus amplificus* at 610 m, sediment accumulation rates are $2\text{--}6 \text{ cm kyr}^{-1}$ including hemipelagic sedimentation above 637 m. This interval has eight age control datums at an average spacing of 0.2 Ma and appears to be well represented by a polynomial fit, with an uncertainty of 0.1–0.5 Ma. It is capped by another accumulation of turbidites, 71 m delivered in 0.4 Ma between the bracketing nannofossil datums (the previous datum and LAD *Discoaster quinqueramus*), although a foraminiferal datum would suggest that it may be even more rapidly emplaced (LAD *Globoquadrina dehiscens*).

We note that uncertainty in the age model comes from both the y axis (depth) and the x axis (age). Age uncertainty in the hemipelagic units comes from the security of magnetostratigraphic constraints and the age control on the nannofossil datums in the many global records from which they derive. Three of the nannofossil datums have assessed uncertainties of 0.2, 0.1 and 0.01 Ma (Routledge et al., 2019), however for other microfossil datums the uncertainty has not been quantitatively assessed. Depth uncertainty derives from the resolution of microfossil sampling, from which the LADs and FADs are identified, with sampling only every 9.5 m (core catcher samples) for foraminifera and nannofossils, with an additional survey at 0.75 m for nannofossils after 8 Ma (Routledge et al., 2019). This means the sampling resolution for FADs and LADs was only every $\sim 0.5 \text{ Ma}$ in the lower hemipelagic unit, but it was every 0.1 to 0.4 Ma within the younger hemipelagic unit in which the main vegetation transition occurs. The two largely hemipelagic intervals with available age control and relatively consistent sediment accumulation rates, provide the best scope for paleoenvironmental reconstruction, and the C₄ transition on the Indian peninsula is constrained to within 0.1 Ma, by by magnetostratigraphy (C3Br.2r) and nannofossil FAD of *Amaurolithus spp.* (Tauxe and Feakins, this issue).

We examined a total of 52 samples from Site U1457 Hole C, at $\sim 7 \text{ m}$ spacing from 500–880 m and $\sim 0.5 \text{ m}$ spacing during the interval of slower (hemipelagic) deposition between 600–670 m. Within the massive turbidites, graded bedding structures were typically 2–15 cm thick thus each sample corresponds to a distinct turbidite flow from shelf to fan. Samples within hemipelagic sedimentation are 2 cm thick and likely integrate $\sim 1 \text{ kyrs}$.

Samples were freeze dried before processing for microfossil counts; and samples for organic analyses were freeze dried then homogenized with a mortar and pestle. Samples were analyzed for a variety of biomarker and geochemical analyses. Epsilon Neodymium (ϵ_{Nd}) values - that is the $^{143}\text{Nd}/^{144}\text{Nd}$ offset from the chondritic uniform reservoir (CHUR), in parts per 10,000 - were previously used to study the provenance of Indus Fan detrital material (Clift et al., 2019). Those ϵ_{Nd} data measured at paired depths with the biomarkers, are reported on the new age model here.

2.2 Lignin phenol quantification

For CuO oxidation to release lignin-derived phenols, powdered samples were placed in Teflon-lined containers together with 0.5 g CuO, 100 mg ammonium iron (II) sulfate

hexahydrate, and 20 mL of 12 M NaOH under N₂ and heated at 170°C for 2.5 h. Then, the water phase was acidified to pH <2 with 6 M HCl and kept for 1 h at room temperature in the dark to prevent reactions of cinnamic acids. After centrifugation (2500 rpm for 30 min), the supernatant was liquid-liquid extracted with ethyl acetate. Extracts were rotary evaporated, spiked with an internal quantification standard (ethyl vanillin), and derivatized with N,O-bis-(trimethylsilyl) trifluoroacetamide and pyridine (70°C, 1 h) to yield trimethylsilyl derivatives before quantification by gas chromatograph (GC) mass spectrometry (MS). We quantified lignin phenols using internal standards on a Trace 1310 GC coupled to an ISQ MS (Thermo Fisher Scientific) using a DB-5ms column (30 m × 0.25 mm, film thickness 0.25 µm). The GC oven temperature was held at 65°C for 2 min, increased from 65–300°C at a rate of 6°C min⁻¹, and held for 20 min. Helium was used as carrier gas (0.8 mL min⁻¹). The MS was operated in electron impact mode at 70 eV and scanned from 50–650 daltons. External quantification standards were used to normalize the response factor for different lignin phenols separately. Vanillyl (vanillin, acetovanillone, vanillic acid), syringyl (syringaldehyde, acetosyringone, syringic acid), and cinnamyl (*p*-coumaric acid, ferulic acid) phenols were summed to represent lignin phenols (VSC).

2.3 Bulk organic carbon isotopic analyses

For bulk organic carbon (OC) measurements, powdered samples were weighed into methanol-rinsed Ag capsules and acidified in a vacuum desiccator at 0.5 atm with an open dish with 50 mL 12N HCl, at 62°C for 60 to 72 h (Whiteside et al., 2011), before drying in a vacuum desiccator with indicating silica gel. Samples were packaged in Sn capsules prior to analysis by a Carlo Erba/Fisons 1108 flash elemental analyzer (EA) equipped with a Costech ZeroBlank carousel, coupled to a DeltaPlus isotope ratio MS via a Finnigan MAT ConFlo II interface for both OC quantification and carbon isotopic analyses ($\delta^{13}\text{C}_{\text{OC}}$; ‰ Vienna Pee Dee Belemnite, VPDB). A blank and three standards (IAEA-N1 [Ammonium sulfate], USGS-40 [Glutamic Acid], and glycine) were run between every 16 samples, and indicate a total OC uncertainty (precision and accuracy) of better than 4%. NBS-19 (limestone) and a calcite laboratory standard were run as unknowns for additional verification of consistency. Carbon isotopic compositions of standards were accurate and precise to better than 0.3‰. N% was measured on non-acidified aliquots and used to calculate OC/N.

2.4 Lipid extraction and biomarker purification

Powdered sediment samples (31.7–95.1 gdw) were extracted with an Accelerated Solvent Extraction system (ASE 350®, Dionex) with 9:1 ratio of dichloromethane (DCM): methanol (MeOH) at 100°C and 1500 psi for two 15-minute cycles. Extracts were separated over a NH₂ sepra column (5 cm x 40 mm Pasteur pipette, 60 Å) eluted with 2:1 ratio of DCM: isopropanol (neutral fraction) and 4% formic acid in diethyl ether (acid fraction). The acid fraction (containing fatty acids) was methylated with methanol of known isotopic composition with 95:5 MeOH:HCl at 70°C for 12 h, before the fatty acid methyl esters (FAMES) were partitioned into hexane and passed through anhydrous sodium sulfate. Samples were further purified over a silica gel column (5 cm x 40 mm Pasteur pipette, 5% water-deactivated silica gel, 100–200 mesh), washing first with hexane, and then eluting the FAME fraction with DCM. The neutral fraction was loaded onto a silica gel column (5 cm x 40 mm Pasteur pipette, 5% water-deactivated silica gel, 100–200 mesh) and eluted with hexane (*n*-alkanes), DCM and MeOH (GDGTs). S was

removed from the *n*-alkanes by eluting in hexane over activated copper. Further purification included urea adduction when necessary for the *n*-alkanes.

2.5 GDGT quantification

The fractions containing glycerol dialkyl glycerol tetraethers (GDGTs) were dissolved in hexane:isopropanol (99:1) and filtered (0.45 µm PTFE) prior to injection on an Agilent 1260 High-Performance Liquid Chromatography (HPLC) coupled to an Agilent 6120 MS. GDGTs were analyzed according to the method of Schouten et al. (2007b), using a Grace Cyano column with 5 min isocratic elution using A, and then a gradient of 90% A/10% B to 82% A/18% B over 35 minutes, where A = hexane and B = hexane:isopropanol (9:1). The column was then backflushed with 100% B for 10 min, and re-equilibrated with 90% A/10% B for another 10 min. Single Ion Monitoring (SIM) mode of the $M + H^+$ ions was used to detect and quantify the isoprenoidal GDGTs produced by marine archaea including GDGTs with 0-3 cyclopentane moieties (GDGT-0 to GDGT-3); crenarchaeol (Cren) with an additional cyclohexane moiety and its regioisomer crenarchaeol' (Cren'). Following Schouten et al. (2007a), TEX₈₆ units were calculated using the equation:

$$TEX_{86} = \frac{[GDGT-2] + [GDGT-3] + [Cren']}{[GDGT-1] + [GDGT-2] + [GDGT-3] + [Cren']}$$

Repeat analysis of a laboratory standard yields a long-term precision of 0.004 TEX₈₆ units. We converted the TEX₈₆ record to mean annual sea surface temperatures (SSTs) using the BAYSPAR calibration (Tierney and Tingley, 2014). For the prior distribution, we used the mean SSTs at the latitude and longitude grid cell (27.6°C) and a standard deviation of 10°C (Locarnini et al., 2010). We report total concentrations of branched (brGDGT) and isoprenoidal (isoGDGTs) GDGTs (ng gdw⁻¹) as well as the Branched and Isoprenoidal Tetraether (BIT) index:

$$BIT = \frac{I + II + III}{I + II + III + IV}$$

where I, II and III represent the abundances of brGDGT and IV represents the abundance of crenarchaeol in each sample (Hopmans et al., 2004). TEX₈₆ values in samples with a BIT > 0.4 were considered to be compromised by terrestrial material and thus unreliable for SST estimation.

2.6 Compound specific carbon and hydrogen isotopic analyses

The $\delta^{13}C$ and δD isotopic composition of *n*-alkanoic acid and *n*-alkanes were analyzed using a Thermo Scientific Trace GC equipped with a Rxi-5ms column (30 m × 0.25 mm, film thickness 0.25 µm) with a PTV injector operated in solvent-split mode, coupled to a Delta V Plus isotope ratio mass spectrometer (IRMS) via an Isolink combustion/pyrolysis furnace (1000/1400°C). Isotopic linearity was monitored daily across a range of peak amplitude (1-8V) CO₂/H₂ gas pulses. For carbon, the isotopic linearity of $\delta^{13}C$ averaged 0.05‰. For hydrogen, the H₃ factor averaged 4.6 ppm mV⁻¹. Two out of five CO₂/H₂ reference peaks were used for standardization of the isotopic analysis, the remainder track stability. We used external standard runs containing a mixture of 15 *n*-alkanes (C₁₆ to C₃₀) with $\delta^{13}C$ values ranging from -33.3 to -26.2‰ and δD values ranging from -254.1 to -9.1‰ (A mix; supplied by A. Schimmelmann,

Indiana University, USA) to normalize to the VPBD-LSVEC carbon isotopic scale and to the Vienna Standard Mean Ocean Water (VSMOW) – Standard Light Antarctic Precipitation (SLAP) hydrogen isotopic scale respectively. The RMS error of the external standards were better than 4.5‰ for δD and 0.2‰ for $\delta^{13}C$. Samples were run in triplicate, and the average standard deviation was 0.5‰ for carbon and 2.2‰ for hydrogen. The isotopic composition of C added during methylation of *n*-alkanoic acids was determined by offline combustion and dual-inlet IRMS ($\delta^{13}C$ MeOH = $-24.7 \pm 0.2\%$, $n=6$), and the isotopic composition of H added during methylation was determined by the esterification of phthalic acid for analysis by GC-IRMS (δD MeOH = $-187 \pm 4\%$, $n=6$). The addition of the methyl group to *n*-alkanoic acids was corrected for by mass balance. The results are reported using delta notation ($\delta^{13}C$ and δD , ‰).

2.7 Palynological counts: pollen and microcharcoal

For palynology, samples were processed using standard chemical palynological processing techniques to extract both pollen and microcharcoal particles. Dry sediment was weighed and spiked with a known quantity of *Lycopodium* spores to allow for evaluation of palynomorph concentrations. Dry sediment was successively treated with hydrochloric acid, hydrofluoric acid, and heavy liquid separation (Brown, 2008). Samples were sieved between a 10 and 250 μm fraction, and the remaining residue was mounted on microscope slides using glycerin jelly. Palynological analysis was conducted on 16 samples in the Louisiana State University's Center for Excellence in Palynology (CENEX) lab. When possible, 300 palynomorphs were tabulated per sample using an Olympus BX41 microscope. Palynomorphs were identified to the lowest taxonomic level possible, and concentrations were calculated as:

$$C = (P_c \times L_t \times T) / (L_c \times W)$$

where C = concentration (per gram of dried sediment, gdw^{-1}), P_c = the number of palynomorphs counted, L_t = the number of *Lycopodium* spores per tablet, T = the total number of *Lycopodium* tablets added per sample, L_c = the number of *Lycopodium* spores counted, W = the weight of dried sediment. Palynomorph data collection included counts of pollen from terrestrial plants including the Poaceae (grasses) reported as a % of all terrestrial pollen (grass pollen %), as well as ratios (non-aboreal:arboreal; NAP %) and preservation (reworked %).

Microcharcoal was studied in 31 samples. Two basic microcharcoal shapes, long (L) and round (R), were identified from the ratio of length (major axis)/width (minor axis), where $L > 2.5$ and $R < 2.5$, where high L/R ratios indicate more grass fire (Miao et al., 2019). Ratios of big/small charcoal were reported, where big ($>50 \mu m$) and small ($<50 \mu m$), with larger sizes a semi-quantitative indicator of fire proximity, conversely small fragments may be transported further or may experience fragmentation during transport. Total charcoal abundance is normalized to organic carbon concentration, to account for sediment dilution effects, in order to assess proportional inputs of charcoal which may reflect fire frequency and magnitude.

3 Results and Discussion

3.1 Stratigraphic context and multi-proxy evidence for provenance of biomarkers

Although the turbidites dominate volumetrically, remobilized material is mixed and exported in geologically brief interludes. Rapid emplacement of turbidites, means that marine influence is minimal and the components are almost entirely terrigenous. In contrast,

hemipelagic facies can be sampled with finer temporal resolution despite slower accumulation rates. Although marine productivity dominates, the slow accumulations also archive eolian inputs of terrigenous material. Turbidite and hemipelagic facies are characterized by different inorganic and organic components reflecting their different mode of emplacement and provenance. We find their different characteristics can be described by four proxies in particular: ϵ_{Nd} , BIT, lignin phenols and plant wax δD (**Figure 3**).

Nd analyses were previously reported for a subset of the samples analyzed here, and Clift *et al.* [2019] reported that turbidites in sediments of U1457 of Neogene age (**Figure 3a**, brown shading) were derived from the Indus River catchment. The Neogene values (ϵ_{Nd} values of -12 to -9 , especially the upper turbidites with ϵ_{Nd} values of -12 to -11), overlap with the Indus River export during the Holocene (ϵ_{Nd} -12 to -10) (Clift *et al.*, 2010), whereas the Pleistocene turbidites are also Indus-derived but more depleted (ϵ_{Nd} -14 to -13) (Clift *et al.*, 2019; Yu *et al.*, 2019). Hemipelagic sediments have ϵ_{Nd} ~ -9.5 to -6.5 and together with other evidence including Sr and clay mineralogy this points to a peninsular India (radiogenic Deccan) source (Clift *et al.*, 2019; Yu *et al.*, 2019) (**Figure 3**, blue shading), with greater variability consistent with variable sourcing and less mixing in comparison to the turbidites.

The BIT index records clear changes in the proportion of marine vs. terrestrial organic inputs with high marine productivity ($\text{BIT} < 0.2$) in hemipelagic sediments and $\text{BIT} > 0.4$ in turbidites indicating a dominance of fluvially-transported, soil-derived membrane lipids (**Figure 3b**). As our focus is the organic component of sediments, we select the BIT proxy with a cutoff of 0.4, to assign sourcing of the biomarkers. Sedimentologic description of the corresponding core section intervals (based on high resolution core section images) confirm that this cutoff adequately separates turbiditic and (usually bioturbated) hemipelagic facies.

High lignin phenol content of turbidites confirms fluvial transport of woody plant components. The low lignin phenol content of the hemipelagic layers (**Figure 3c**) likely indicates negligible fluvial inputs and instead suggests that any terrigenous plant debris (plant wax, pollen and charcoal) was likely wind-blown. Based on Holocene plant wax (Dahl *et al.*, 2005) and Nd evidence for radiogenic, likely Deccan influences (Clift *et al.*, 2019; Yu *et al.*, 2019), we assume a proximal Indian peninsula source for the terrigenous plant debris within Neogene hemipelagic sediments. Studies of Sr-Nd and clay mineralogy have ruled out an Arabian source of dust during the Pleistocene (Yu *et al.*, 2019), as have studies of plant wax provenance in the Holocene, instead finding Indian peninsula sourcing via low level wind transport in winter easterlies and in the upper troposphere easterlies in summer (Dahl *et al.*, 2005).

Plant wax δD values show abrupt transitions between D-depleted values in the turbidites (-220 to -180‰) and D-enriched values in the hemipelagic sediments (-190 to -130‰ , **Figure 3d**). This is consistent with the BIT evidence for facies changes, the lignin evidence for transport mechanism, and Nd evidence for source switching, between Indus River-derived plant waxes in the turbidites (D-depleted) and those that are wind-blown from India into the hemipelagic units.

3.2 Multi-proxy evidence for C₄ grassland expansion

We have generated a multi-proxy record of vegetation change based on carbon isotopes and palynomorphs (pollen, microcharcoal), and parsed these using the BIT index to distinguish between facies and thus separate archives of terrestrial vegetation in the Indus catchment (**Figure 4a**; turbidites) and peninsular India (**Figure 4b**; hemipelagic), following source characterization described in Section 3.1. We report $\delta^{13}\text{C}$ values of long chain *n*-alkane (C_{31}) and *n*-alkanoic acid (C_{30}) homologues, both presumed to derive from the waxy coating on plant leaves, as well as for

bulk organic matter ($\delta^{13}\text{C}_{\text{OC}}$) that comprises mostly terrestrial material in turbidites and a mixture of marine and terrestrial production in the hemipelagic facies.

Owing to their high sedimentation rates, turbidites provide only brief glimpses of Indus River catchment conditions at 8.2–7.6 Ma and ~7.4 Ma (grouped as 8.2–7.4 Ma), and 5.9–5.6 Ma (**Figure 4a**), with no internal age control. Nonetheless, the turbidites do allow us to contrast conditions in the Indus catchment before and after the main C_4 grassland transition: all proxies ($\delta^{13}\text{C}_{\text{OC}}$, $\delta^{13}\text{C}_{31\text{alk}}$, $\delta^{13}\text{C}_{30\text{acid}}$, grass pollen and L/R charcoal) indicate C_4 grass components increased between 7.4 and 5.9 Ma, suggesting an expansion of C_4 grasslands across the Indus River catchment. Comparing the turbidites bracketing the C_4 grassland expansion, we find a shift from -23.8‰ ($\pm 0.5\text{‰}$, $n = 13$) to -20.8‰ ($\pm 1.2\text{‰}$, $n = 9$) in $\delta^{13}\text{C}_{\text{OC}}$, from -32.1‰ ($\pm 0.5\text{‰}$, $n = 10$) to -26.2‰ ($\pm 0.3\text{‰}$, $n = 7$) in C_{31} *n*-alkanes, from -28.9‰ ($\pm 1.7\text{‰}$, $n = 12$) to -21.7‰ ($\pm 1.2\text{‰}$, $n = 7$) in C_{30} *n*-alkanoic acids. Although a carbon isotopic shift is observed in bulk OC as well as the plant wax components, the relative ^{13}C -enrichment is muted in the bulk, and the absolute values of the OC are ^{13}C -enriched as expected based on the isotopic offset in plants between bulk plants and plant wax and as would be the case if admixed with marine contributions. The OC/N (mean 6.0 ± 1.3 , $n = 21$) indicates some marine inputs to the bulk OC emphasizing the need for compound specific approaches to isolate terrestrial organic matter from that of marine inputs. However, the two plant wax compound classes reveal some differences. Before 7.4 Ma the *n*-alkanes ($-32.1\text{‰} \pm 0.5\text{‰}$, $n = 10$) suggest a pure C_3 ecosystem, such as woodlands, whereas the *n*-alkanoic acids show more variability ($-28.9\text{‰} \pm 1.7\text{‰}$, $n = 12$) that could indicate some C_4 contributions in the 8.2–7.4 Ma turbidites. This offset may reflect differential production of the two compound classes in plants, as there are much greater concentrations of C_{29} and C_{31} *n*-alkanes on the leaves of tropical trees compared to grasses (Garcin et al., 2014), such that C_3 trees may dominate these *n*-alkanes, as also reported in the Mississippi River catchment (Suh et al., 2019). Consistent with this understanding, the C_{33} *n*-alkanes are enriched by up to 1.9‰ (**Supplemental Information, Figure S5a**), indicating less ‘dampening’ of the grass signal by trees. The C_{30} *n*-alkanoic acid appears to be even more sensitive to the C_4 grassland expansion, with two tentative explanations: leaf production or catchment sourcing. There is little information on relative concentrations of *n*-alkanoic acids produced by grasses, however tropical trees do not produce have a strongly modal distribution of *n*-alkanoic acids as they do with *n*-alkanes (Feakins et al., 2016), such that *n*-alkanoic acids are unlikely to be biased to tree production. Other differences between compound class could arise during fluvial export. Observations from modern catchments that have found that *n*-alkanoic acids tend to have a bias to lowland proximal sources greater than the lowland area would suggest, whereas the *n*-alkanes may retain signatures that are more distally-sourced within the catchment arising from inland or higher elevation erosion (Feakins et al., 2018; Hemingway et al., 2016).

In the Indus River system, we may therefore expect more signs of the upland C_3 forested flanks of the Himalayas in the *n*-alkanes record even after the lowland transitioned to C_4 grassland. Alternatively, isotopic offsets between compound classes could result from non-plant, marine algal or sedimentary microbial input of *n*-alkanoic acids (Kusch et al., 2010; Makou et al., 2018), which could raise the $\delta^{13}\text{C}$ values. However why an increase in the microbial contribution would coincide with the vegetation change is unknown and seems unlikely unless this were related to a change in catchment soils and sediment dynamics with the ecological transition. Microfossil evidence confirms the C_4 grassland expansion identified in the bulk organic and plant wax biomarker carbon isotopic evidence. We find that pollen confirms an

increase in *Poaceae* (grasses) from 16% ($\pm 18\%$, $n = 3$) to 59% ($n = 1$) and grass-derived charcoal (L/R) from 21% (± 55 , $n = 8$) to 73% (± 155 , $n = 8$) between the turbidite sets ending at 7.4 Ma and the next set beginning at 5.9 Ma (**Figure 4a**) indicates an increase in grass fire. Although it is challenging to convert carbon isotopes to $C_4\%$ given the spread in C_3 plants associated with canopy closure resulting in non-unique interpretations, the microfossil evidence supports the n -alkanoic acid evidence and allows for roughly a quarter of the lowland biosphere being C_4 grasses or shrubs in the Indus catchment at 7.4 Ma, supporting the interpretation of an early rise of C_4 detected in the n -alkanoic acids as a signal of grassland expansion.

In the hemipelagic sediments, we are able to develop a vegetation reconstruction for peninsular India spanning 9.9 to 5.9 Ma. In the earliest part of the record, plant wax carbon isotopic values are close to -30% suggesting C_3 ecosystems, and pollen data indicate a mixture of conifers (which grow in montane zones today), Amaranthaceae (subshrubs), grasses and mangroves suggesting broad sourcing and mixing from mountain to delta. Each of the carbon isotope proxies shows some enrichment around or after 9 Ma. $\delta^{13}C_{OC}$ values show a gradual modest increase after 9 Ma, and the $\delta^{13}C_{30}$ n -alkanoic acids show a rise at 8.7 Ma with a single enriched sample (and a large amount of non-arboreal pollen) before a reversal. The main C_4 expansion is complete before 7.2 Ma. $\delta^{13}C_{OC}$ values increase in one main step from -20.9 to -18.8% between 7.3 and 7.2 Ma. The $\delta^{13}C_{30}$ n -alkanoic acid shifts from -29.3% at 7.4 Ma to -23.7% at 7.2 Ma. The $\delta^{13}C$ of the C_{31} n -alkanes increase from -30.2 to -26.8% from 7.4 to 7.2 Ma indicating an abrupt ^{13}C -enrichment. Correspondence between the carbon isotopic composition of the different proxies reveals their common source in terrestrial vegetation. Subtle differences between the three carbon isotopic archives reflects sensitivity to productivity and degradation pathways that if understood, may reveal useful details. The C_{30} n -alkanoic acid record indicates a stronger C_4 expansion than the C_{31} n -alkanes in these hemipelagic sediments, as in the turbidites. Evidence from microfossils also informs on the grassland expansion in the hemipelagic sediments. The variability in pollen between samples implies heterogeneity in the wind-exported signal, which may explain divergence in the two compound classes of plant waxes carbon isotope proxies. *Poaceae* (grass) pollen increase from 8% at 7.5 Ma to 47% at 7.4 Ma, and reaching 77% at 7.2 Ma. Thus pollen are highly complementary to the carbon isotope evidence corroborating the timing of the main C_4 grass rise enveloping the shift in all three archives. Long/round charcoal increases, from $<50\%$ before the C_4 transition to $>50\%$ after the transition (**Figure 4b**), reflecting a proportional increase in grasses in combustion (Miao et al., 2019), as would be expected with fire accompanying grassland expansion.

Each of the proxies records the C_4 transition and the subtle differences reveal the challenges of quantitatively establishing the proportion of grassland on the landscape from each of the proxies. As the stronger signals of ^{13}C -enrichment in the n -alkanoic acids versus n -alkanes are derived from different transport pathways (fluvial vs. aeolian) and different source regions (Indus catchment vs. peninsular India) in the two facies, this suggests that the difference in proxy sensitivity originates from the plant wax production in tree leaves swamping the C_{31} n -alkane record and underrepresenting the C_4 expansion, compared to the stronger signal of C_4 expansion seen in the C_{30} n -alkanoic acids and the microfossil evidence (pollen and grass-derived charcoal).

3.3 Did fire drive the C_4 grassland expansion?

As C_4 grasslands expanded in both regions (**Figure 5a**), the increased charcoal abundance (**Supplementary Information**) suggests that fire promoted savanna grassland expansion and that fire increased as a result of the presence of grasslands (grasslands are more

flammable than forests). Other studies have similarly used various measures of fire including charcoal and polycyclic aromatic hydrocarbons (PAHs) to show an increase in fire as grasslands expand, including during the late Miocene C₄ expansion documented in the Siwaliks (Karp et al., 2018) and in the Pliocene in West Africa (Hoetzel et al., 2013). Here we use the shape of charcoal (Miao et al., 2019), to identify the proportion of grass-derived charcoal (increased L/R, **Figure 5b**). The increased prevalence of grass fires provides evidence of the feedback role of fire in maintaining grasslands (Beckage and Ellingwood, 2008), as has been shown in modern climates with bistable ecosystems in Africa and South America (Dantas et al., 2016; Staver et al., 2011). Here too in the Indus catchment and peninsula India, it appears likely that fires maintained savanna grassland ecosystems once they appeared, as the transition appears abrupt and unidirectional in the late Miocene, yet in the modern ecosystems C₄ grasses are not dominant, suggesting bistability, at least between forest-grassland in peninsular India (Das et al., 2015), whereas lowland Indus is too dry for grasslands, instead supporting xeric taxa in the late Holocene (Ivory and Lezine, 2009). Modern studies indicate that it is low dry season precipitation amounts that is particularly critical in allowing fire to spread (Das et al., 2015) with the ignition source being lightning that itself exhibits threshold behavior and is climate dependent (Beckage and Ellingwood, 2008). The fire feedback effect may have been enhanced in the Miocene if oxygen levels were elevated through carbon burial as has been hypothesized (Derry and France-Lanord, 1996), helping to maintain C₄ grasslands after they appeared, although no such oxygen evidence is available. An alternate disturbance mechanism comes from cyclonic activity, which can uproot large trees and promote grassland replacement (Beckage and Ellingwood, 2008), with some evidence that Arabian Sea cyclonic activity is increasing associated with anthropogenic warming (Murakami et al., 2017), perhaps also during Miocene warmth.

3.4 Did a change in monsoon rainfall drive the C₄ grassland expansion?

In this study, we sought to test the long-standing question as to whether a change in monsoon precipitation drove the C₄ expansion, using the hydrogen isotopic composition as recorded by plant wax biomarkers as a proxy for precipitation. Within the turbidites, the δD values of C₃₁ *n*-alkanes vary between -213 to -198‰ (mean $-206 \pm 8\text{‰}$, $n = 5$, s.e.m. 4‰) and for the C₃₀ *n*-alkanoic acids between -205 to -166‰ (mean $-191 \pm 12\text{‰}$, $n = 13$, s.e.m. 4‰). Within the hemipelagic sediments, the δD values of C₃₁ *n*-alkanes varies between -190 to -162‰ (mean $-175 \pm 9\text{‰}$, $n = 16$, s.e.m. 2‰) and for the C₃₀ *n*-alkanoic acids between -160 to -139‰ (mean $-148 \pm 7\text{‰}$, $n = 8$, s.e.m. 2‰). Thus the plant waxes in the turbidites are distinctly more D-depleted than the hemipelagic sediments, with no overlapping range. This strongly suggests the different plant wax sourcing of these facies determines their isotopic composition.

In order to calculate estimates of precipitation δD we select a fractionation for tropical forests (Feakins et al., 2016), which is close to the global average value (Sachse et al., 2012), but carries the benefit of calibration of both compound classes used here, and surveyed in a large dataset. Using an $\epsilon_{\text{alkane/precip}}$ of -129‰ and $\epsilon_{\text{acid/precip}}$ of -121‰ (Feakins et al., 2016) for tropical trees, the central estimates for δD_{precip} are -90‰ based on *n*-alkanes and -80‰ based on *n*-alkanoic acids for sources in the Indus catchment (**Figure 5c**). The compound class offset fits with the hypothesis that *n*-alkanes may integrate more of the catchment and that *n*-alkanoic acids may be dominantly lowland derived (Hemingway et al., 2016). Uncertainties in the appropriate fractionation (which have not been determined for this region's vegetation), are likely on the order of $>10\text{‰}$. Reconstructed values for the Indus River turbidites are much more D-depleted

than modern coastal precipitation sampled by the Global Network of Isotopes in Precipitation (GNIP), in Karachi, Pakistan (IAEA/WMO, 2019) and likely much of the lowland catchment. Indus River turbidites are also more D-depleted than the summer-season precipitation in the NE Indus catchment at the GNIP station in Jammu, Kashmir. Although D-depleted precipitation may occur with either winter or summer rainfall maxima during individual storms, observational data indicate that monthly D-depletion is greatest for the highly distilled westerlies and cold-season precipitation (JF monthly mean values $<-60\%$ at Kabul, Afghanistan) influencing the NW Indus catchment. Summer rainfall in the lowlands is less well constrained by Global Network of Isotopes in Precipitation (GNIP) data, but would likely be between -50% (signaling deep convection) and $+50\%$ (evaporation during descent) based on examination of monthly values for Karachi and very limited data available for Mumbai (IAEA/WMO, 2019). The turbidites instead contain plant wax signals much more consistent with modern winter-dominated precipitation in locations similar to Kabul, Afghanistan (see **Supplementary Information** for modern climate maps and precipitation isotope data).

Although we are not aware of any modern or Holocene estimates of plant wax δD values transported by the Indus River or exported from the Indus catchment, comparison to processes in the Ganges-Brahmaputra system (Galy et al., 2011) and other tropical rivers (Feakins et al., 2018; Hemingway et al., 2016) demonstrated export of a minor upland (distal) component mixed with a dominantly lowland (proximal) signature, and thus a lowland (proximal) bias in the exported signal. Yet, the observed Indus River turbidite δD values are D-depleted compared to expected precipitation in the lowland reaches. Thus it is possible that these late Miocene Indus River exported signatures of plant wax δD in the turbidite facies are more depleted than we would expect for the modern situation, perhaps implying more winter season precipitation (like Kabul, Afghanistan) within a greater portion of the catchment. Alternatively, it is plausible that the Indus catchment may see a greater proportion of upland signal exported from forested regions, given the catchment has a much drier lowland than the Ganges, Congo or Amazon, with reduced productivity in the lowland grasslands and shrublands. The hypothesis that the Indus River may export the upland biomarker signal to the sea, could be tested with future work in modern river or Holocene delta sediments in the Indus River system or in other large montane river networks with dry lowlands.

Given the vegetation transition it may be appropriate to consider selection of a smaller fractionation to represent arid C_3 ecosystems (Feakins and Sessions, 2010; Feakins et al., 2014) and a larger fractionation to represent C_4 grasslands (Smith and Freeman, 2006) however as there are no locally available calibrations we opt for a fixed fractionation for our reconstructions. We note that if a smaller epsilon were appropriate (as for dryland trees perhaps relevant to the Holocene Indus) then this would make the precipitation more D-depleted (by up to 30%) (Feakins and Sessions, 2010), pointing further to winter precipitation or upland sourcing in that case. A larger fractionation for grasses after the C_4 expansion would increase the offset in precipitation reconstructed between the two turbidites (**Figure 5c**), and could imply a drying, or more lowland source. However, any such drying would imply the tree component would shift from moist to dry forest values and the overall plant community average may have very little overall change in fractionation, thus we do not apply a correction given the uncertainties. We further emphasize that all interpretations must remain cautious due to very limited δD data for the turbidites.

Within the hemipelagic sediments, applying the tropical tree fractionation implies a δD_{precip} of -78 to -47% based on n -alkanes or -44 to -20% based on n -alkanoic acids for

sources in peninsular India (**Figure 5c**), which overlaps with the modern range of precipitation isotopes during the Indian Summer Monsoon (**Supplementary Information**) and indicates similar to modern or perhaps wetter conditions during the late Miocene, with similar circulation and distillation patterns over the Indian peninsula, consistent with evidence for topographic steering and orographic rainout patterns being in place by this time (Acosta and Huber, 2020). The wet tropical forest fractionation appears appropriate for the Western Ghats today and would likewise be appropriate before the C₄ transition. Once again, we do not apply a vegetation correction for possible hydrogen isotope fractionation changes after C₄ expansion, as the difference between tropical C₃ trees and tropical C₄ grasses is not large. Using the slightly larger $\epsilon_{\text{alkane/precip}}$ fractionation for C₄ grasses of -134‰ (Sachse et al., 2012) would adjust the data points after the C₄ transition by $+15\text{‰}$, within the spread of data, and thus would not alter interpretations for the lack of detectable hydroclimate change in peninsular India.

Although we find a large hydrogen isotopic range in U1457 (**Figure 5c**), this is largely controlled by the change in provenance between facies. The fluvially-exported plant waxes in the turbidites (accompanied by river transported lignin, **Figure 5d**; and soil derived lipids, with a high BIT, **Figure 5e**) carry an isotopic composition that is considerably more D-depleted than the wind-blown plant waxes from peninsular India in the hemipelagic sediments (characterized by almost no lignin, **Figure 5d**; and marine-dominated lipids, with a low BIT, **Figure 5e**). Although our δD data are sparse, we find no systematic shift in δD within each facies within the available data. Importantly, we find negligible change in the hydrogen isotopic composition of plant waxes in either archive before and after the C₄ transition at 7.2 Ma in peninsular India and between bracketing archives at 7.4 and 5.9 Ma for the Indus catchment.

This lack of plant wax δD shift associated with a plant wax $\delta^{13}\text{C}$ shift is also seen in records from West Africa at ~ 10 Ma (Polissar et al., 2019), however revisiting their data from DSDP Sites 235 and 241, offshore Somali, do in fact show a 20‰ δD shift associated with the C₄ expansion. Their data are therefore similar to a prior report of a ~ 10 Ma C₄ expansion (Feakins et al., 2013), accompanied by a 40‰ δD shift, that was interpreted as a drying (Feakins, 2013), from DSDP Site 231 in the Gulf of Aden. Thus the absence of a hydrogen isotope shift at the C₄ transition in U1457 is similar to reports from West Africa but differs from the pattern observed in NE Africa. This may denote variations in the degree to which hydroclimate change accompanies the C₄ transition in different regions (Higgins and Scheiter, 2012). Yet, each of these studies (Feakins, 2013; Polissar et al., 2019; and this study), has been conducted at very low resolutions (as low as every ~ 1 Ma), and future work could usefully seek to probe Miocene hydroclimate variability with higher resolution plant wax δD studies. And, as the sensitivity of precipitation isotopes to detect hydroclimate transitions varies between climate regimes and regions, the question merits testing with additional proxies for hydroclimate.

3.33 Did late Miocene cooling drive the C₄ grassland expansion?

Another driver of the ecological transition has been proposed: related to a global cooling in the late Miocene (Herbert et al., 2016). Here we generate new data for ocean temperatures to test this hypothesis. The turbidite facies have high BIT values (>0.4) which indicates that soil inputs of GDGTs dominate, compromising the TEX₈₆ thermometer (**Figure 5e**). In the hemipelagic sediments however, the BIT values were low (mostly <0.2) indicating mainly a marine source of GDGTs, thus TEX₈₆ based temperature estimates were converted using the BAYSPAR calibration (**Figure 5f**). Ocean temperature estimates have central values of 28 to 24°C. The absolute values of the reconstructed ocean temperatures have uncertainties that are

rigorously quantified and that are on the order of $\sim 4^{\circ}\text{C}$. Previous work in the Arabian sea and Gulf of Aden suggests that TEX₈₆ may represent sub-surface rather than surface waters (Liddy et al., 2016; Tierney et al., 2016). Given these uncertainties more emphasis should be placed on the trends in the temperature data and we see a cooling of 4°C across 10–5.5 Ma, with a 1°C step shift close to 7.3 Ma (**Figure 5f**). This differs from a nearby alkenone record which remained at the upper limit of that paleothermometer at nearby ODP Site 722 throughout this span (Huang et al., 2007). Although the absolute temperatures may be underestimates, the cooling trend is likely robust and detects a late Miocene cooling of surface waters, such as via increased upwelling in the Oman margin or cooling of the deep ocean and thus cooling of the upwelled water mass. The finding of a late Miocene cooling trend here is furthermore consistent with reports of a global late Miocene cooling in an analysis of available alkenone-based sea surface temperature (SST) estimates (Herbert et al., 2016), and in foraminiferal Mg/Ca paleothermometers (Holbourn et al., 2018). It is also consistent with the evidence for deep ocean cooling (Holbourn et al., 2013; Lear et al., 2000), as well as reports of cooling in upwelling zones in the Arabian Sea (whether due to upwelling intensification or cooler upwelled waters) based on the abundance of cold-loving *Globigerina bulloides* foraminifera (Gupta et al., 2015; Kroon, 1988). Changes in SSTs and in particular SST gradients are found likely to affect monsoon rainfall in idealized (Lutsko et al., 2019) and high resolution climate model experiments (Acosta and Huber, 2020).

We present the TEX₈₆-based BAYSPAR temperature reconstructions in the context of an orbitally-resolved foraminiferal record from the South China Sea where benthic $\delta^{18}\text{O}$ record high latitude temperatures and Antarctic ice volume instability with obliquity pacing (Holbourn et al., 2018) (**Figure 5f**). Both estimates are on the GTS12 timescale and well-dated, thus we can compare the discrete TEX₈₆-based estimates with the orbitally-resolved record, finding good agreement. For example, the warmest TEX₈₆ estimate of 28.5°C at 7.37 Ma corresponds to a warm and low ice volume “interglacial” (one of the lightest benthic $\delta^{18}\text{O}$ values). Independent corroboration of locally warm temperatures in that same sample at 7.37 Ma comes from the palynomorph analysis that includes dinoflagellates, and a species, *Operculodinium israelianum*, that is normally found in waters with summer maxima $>29.2^{\circ}\text{C}$ (Marret and Zonneveld, 2003), together providing compelling evidence that Indian Ocean SSTs warmed in phase with the high latitudes. At the other extreme, we find several consecutive cooler estimates near 7.2 Ma (**Figure 5C**), that correspond to a “prolonged cold event” (Holbourn et al., 2018) where the obliquity-paced cycles appear to “skip a beat” at an eccentricity low.

Basic C₄ energetic theory predicts that the combination of low $p\text{CO}_2$ and warmer temperatures promote C₄ competitive advantage (Ehleringer et al., 1997), however additional factors influence competition and the interplay of factors is complex. Mean annual precipitation and its seasonal distribution is a key factor in tree cover versus grassland (Cotton et al., 2016; Guan et al., 2014). $p\text{CO}_2$ thresholds vary with rainfall amount, with C₄ found in drier climates at lower $p\text{CO}_2$ (Higgins and Scheiter, 2012), and conversely in the higher than pre-Industrial $p\text{CO}_2$ levels of the Miocene, C₄ tropical grasslands may have been found in rainfall regimes that today support forests. In tropical mountains of western peninsular India, cool season temperatures are found to limit tropical trees, but warm summers allow C₄ grassland patches at ~ 2000 m above sea level today and the ecosystems are thought to be bistable (Das et al., 2015). In bistable ecosystems, where either forest or grassland could exist, other factors of recruitment and disturbance may control the forest-grassland transition including fire (Beckage and Ellingwood, 2008; Higgins and Scheiter, 2012). Transitions are expected to be abrupt and with a hysteresis

effect, for example, as grasses are more flammable than trees so the threshold fire frequency to maintain grasses is lower than that needed to clear forest (Beckage and Ellingwood, 2008).

During the last Pleistocene glacial, abundant C₄ grasslands have been reported from the lowlands of the Ganges-Brahmaputra catchment (during Heinrich Stadial 1) interpreted as a response to lowered $p\text{CO}_2$ and drying, followed by C₃ replacement and mixed ecosystems in the Holocene as monsoon rainfall increased (Agrawal et al., 2014; Contreras-Rosales et al., 2014; Hein et al., 2017). Similarly, C₄ grasslands expanded in the Western Ghats where forests were increasingly limited by cooler temperatures and reduced summer rainfall (Caner et al., 2007). In contrast dry conditions in the (already dry) glacial in the Indus catchment did not promote grasses, and instead grasses expanded in the wetter early Holocene (Ansari and Vink, 2007). Thus the triggers for C₄ expansion vary within the region with varied different tipping points for grassland conditions according to the local context.

Whether the late Miocene C₄ expansion was driven by similar-to-glacial-transition drivers – a drop in $p\text{CO}_2$, cooling and drying – remains an open question. Evidence for a late Miocene $p\text{CO}_2$ drop has long been elusive since first proposed (Cerling et al., 1997). It was initially rejected by alkenone based estimates (Pagani et al., 1999), but more recently found some support from diatom associated organics, calcification and boron isotopes, although the data remain very preliminary (Badger et al., 2013; Bolton et al., 2016; Mejía et al., 2017; Sosdian et al., 2018). New solutions may emerge e.g. refinements to the alkenone method (Zhang et al., 2019) and there is a keen interest in detailed, robust $p\text{CO}_2$ reconstructions in order to understand the drivers of this late Miocene ecological transformation. During the prolonged cold event near 7.2 Ma, Holbourn *et al.*, (2018) infer a drop in $p\text{CO}_2$ based on a greater difference between the $\delta^{13}\text{C}$ of planktic and benthic foraminera. However, alternative possibilities could explain the increased offset, including habitat changes in foraminiferal communities, changes in ocean ventilation influencing the benthic values (Keigwin, 1979) and enrichment of the carbon isotopic composition of the terrestrial biosphere altering that of the atmosphere and surface ocean – i.e. the C₄ expansion could lead to a ^{13}C -depletion in the residual pool in the atmosphere and planktic carbonates (Derry and France-Lanord, 1996). While, the C₄ expansion is time-transient around the tropics, the grassland expansion described here, at first gradual and then abruptly switching at 7.2 Ma described here, seems to coincide with gradual and then abrupt change at 7.2 Ma with the final phase of the oceanic carbon isotope shift complete by 7.0 Ma (Holbourn et al., 2018). This correspondence allows for the possibility that the terrestrial vegetation drives the abrupt change in the ocean isotopic balance.

At this time, detailed and multi-proxy confirmatory evidence of the carbon dioxide record for the late Miocene – and any possible ties to C₄ expansion – remains elusive, including in recent climate model experiments for North America with a responsive biosphere (Lee, 2019). As independent evolutionary C₄ innovations occurred across the tropics, we should anticipate that the ecological response of the local species may differ and would be triggered by the specific interplay of climatic factors that trigger local ecologic tipping points. The likely answer then to what caused the C₄ expansion appears to be multi-factorial regional drivers and regional ecological response (Osborne, 2008). The timing of the response to even a global forcing such as declining carbon dioxide levels would be expected to yield time-transitions based on regional tipping points (Higgins and Scheiter, 2012).

4 Conclusions

In this study of late Miocene sediments of Indus Fan at IODP Site U1457, we find distinct changes in sedimentation rate and composition between turbidites and hemipelagic sediments. We present a multi-proxy biomarker and microfossil study of the terrigenous components in these sediments and determine that provenance switches between facies, implying two discontinuous archives within one core. Our primary marker for the facies-based interpretation is the lipid index, the BIT ratio indicating the dominance of soil microbial inputs (BIT > 0.4 in turbidites) or marine microbial inputs (BIT < 0.2 in hemipelagics). Supporting evidence comes from lignin phenols, which establish transport method (fluvial vs. aeolian), with only fluvial transport delivering lignin to turbidites but not to the hemipelagic facies. Plant wax δD provides confirmatory evidence of source switching with more D-depletion in the plant wax from the Indus derived turbidites, compared to D-enrichment in wind-blown waxes from the Indian peninsula.

We observe a vegetation change in each source region within our facies-specific reconstructions. In the turbidites we have massive accumulations but only brief windows of time represented. The available turbidites can only bracket the C₄ expansion in the Indus River catchment to between 7.4 and 5.9 Ma. We find the hemipelagic sediments allow for a detailed investigation of the nature of the transition on peninsular India. Although there is likely some C₄ present by 8.7 Ma, the main transition is revealed in bulk OC, plant wax *n*-alkanes and *n*-alkanoic acids to occur across 7.4 to 7.2 Ma (dated to within 0.1 Ma), with supporting data from grass pollen and grass-derived charcoal.

We see a marked increase in charcoal abundance suggesting more fire on the landscape cleared forests, and that grass fire (elongate charcoal) increased proportionally as grasslands expanded. The flammability of grasses, known to be a key feedback mechanism in maintaining grasslands in bistable ecosystem conditions, allowed for the persistence of C₄ grasslands once they appeared in both regions and explains the step shift in vegetation observed on peninsular India.

Paired analysis of plant wax δD and $\delta^{13}C$ does not diagnose whether the monsoon precipitation changes drove the C₄ expansion in the region, as the δD value of plant waxes and thus the reconstructed isotopes of precipitation are found to be dominated by source changes in this archive. Within facies we are unable to identify a clear change in precipitation isotopic composition associated with the C₄ expansion. The sparse data suggest that the hydroclimate of the Indian peninsula was broadly similar to today, at least as detected by precipitation isotopes, although the Indus River catchment may have been more strongly influenced by westerly winter precipitation (more D-depleted) than today. A change in monsoon rainfall may not be the culprit for C₄ expansion or possible grasslands could respond to a precipitation change that was not detected by these plant wax precipitation isotopic proxies, such as a reduction of dry season precipitation that would also promote the observed increase in fire. While we have no evidence for drying here other than the increase of grasslands and fire, bistable ecosystems that display hysteresis behavior may not require much of a climatic shift to trigger a transition of the kind observed in the late Miocene.

Lowering of atmospheric *p*CO₂ may have both cooled the global climate and also directly increased the competitiveness of C₄ grasses – *p*CO₂ reconstructions allow this but remain tentative for the late Miocene. A late Miocene cooling has been identified elsewhere, and our TEX₈₆ data shows this cooling was reproduced locally in the Arabian Sea. Arabian Sea cooling was likely in part driven by cooling of the high latitudes and thus deep waters affecting the temperature of upwelled waters and in part by radiative changes associated with a drop in

$p\text{CO}_2$. Although our low resolution record aliases some of the orbital variability seen elsewhere, we have finely resolved sampling that shows a cooling of $> 1^\circ\text{C}$ at 7.3 Ma (dated to within 0.1 Ma) and overall variability and cooling on the order of 4°C . It is therefore possible that the cooling of the Arabian Sea was implicated in the ecological transition, perhaps by drying the dry season, which would be the most likely explanation for the increase in fire observed here. However, the ocean-atmosphere connection that hypothetically linked cooler with reduced rainfall, remains elusive because of the lack of a rainfall shift detected by hydrogen isotopes in plant waxes here. It is possible that only modest climatic changes are needed to trigger and maintain ecological switches in bistable ecosystem settings once the C_4 innovation appeared.

Acknowledgments, Samples, and Data

We declare no financial conflicts of interests for any author, or their affiliations. This research was funded by the US National Science Foundation (OCE 14-50528 to Consortium for Ocean Leadership, sub-awards GG0093093-01 to S. Feakins, sub-award GG0093093-01 to S. Warny, sub-award GG0093093-01 to L. Tauxe and EAR 1547263 to L. Tauxe); microcharcoal analyses were supported by the National Natural Science Foundation of China grant 41772181. This research used samples and/or data collected by the International Ocean Discovery Program, supported by funding from the US National Science Foundation and other member nations. We thank all participants of the shipboard science party and crew on Expedition 355. We thank the following for laboratory assistance: A. Figueroa, J. Sunwoo, and C. Johnson. Data files are archived at the NOAA paleoclimatology database at <https://www.ncdc.noaa.gov/paleo/study/28792> (Feakins, 2020).

Author Contribution Statement: S.F. designed and led the study. H.L and L.T. were shipboard scientists. The following contributed analyses and interpretations: H.L and S.F. (plant wax); L.T. (age model); V.G. (bulk organics); J.T. (membrane lipids); X.F., (lignin); Y.M. (charcoal) and S.W. (pollen). S.F. analyzed and interpreted the multi-proxy data, graphed the results and wrote the manuscript. All authors contributed to the study.

References

- Acosta, R.P. and Huber, M. (2020) Competing topographic mechanisms for the Summer Indo-Asian Monsoon. *Geophysical Research Letters*. <http://dx.doi.org/10.1029/2019gl085112>.
- Agrawal, S., Galy, V., Sanyal, P. and Eglinton, T. (2014) C_4 plant expansion in the Ganga Plain during the last glacial cycle: Insights from isotopic composition of vascular plant biomarkers. *Organic Geochemistry* 67, 58-71. <http://dx.doi.org/10.1016/j.orggeochem.2013.12.007>.
- Ansari, M.H. and Vink, A. (2007) Vegetation history and palaeoclimate of the past 30 kyr in Pakistan as inferred from the palynology of continental margin sediments off the Indus Delta. *Review of Palaeobotany and Palynology* 145, 201-216. <https://doi.org/10.1016/j.revpalbo.2006.10.005>.
- Badger, M.P.S., Lear, C.H., Pancost, R.D., Foster, G.L., Bailey, T.R., Leng, M.J. and Abels, H.A. (2013) CO_2 drawdown following the middle Miocene expansion of the Antarctic Ice Sheet. *Paleoceanography* 28, 42-53. <https://doi.org/10.1002/palo.20015>.
- Beckage, B. and Ellingwood, C. (2008) Fire Feedbacks with Vegetation and Alternative Stable States. *Complex Systems* 18, 159-173. <https://doi.org/10.25088/ComplexSystems.18.1.159>.
- Bolton, C.T., Hernández-Sánchez, M.T., Fuertes, M.-Á., González-Lemos, S., Abrevaya, L., Mendez-Vicente, A., Flores, J.-A., Probert, I., Giosan, L., Johnson, J. and Stoll, H.M. (2016) Decrease in coccolithophore calcification and CO_2 since the middle Miocene. *Nature Communications* 7, 10284. <http://dx.doi.org/10.1038/ncomms10284>.
- Bourget, J., Zaragosi, S., Rodriguez, M., Fournier, M., Garlan, T. and Chamot-Rooke, N. (2013) Late Quaternary megaturbidites of the Indus Fan: Origin and stratigraphic significance. *Marine Geology* 336, 10-23. <http://dx.doi.org/10.1016/j.margeo.2012.11.011>.
- Brown, C. (2008) *Palynological Techniques*. AASP Special Publications.
- Calvès, G., Huuse, M., Clift, P.D. and Brusset, S. (2015) Giant fossil mass wasting off the coast of West India: The Nataraja submarine slide. *Earth and Planetary Science Letters* 432, 265-272. <https://doi.org/10.1016/j.epsl.2015.10.022>.

- Caner, L., Seen, D.L., Gunnell, Y., Ramesh, B.R. and Bourgeon, G. (2007) Spatial heterogeneity of land cover response to climatic change in the Nilgiri highlands (Southern India) since the last glacial maximum. *The Holocene* 17, 195-205. <http://dx.doi.org/10.1177/0959683607075833>.
- Cerling, T.E., Harris, J.M., MacFadden, B.J., Leakey, M.G., Quade, J., Eisenmann, V. and Ehleringer, J.R. (1997) Global vegetation change through the Miocene/Pliocene boundary. *Nature* 389, 153-158.
- Clift, P., Giosan, L. and Carter, A. (2010) Monsoon control over erosion patterns in the Western Himalaya: possible feed-backs into the tectonic evolution. *Monsoon Evolution and Tectonic-Climate Linkage in Asia*. Special Publication, 342, 181-213.
- Clift, P.D., Zhou, P., Stockli, D.F. and Blusztajn, J. (2019) Regional Pliocene exhumation of the Lesser Himalaya in the Indus drainage. *Solid Earth* 10, 647-661. <http://dx.doi.org/10.5194/se-10-647-2019>.
- Contreras-Rosales, L.A., Jennerjahn, T., Tharammal, T., Meyer, V., Lückge, A., Paul, A. and Schefuß, E. (2014) Evolution of the Indian Summer Monsoon and terrestrial vegetation in the Bengal region during the past 18 ka. *Quaternary Science Reviews* 102, 133-148. <https://doi.org/10.1016/j.quascirev.2014.08.010>.
- Cotton, J.M., Cerling, T.E., Hoppe, K.A., Mosier, T.M. and Still, C.J. (2016) Climate, CO₂, and the history of North American grasses since the Last Glacial Maximum. *Science Advances* <http://dx.doi.org/2.10.1126/sciadv.1501346>.
- Dahl, K.A., Oppo, D.W., Eglinton, T.I., Hughen, K.A., Curry, W.B. and Sirocko, F. (2005) Terrestrial plant wax inputs to the Arabian Sea: Implications for the reconstruction of winds associated with the Indian Monsoon. *Geochimica Et Cosmochimica Acta* 69, 2547-2558. <http://dx.doi.org/10.1016/j.gca.2005.01.001>.
- Dantas, V.d.L., Hirota, M., Oliveira, R.S. and Pausas, J.G. (2016) Disturbance maintains alternative biome states. *Ecology Letters* 19, 12-19. <http://dx.doi.org/10.1111/ele.12537>.
- Das, A., Nagendra, H., Anand, M. and Bunyan, M. (2015) Topographic and Bioclimatic Determinants of the Occurrence of Forest and Grassland in Tropical Montane Forest-Grassland Mosaics of the Western Ghats, India. *Plos One* 10, e0130566. <https://doi.org/10.1371/journal.pone.0130566>.
- Derry, L.A. and France-Lanord, C. (1996) Neogene growth of the sedimentary organic carbon reservoir. *Paleoceanography* 11, 267-275.
- Ehleringer, J.R., Cerling, T.E. and Helliker, B.R. (1997) C-4 photosynthesis, atmospheric CO₂ and climate. *Oecologia* 112, 285-299.
- Feakins, S., deMenocal, P. and Eglinton, T. (2005) Biomarker records of Late Neogene Changes in East African Vegetation. *Geology* 33, 977-980.
- Feakins, S., Levin, N., Liddy, H., Sieracki, A., Eglinton, T. and Bonnefille, R. (2013) Northeast African vegetation change over 12 million years. *Geology* 41, 295-298. <http://dx.doi.org/10.1130/G33845.1>.
- Feakins, S. and Sessions, A.L. (2010) Controls on the D/H ratios of plant leaf waxes from an arid ecosystem. *Geochimica Et Cosmochimica Acta* 74, 2128-2141. <http://dx.doi.org/10.1016/j.gca.2010.01.016>.
- Feakins, S.J. (2013) Pollen-corrected leaf wax D/H reconstructions of northeast African hydrological changes during the late Miocene. *Palaeogeography, Palaeoclimatology, Palaeoecology* 374, 62-71. <http://dx.doi.org/10.1016/j.palaeo.2013.01.004>.
- Feakins, S.J. (2020) Indus Fan, plant biomarkers and microfossils, 10-5 million years, In supplement to: Feakins S.J., Liddy, H.M., Tauxe, L., Galy, V., Feng, X; Tierney, J.E.; Miao, Y.; Warny, S., Late Miocene C4 expansion and hydrological change in the Indus River Catchment, *Paleoceanography and Paleoclimatology*. <https://www.ncdc.noaa.gov/paleo/study/28792>. NOAA
- Feakins, S.J., Bentley, L.P., Salinas, N., Shenkin, A., Blonder, B., Goldsmith, G.R., Ponton, C., Arvin, L.J., Wu, M.S., Peters, T., West, A.J., Martin, R.E., Enquist, B.J., Asner, G.P. and Malhi, Y. (2016) Plant leaf wax biomarkers capture gradients in hydrogen isotopes of precipitation from the Andes and Amazon. *Geochimica Et Cosmochimica Acta* 182, 155-172. <http://dx.doi.org/10.1016/j.gca.2016.03.018>.
- Feakins, S.J., Kirby, M.E., Cheetham, M.I., Ibarra, Y. and Zimmerman, S.R.H. (2014) Fluctuation in leaf wax D/H ratio from a southern California lake records significant variability in isotopes in precipitation during the late Holocene. *Organic Geochemistry* 66, 48-59. <http://dx.doi.org/10.1016/j.orggeochem.2013.10.015>.
- Feakins, S.J., Wu, M.S., Ponton, C., Galy, V. and West, A.J. (2018) Dual isotope evidence for sedimentary integration of plant wax biomarkers across an Andes-Amazon elevation transect. *Geochimica Et Cosmochimica Acta* 242, 64-81. <https://doi.org/10.1016/j.gca.2018.09.007>.
- France-Lanord, C. and Derry, L.A. (1997) Organic carbon burial forcing of the carbon cycle from Himalayan erosion. *Nature* 390, 65-67.
- Freeman, K.H. and Colarusso, L.A. (2001) Molecular and isotopic records of C4 grassland expansion in the late Miocene. *Geochimica Et Cosmochimica Acta* 65, 1439-1454.

- Galy, V., Eglinton, T., France-Lanord, C. and Sylva, S. (2011) The provenance of vegetation and environmental signatures encoded in vascular plant biomarkers carried by the Ganges-Brahmaputra rivers. *Earth and Planetary Science Letters* 304, 1-12. <https://doi.org/10.1016/j.epsl.2011.02.003>.
- Garcin, Y., Schefuß, E., Schwab, V.F., Garreta, V., Gleixner, G., Vincens, A., Todou, G., Séné, O., Onana, J.-M., Achoundong, G. and Sachse, D. (2014) Reconstructing C3 and C4 vegetation cover using n-alkane carbon isotope ratios in recent lake sediments from Cameroon, Western Central Africa. *Geochimica Et Cosmochimica Acta* 142, 482-500. <http://dx.doi.org/10.1016/j.gca.2014.07.004>.
- Ghosh, S., Sanyal, P. and Kumar, R. (2017) Evolution of C4 plants and controlling factors: Insight from n-alkane isotopic values of NW Indian Siwalik paleosols. *Organic Geochemistry* 110, 110-121. <https://doi.org/10.1016/j.orggeochem.2017.04.009>.
- Gradstein, F.M., Ogg, J.G., Schmitz, M.D. and Ogg, G.M. (2012) *The Geologic Time Scale 2012*. Elsevier, Boston.
- Guan, K., Wood, E.F., Medvigy, D., Kimball, J., Pan, M., Caylor, K.K., Sheffield, J., Xu, X. and Jones, M.O. (2014) Terrestrial hydrological controls on land surface phenology of African savannas and woodlands. *Journal of Geophysical Research: Biogeosciences* 119, 1652-1669. <https://doi.org/10.1002/2013jg002572>.
- Gupta, A.K., Yuvaraja, A., Prakasam, M., Clemens, S.C. and Velu, A. (2015) Evolution of the South Asian monsoon wind system since the late Middle Miocene. *Palaeogeography, Palaeoclimatology, Palaeoecology* 438, 160-167. <https://doi.org/10.1016/j.palaeo.2015.08.006>.
- Hein, C.J., Galy, V., Galy, A., France-Lanord, C., Kudrass, H. and Schwenk, T. (2017) Post-glacial climate forcing of surface processes in the Ganges–Brahmaputra river basin and implications for carbon sequestration. *Earth and Planetary Science Letters* 478, 89-101. <https://doi.org/10.1016/j.epsl.2017.08.013>.
- Hemingway, J.D., Schefuß, E., Dinga, B.J., Pryer, H. and Galy, V.V. (2016) Multiple plant-wax compounds record differential sources and ecosystem structure in large river catchments. *Geochimica Et Cosmochimica Acta* 184, 20-40. <http://dx.doi.org/10.1016/j.gca.2016.04.003>.
- Herbert, T.D., Lawrence, K.T., Tzanova, A., Peterson, L.C., Caballero-Gill, R. and Kelly, C.S. (2016) Late Miocene global cooling and the rise of modern ecosystems. *Nature Geoscience* 9, 843. <https://doi.org/10.1038/ngeo2813>
- Higgins, S.I. and Scheiter, S. (2012) Atmospheric CO2 forces abrupt vegetation shifts locally, but not globally. *Nature* 488, 209-212. <http://dx.doi.org/10.1038/nature11238>.
- Hoetzel, S., Dupont, L., Schefus, E., Rommerskirchen, F. and Wefer, G. (2013) The role of fire in Miocene to Pliocene C4 grassland and ecosystem evolution. *Nature Geosci* 6, 1027-1030. <http://dx.doi.org/10.1038/ngeo1984>.
- Holbourn, A., Kuhnt, W., Clemens, S., Prell, W. and Andersen, N. (2013) Middle to late Miocene stepwise climate cooling: Evidence from a high-resolution deep-water isotope curve spanning 8 million years. *Paleoceanography*, 2013PA002538. <http://dx.doi.org/10.1002/2013pa002538>.
- Holbourn, A.E., Kuhnt, W., Clemens, S.C., Kochhann, K.G.D., Jöhnck, J., Lübbers, J. and Andersen, N. (2018) Late Miocene climate cooling and intensification of southeast Asian winter monsoon. *Nature Communications* 9, 1584. <http://dx.doi.org/10.1038/s41467-018-03950-1>.
- Hopmans, E.C., Weijers, J.W.H., Schefuß, E., Herfort, L., Sinninghe Damsté, J.S. and Schouten, S. (2004) A novel proxy for terrestrial organic matter in sediments based on branched and isoprenoid tetraether lipids. *Earth and Planetary Science Letters* 224, 107-116. <https://doi.org/10.1016/j.epsl.2004.05.012>.
- Huang, Y., Clemens, S.C., Liu, W., Wang, Y. and Prell, W.L. (2007) Large-scale hydrological change drove the late Miocene C4 plant expansion in the Himalayan foreland and Arabian Peninsula. *Geology* 35, 531-534. <http://dx.doi.org/10.1130/G23666A>.
- IAEA/WMO (2019) Global Network of Isotopes in Precipitation. The GNIP Database. Accessible at: <http://www.iaea.org/water>.
- Ivory, S.J. and Lezine, A.M. (2009) Climate and environmental change at the end of the Holocene Humid Period: A pollen record off Pakistan. *Comptes Rendus Geoscience* 341, 760-769. [10.1016/j.crte.2008.12.009](https://doi.org/10.1016/j.crte.2008.12.009).
- Karp, A.T., Behrensmeyer, A.K. and Freeman, K.H. (2018) Grassland fire ecology has roots in the late Miocene. *Proceedings of the National Academy of Sciences* 115, 12130-12135. <http://dx.doi.org/10.1073/pnas.1809758115>.
- Keigwin, L.D. (1979) Late Cenozoic stable isotope stratigraphy and paleoceanography of DSDP sites from the east equatorial and central north Pacific Ocean. *Earth and Planetary Science Letters* 45, 361-382. [https://doi.org/10.1016/0012-821X\(79\)90137-7](https://doi.org/10.1016/0012-821X(79)90137-7).
- Kessarkar, P.M., Rao, V.P., Ahmad, S.M. and Babu, G.A. (2003) Clay minerals and Sr–Nd isotopes of the sediments along the western margin of India and their implication for sediment provenance. *Marine Geology* 202, 55-69. [https://doi.org/10.1016/S0025-3227\(03\)00240-8](https://doi.org/10.1016/S0025-3227(03)00240-8).

- Kroon, D. (1988) Distribution of extant planktic foraminiferal assemblages in Red Sea and Northern Indian Ocean surface waters, in: Kroon, D. (Ed.), *Planktonic foraminifera as tracers of ocean-climate history*. Free University Press, Amsterdam, p. 346.
- Kroon, D., Steens, T. and Troelstra, S.R. (1991) Onset of monsoonal related upwelling in the western Arabian Sea as revealed by planktonic foraminifera. *Proc., scientific results, ODP, Leg 117, Oman Margin/Neogene package*, 257-263.
- Kusch, S., Rethemeyer, J., Schefuß, E. and Mollenhauer, G. (2010) Controls on the age of vascular plant biomarkers in Black Sea sediments. *Geochimica Et Cosmochimica Acta* 74, 7031-7047. [10.1016/j.gca.2010.09.005](https://doi.org/10.1016/j.gca.2010.09.005).
- Lear, C., Elderfield, H. and Wilson, P. (2000) Cenozoic Deep-Sea Temperatures and Global Ice Volumes from Mg/Ca in Benthic Foraminiferal Calcite. *Science* 287, 269-272.
- Lear, C.H., Coxall, H.K., Foster, G.L., Lunt, D.J., Mawbey, E.M., Rosenthal, Y., Sosdian, S.M., Thomas, E. and Wilson, P.A. (2015) Neogene ice volume and ocean temperatures: Insights from infaunal foraminiferal Mg/Ca paleothermometry. *Paleoceanography* 30, 1437-1454. [http://dx.doi.org/10.1002/2015pa002833](https://doi.org/10.1002/2015pa002833).
- Lee, J.-E. (2019) Understanding Neogene Oxygen Isotopes in the Southern Great Plains Using Isotope-Enabled General Circulation Model Simulations. *Journal of Geophysical Research: Atmospheres* 124, 2452-2464. [http://dx.doi.org/10.1029/2018jd028894](https://doi.org/10.1029/2018jd028894).
- Levin, N.E. (2015) Environment and Climate of Early Human Evolution. *Annual Review of Earth and Planetary Sciences* 43, 405-429. [http://dx.doi.org/10.1146/annurev-earth-060614-105310](https://doi.org/10.1146/annurev-earth-060614-105310).
- Liddy, H.M., Feakins, S.J. and Tierney, J.E. (2016) Cooling and drying in northeast Africa across the Pliocene. *Earth and Planetary Science Letters* 449, 430-438. [http://dx.doi.org/10.1016/j.epsl.2016.05.005](https://doi.org/10.1016/j.epsl.2016.05.005).
- Locarnini, A.R., Mishonov, A., Antonov, I.J., Boyer, P.T., Garcia, H., Baranova, K.O., Zweng, M. and Johnson, D. (2010) *World Ocean Atlas 2009, Vol. 1: Temperature*, p. 184.
- Lutsko, N.J., Marshall, J. and Green, B. (2019) Modulation of Monsoon Circulations by Cross-Equatorial Ocean Heat Transport. *Journal of Climate* 32, 3471-3485. [http://dx.doi.org/10.1175/jcli-d-18-0623.1](https://doi.org/10.1175/jcli-d-18-0623.1).
- Makou, M., Eglinton, T., McIntyre, C., Montluçon, D., Antheaume, I. and Grossi, V. (2018) Plant Wax n-Alkane and n-Alkanoic Acid Signatures Overprinted by Microbial Contributions and Old Carbon in Meromictic Lake Sediments. *Geophysical Research Letters* 45, 1049-1057. [http://dx.doi.org/10.1002/2017GL076211](https://doi.org/10.1002/2017GL076211).
- Marret, F. and Zonneveld, K.A.F. (2003) Atlas of modern organic-walled dinoflagellate cyst distribution. *Review of Palaeobotany and Palynology* 125, 1-200. [https://doi.org/10.1016/S0034-6667\(02\)00229-4](https://doi.org/10.1016/S0034-6667(02)00229-4).
- Mejía, L.M., Méndez-Vicente, A., Abrevaya, L., Lawrence, K.T., Ladlow, C., Bolton, C., Cacho, I. and Stoll, H. (2017) A diatom record of CO₂ decline since the late Miocene. *Earth and Planetary Science Letters* 479, 18-33. <https://doi.org/10.1016/j.epsl.2017.08.034>.
- Métivier, F., Gaudemer, Y., Tapponnier, P. and Klein, M. (1999) Mass accumulation rates in Asia during the Cenozoic. *Geophysical Journal International* 137, 280-318. [http://dx.doi.org/10.1046/j.1365-246X.1999.00802.x](https://doi.org/10.1046/j.1365-246X.1999.00802.x).
- Miao, Y., Wu, F., Warny, S., Fang, X., Lu, H., Fu, B., Song, C., Yan, X., Escarguel, G., Yang, Y., Meng, Q. and Shi, P. (2019) Miocene fire intensification linked to continuous aridification on the Tibetan Plateau. *Geology* 47, 303-307. [http://dx.doi.org/10.1130/g45720.1](https://doi.org/10.1130/g45720.1).
- Milliman, J.D. and Syvitski, J.P.M. (1992) Geomorphic/Tectonic Control of Sediment Discharge to the Ocean: The Importance of Small Mountainous Rivers. *The Journal of Geology* 100, 525-544. [10.1086/629606](https://doi.org/10.1086/629606).
- Murakami, H., Vecchi, G.A. and Underwood, S. (2017) Increasing frequency of extremely severe cyclonic storms over the Arabian Sea. *Nature Climate Change* 7, 885-889. [http://dx.doi.org/10.1038/s41558-017-0008-6](https://doi.org/10.1038/s41558-017-0008-6).
- Osborne, C.P. (2008) Atmosphere, ecology and evolution: what drove the Miocene expansion of C-4 grasslands? *Journal of Ecology* 96, 35-45. [http://dx.doi.org/10.1111/j.1365-2745.2007.01323.x](https://doi.org/10.1111/j.1365-2745.2007.01323.x).
- Pagani, M., Freeman, K.H. and Arthur, M.A. (1999) Late Miocene atmospheric CO₂ concentrations and the expansion of C-4 grasses. *Science* 285, 876-879.
- Pandey, D., Clift, P., Kulhanek, D., Andò, S., Bendle, J., Bratenkov, S., Griffith, E., Gurumurthy, G., Hahn, A. and Iwai, M. (2015) Expedition 355 Preliminary Report: Arabian Sea Monsoon. *Proceedings of the International Ocean Discovery Program* 2015, 1-46.
- Polissar, P.J., Rose, C., Uno, K.T., Phelps, S.R. and deMenocal, P. (2019) Synchronous rise of African C4 ecosystems 10 million years ago in the absence of aridification. *Nature Geoscience*, 12, 657-660. <https://doi.org/10.1038/s41561-019-0399-22>.
- Prins, M.A., Postma, G. and Weltje, G.J. (2000) Controls on terrigenous sediment supply to the Arabian Sea during the late Quaternary: the Makran continental slope. *Marine Geology* 169, 351-371. [https://doi.org/10.1016/S0025-3227\(00\)00087-6](https://doi.org/10.1016/S0025-3227(00)00087-6).

- Quade, J. and Cerling, T.E. (1995) Expansion of C-4 Grasses in the Late Miocene of Northern Pakistan - Evidence from Stable Isotopes in Paleosols. *Palaeogeography Palaeoclimatology Palaeoecology* 115, 91-116.
- Quade, J., Cerling, T.E. and Bowman, J.R. (1989) Development of Asian Monsoon Revealed by Marked Ecological Shift During the Latest Miocene in Northern Pakistan. *Nature* 342, 163-166.
- Rodriguez, M., Chamot-Rooke, N., Huchon, P., Fournier, M. and Delescluse, M. (2014) The Owen Ridge uplift in the Arabian Sea: Implications for the sedimentary record of Indian monsoon in Late Miocene. *Earth and Planetary Science Letters* 394, 1-12.
- Routledge, C.M., Kulhanek, D.K., Tauxe, L., Scardia, G., Singh, A.D., Steinke, S., Griffith, E.M. and Saraswat, R. (2019) A revised chronostratigraphic framework for International Ocean Discovery Program Expedition 355 sites in Laxmi Basin, eastern Arabian Sea. *Geological Magazine*, 1-18. <https://doi.org/10.1017/S0016756819000104>.
- Sachse, D., Billault, I., Bowen, G.J., Chikaraishi, Y., Dawson, T.E., Feakins, S.J., Freeman, K.H., Magill, C.R., McInerney, F.A., van der Meer, M.T.J., Polissar, P., Robins, R.J., Sachs, J.P., Schmidt, H.-L., Sessions, A.L., White, J.W.C., West, J.B. and Kahmen, A. (2012) Molecular Paleohydrology: Interpreting the Hydrogen-Isotopic Composition of Lipid Biomarkers from Photosynthesizing Organisms. *Annual Review of Earth and Planetary Sciences* 40, 221-249. <https://doi.org/10.1146/annurev-earth-042711-105535>.
- Schouten, S., Huguot, C., Hopmans, E.C., Kienhuis, M.V.M. and Damste, J.S.S. (2007a) Analytical methodology for TEX86 paleothermometry by high-performance liquid chromatography/atmospheric pressure chemical ionization-mass spectrometry. *Analytical Chemistry* 79, 2940-2944.
- Schouten, S., van der Meer, M.T.J., Hopmans, E.C., Rijpstra, W.I.C., Reysenbach, A.L., Ward, D.M. and Damste, J.S.S. (2007b) Archaeal and bacterial glycerol dialkyl glycerol tetraether lipids in hot springs of Yellowstone National Park. *Applied and Environmental Microbiology* 73, 6181-6191.
- Smith, F. and Freeman, K. (2006) Influence of physiology and climate on dD of leaf wax n-alkanes from C3 and C4 grasses. *Geochimica Et Cosmochimica Acta* 70, 1172-1187.
- Sosdian, S.M., Greenop, R., Hain, M.P., Foster, G.L., Pearson, P.N. and Lear, C.H. (2018) Constraining the evolution of Neogene ocean carbonate chemistry using the boron isotope pH proxy. *Earth and Planetary Science Letters* 498, 362-376. <https://doi.org/10.1016/j.epsl.2018.06.017>.
- Staver, A.C., Archibald, S. and Levin, S.A. (2011) The Global Extent and Determinants of Savanna and Forest as Alternative Biome States. *Science* 334, 230-232. <https://doi.org/10.1126/science.1210465>.
- Suh, Y.J., Diefendorf, A.F., Bowen, G.J., Cotton, J.M. and Ju, S.-J. (2019) Plant wax integration and transport from the Mississippi River Basin to the Gulf of Mexico inferred from GIS-enabled isoscapes and mixing models. *Geochimica Et Cosmochimica Acta* 257, 131-149. <https://doi.org/10.1016/j.gca.2019.04.022>.
- Tauxe, L. and Feakins, S.J. (this issue) A re-assessment of the timing of the late Miocene C3-C4 transition. *Paleoceanography and Paleoclimatology manuscript*: 2020PA003857.
- Tierney, J.E., Pausata, F.S.R. and deMenocal, P. (2016) Deglacial Indian monsoon failure and North Atlantic stadials linked by Indian Ocean surface cooling. *Nature Geosci* 9, 46-50. <https://doi.org/10.1038/ngeo2603> <http://www.nature.com/ngeo/journal/v9/n1/abs/ngeo2603.html#supplementary-information>.
- Tierney, J.E. and Tingley, M.P. (2014) A Bayesian, spatially-varying calibration model for the TEX86 proxy. *Geochimica Et Cosmochimica Acta* 127, 83-106. <https://doi.org/10.1016/j.gca.2013.11.026>.
- Uno, K.T., Cerling, T.E., Harris, J.M., Kunitatsu, Y., Leakey, M.G., Nakatsukasa, M. and Nakaya, H. (2011) Late Miocene to Pliocene carbon isotope record of differential diet change among East African herbivores. *Proceedings of the National Academy of Sciences of the United States of America* 108, 6509-6514.
- Uno, K.T., Polissar, P.J., Jackson, K.E. and deMenocal, P.B. (2016) Neogene biomarker record of vegetation change in eastern Africa. *Proceedings of the National Academy of Sciences* 113, 6355-6363. <https://doi.org/10.1073/pnas.1521267113>.
- Vögeli, N., Najman, Y., van der Beek, P., Huyghe, P., Wynn, P.M., Govin, G., van der Veen, I. and Sachse, D. (2017) Lateral variations in vegetation in the Himalaya since the Miocene and implications for climate evolution. *Earth and Planetary Science Letters* 471, 1-9. <https://doi.org/10.1016/j.epsl.2017.04.037>.
- Whiteside, J.H., Olsen, P.E., Eglinton, T.I., Cornet, B., McDonald, N.G. and Huber, P. (2011) Pangean great lake paleoecology on the cusp of the end-Triassic extinction. *Palaeogeography, Palaeoclimatology, Palaeoecology* 301, 1-17. <http://dx.doi.org/10.1016/j.palaeo.2010.11.025>.
- Yu, Z., Colin, C., Wan, S., Saraswat, R., Song, L., Xu, Z., Clift, P., Lu, H., Lyle, M., Kulhanek, D., Hahn, A., Tiwari, M., Mishra, R., Miska, S. and Kumar, A. (2019) Sea level-controlled sediment transport to the eastern Arabian Sea over the past 600 kyr: Clay minerals and SrNd isotopic evidence from IODP site U1457. *Quaternary Science Reviews* 205, 22-34. <https://doi.org/10.1016/j.quascirev.2018.12.006>.

Zhang, Y.G., Pearson, A., Benthien, A., Dong, L., Huybers, P., Liu, X. and Pagani, M. (2019) Refining the alkenone-pCO₂ method I: Lessons from the Quaternary glacial cycles. *Geochimica Et Cosmochimica Acta* 260, 177-191. <https://doi.org/10.1016/j.gca.2019.06.032>.

Figure Captions:

Figure 1: Location of IODP Site U1457.

Figure 2: Age model for IODP Site U1457C using age control first and last appearance datums for foraminifera (PF, black triangles) and nannofossils (CN, blue triangles) (Routledge et al., 2019) with refined paleomagnetic chronostratigraphy (Pmag, red triangles) (Tauxe and Feakins, this issue). Age model and uncertainty (yellow shading) for hemipelagic sediments, with turbidites under-constrained resulting in a linear age model between 2 points (dashed lines), and erosional unconformities (dotted lines).

Figure 3: Source characterization based on multi-proxy analyses of the same set of samples from Indus Fan IODP Site U1457C. Showing depth distribution of (a) ϵ_{Nd} (Clift et al., 2019) (red circles); (b) the BIT index (blue circle); (c) summed lignin phenols relative to OC concentration (orange circles); and (d) plant wax δD showing example homologues of two compound classes: the C₃₁ *n*-alkanes (white triangles) and C₃₀ *n*-alkanoic acid (black squares). Bivariate plots showing BIT versus (e) ϵ_{Nd} (Clift et al., 2019) (red circles), (f) lignin phenols/OC (orange circles), (g) δD of C₂₈ *n*-alkanoic acid (red squares), C₃₀ *n*-alkanoic acid (black squares), C₂₉ *n*-alkanes (grey triangles) and C₃₁ *n*-alkanes (white triangles); within compound variation between homologues is less than between compound class, alkanes are consistently D-depleted relative to alkanoic acids.

Figure 4: Vegetation indicators from the Indus Fan IODP Site U1457C showing (a) fluvially-exported material from the Indus River catchment delivered via turbidite currents and recovered from low recovery, but expansive turbidites representing short intervals of time (with bounding age control only); and (b) hemipelagic sediments with an inferred peninsula India source of wind-blown plant proxies (with age control on the C₄ transition being complete by 7.2 Ma to within 0.1 Ma). Within each panel the left hand axis shows $\delta^{13}\text{C}$ plant wax isotope records of *n*-alkane C₃₁ (brown triangles) and *n*-alkanoic acid C₃₀ (gold squares) as well as bulk organic carbon isotopic values (white circles). The right hand axis shows microfossil records including non-arboreal (i.e. shrubs and grasses) pollen (gold triangles) and grass pollen (green squares) each as % of all pollen, and long/round charcoal (pink circles) as % of all charcoal (indicates grass-derived charcoal). As the archive represents two delivery mechanisms, separated into panels a and b, the missing time in each archive is indicated (grey shading, i.e. no data).

Figure 5: Multi-proxy investigation of the possible drivers of C₄ expansion in each region, showing fluvial-turbidite Indus sourcing (brown circles) and hemipelagic sediments with aeolian inputs of terrigenous material from peninsular India (blue triangles) to Indus Fan IODP Site U1457. (a) $\delta^{13}\text{C}$ values of C₃₁ *n*-alkanes, (b) microcharcoal long/round (L/R) shape, (c) δD values of paleoprecipitation as reconstructed from C₃₁ *n*-alkanes, (d) Lignin VSC phenols, (e) BIT index, (f) TEX₈₆ based sub-surface temperatures calibrated using BAYSPAR and comparison to $\delta^{18}\text{O}_{\text{benthic}}$ (gray line) (Holbourn et al., 2018).









