

1-1-2020

Excluded minors are almost fragile

Nick Brettell

Technische Universiteit Eindhoven

Ben Clark

Louisiana State University

James Oxley

Louisiana State University

Charles Semple

University of Canterbury

Geoff Whittle

Victoria University of Wellington

Follow this and additional works at: https://digitalcommons.lsu.edu/mathematics_pubs

Recommended Citation

Brettell, N., Clark, B., Oxley, J., Semple, C., & Whittle, G. (2020). Excluded minors are almost fragile. *Journal of Combinatorial Theory. Series B*, 140, 263-322. <https://doi.org/10.1016/j.jctb.2019.05.008>

This Article is brought to you for free and open access by the Department of Mathematics at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Durham Research Online

Deposited in DRO:

15 November 2019

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Brettell, Nick and Clark, Ben and Oxley, James and Semple, Charles and Whittle, Geoff (2020) 'Excluded minors are almost fragile.', *Journal of combinatorial theory, series B.*, 140 . pp. 263-322.

Further information on publisher's website:

<https://doi.org/10.1016/j.jctb.2019.05.008>

Publisher's copyright statement:

© 2019 This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

EXCLUDED MINORS ARE ALMOST FRAGILE

NICK BRETTELL, BEN CLARK, JAMES OXLEY, CHARLES SEMPLE,
AND GEOFF WHITTLE

ABSTRACT. Let M be an excluded minor for the class of \mathbb{P} -representable matroids for some partial field \mathbb{P} , and let N be a 3-connected strong \mathbb{P} -stabilizer that is non-binary. We prove that either M is bounded relative to N , or, up to replacing M by a Δ - Y -equivalent excluded minor, we can choose a pair of elements $\{a, b\}$ such that either $M \setminus \{a, b\}$ is N -fragile, or $M^* \setminus \{a, b\}$ is N^* -fragile.

1. INTRODUCTION

One of the longstanding goals of matroid theory is to find excluded-minor characterisations of classes of representable matroids. Results to date include Tutte's excluded-minor characterisation of binary and regular matroids [21]; Bixby's and, independently, Seymour's excluded-minor characterisation of ternary matroids [1, 20]; Geelen, Gerards and Kapoor's excluded-minor characterisation of $\text{GF}(4)$ -representable matroids [8]; and Hall, Mayhew and van Zwam's excluded-minor characterisation of the near-regular matroids, that is, the matroids representable over all fields with at least three elements [9].

The immediate problem that looms large is that of finding the excluded minors for the class of $\text{GF}(5)$ -representable matroids. While this problem is beyond the range of current techniques, a road map for an attack is outlined in [16]. In essence, this road map reduces the problem to a finite sequence of problems of a type that we now describe. First note that regular matroids and many other naturally arising classes of representable matroids such as near-regular, dyadic and $\sqrt[6]{1}$ -matroids [22] can be described as classes of matroids representable over an algebraic structure called a *partial field*. We wish to find the excluded-minor characterisation for the class of \mathbb{P} -representable matroids for some fixed partial field \mathbb{P} . We have a 3-connected matroid N with the property that every \mathbb{P} -representation of N extends *uniquely* to a \mathbb{P} -representation of any 3-connected \mathbb{P} -representable matroid having N as a minor. Such a matroid N is called a *strong stabilizer* for the class of \mathbb{P} -representable matroids. With these ingredients, the goal is to bound the size of an excluded minor for the class of \mathbb{P} -representable matroids having the strong stabilizer N as a minor. This situation is a more general version of the one that arises in the proof of the excluded-minor characterisation of $\text{GF}(4)$ -representable matroids [8]. There, the partial field is $\text{GF}(4)$ and the strong stabilizer is $U_{2,4}$. (See also the introduction to [4] for more detail on this strategy.)

Date: September 21, 2018.

The first, fourth, and fifth authors were supported by the New Zealand Marsden Fund.

Ideally, we would develop techniques that would reduce problems of the above type to routine computation. But an annoying barrier arises. Let N be a matroid. A matroid M is N -fragile if, for all elements e of M , at most one of $M \setminus e$ or M/e has an N -minor. It seems that, for a strong stabilizer N for a partial field \mathbb{P} , to bound the size of an excluded minor for \mathbb{P} -representable matroids that contains N as a minor, we need to have some insight into the structure of \mathbb{P} -representable N -fragile matroids. The goal of this paper is to demonstrate that this is, in essence, the fundamental problem. We prove that if M is an excluded minor for the class of \mathbb{P} -representable matroids having an N -minor, then either the size or rank of M is bounded relative to N , or, up to replacing M by a Δ - Y -equivalent excluded minor, M (or its dual) has a pair of elements $\{a, b\}$ such that $M \setminus a, b$ is an N -fragile (or N^* -fragile) matroid. More specifically, we prove the following:

Theorem 1.1. *Let \mathbb{P} be a partial field, let M be an excluded minor for the class of \mathbb{P} -representable matroids, and let N be a non-binary strong stabilizer for the class of \mathbb{P} -representable matroids, where M has an N -minor. For some matroid M_1 that is Δ - Y -equivalent to M , and some (M', N') in $\{(M_1, N), (M_1^*, N^*)\}$, the matroid M' is an excluded minor having an N' -minor such that at least one of the following holds:*

- (i) $|E(M')| \leq |E(N')| + 9$;
- (ii) $r(M') \leq r(N') + 7$; or
- (iii) *there is a pair $\{a, b\} \subseteq E(M)$ such that $M' \setminus a, b$ is a 3-connected N' -fragile matroid with an N' -minor.*

We defer the definition of the Δ - Y -equivalence to the next section.

Theorem 1.1 tells us that an excluded minor for \mathbb{P} -representable matroids will either have bounded size or will be very close to an N -fragile matroid. Current techniques for bounding the size of an excluded minor in the latter case rely on obtaining explicit information about the structure of N -fragile matroids and this needs to be done on a case-by-case basis. Even for quite simple matroids this can be a difficult problem. Here is an example. Recall the non-Fano matroid F_7^- . The barrier to finding the excluded minors for the class of dyadic matroids is that we do not understand the structure of dyadic F_7^- -fragile matroids and such an understanding seems some way off.

On the other hand, $U_{2,5}$ and $U_{3,5}$ are strong stabilizers for representability over two interesting partial fields and we do know the structure of $U_{2,5}$ - and $U_{3,5}$ -fragile matroids within these classes [7]. The first is the partial field \mathbb{H}_5 which was introduced by Pendavingh and van Zwam [16]. The class of matroids representable over this field is the class obtained by taking the 3-connected matroids that have exactly six inequivalent representations over $\text{GF}(5)$ and closing the class under minors. This class forms the bottom layer of Pendavingh and van Zwam's hierarchy of $\text{GF}(5)$ -representable matroids. Finding excluded minors for this class would be a key first step towards finding the excluded minors for matroids representable over $\text{GF}(5)$.

The other partial field is the 2-regular or 2-uniform partial field, denoted \mathbb{U}_2 . This is a member of a family of partial fields. The matroids representable over \mathbb{U}_0 and \mathbb{U}_1 are the regular and near-regular matroids respectively. Regular matroids are the matroids representable over all fields, and near-regular

matroids are the matroids representable over all fields with at least three elements. Let \mathcal{M}_4 denote the matroids representable over all fields of size at least four. It would certainly be interesting to have a characterisation of the class \mathcal{M}_4 . The class of \mathbb{U}_2 -representable matroids is contained in \mathcal{M}_4 , and it is known [19] that this class is a proper subclass of \mathcal{M}_4 . Nonetheless, knowing the excluded minors for \mathbb{U}_2 would be a key step towards characterising the class \mathcal{M}_4 . The interesting matroids to uncover are the excluded minors for \mathbb{U}_2 that belong to \mathcal{M}_4 . Attention could then be focussed on members of \mathcal{M}_4 having these matroids as minors. It is possible that these will form highly structured classes of bounded branch width.

With the results of this paper, and the characterisation of the \mathbb{U}_2 - and \mathbb{H}_5 -representable $U_{2,5}$ - and $U_{3,5}$ -fragile matroids, there is real hope that obtaining the full list of excluded minors for these classes is an achievable goal. Beyond these classes all bets are off. Experience with graph minors tells us that we must expect to hit a wall quite soon — consider, for example, the excluded minors for the class of toroidal graphs or the class of Δ - Y -reducible graphs [25]. We know from [10] that there are at least 564 excluded minors for $\text{GF}(5)$ -representable matroids. It is possible that obtaining the full list will be forever beyond our reach. But the quest is surely a worthy one.

2. PRELIMINARIES AND THE MAIN THEOREMS

In this section we gather preliminaries that are used throughout the paper. We will then be able to state the main results: Theorems 2.30 and 2.31. In particular, Theorem 2.31 implies Theorem 1.1. Most of the relevant results and terminology on matroid connectivity can either be found in Oxley [12] or in the recent literature on removing elements relative to a fixed basis [3, 14, 24]. The results and terminology on matroid representation theory can be found in [11, 16, 17].

We write “by orthogonality” to refer to the property that a circuit and a cocircuit cannot meet in one element. In the context of partitions of the form $(X, \{e\}, Y)$, we will also write “by orthogonality” to refer to an application of the next lemma.

Lemma 2.1. *Let e be an element of a matroid M , and let $(X, \{e\}, Y)$ be a partition of $E(M)$. Then $e \in \text{cl}(X)$ if and only if $e \notin \text{cl}^*(Y)$.*

Connectivity. The following results are well known.

Lemma 2.2. *Let M be a 3-connected matroid. If X is a rank-2 subset of $E(M)$ and $|X| \geq 4$, then $M \setminus x$ is 3-connected for all $x \in X$.*

Lemma 2.3 (Bixby’s Lemma [2]). *Let M be a 3-connected matroid, and let $e \in E(M)$. Then $\text{si}(M/e)$ or $\text{co}(M \setminus e)$ is 3-connected.*

The next three results state elementary properties of 3-separations that we shall use frequently. We use the notation $e \in \text{cl}^{(*)}(X)$ to mean $e \in \text{cl}(X)$ or $e \in \text{cl}^*(X)$.

Lemma 2.4. *Let X be an exactly 3-separating set in a 3-connected matroid, and suppose that $e \in E(M) - X$. Then $X \cup e$ is 3-separating if and only if $e \in \text{cl}^{(*)}(X)$.*

Lemma 2.5. *Let (X, Y) be an exactly 3-separating partition of a 3-connected matroid M . Suppose $|X| \geq 3$ and $x \in X$. Then*

- (i) $x \in \text{cl}^{(*)}(X - x)$; and
- (ii) $(X - x, Y \cup x)$ is exactly 3-separating if and only if x is in exactly one of $\text{cl}(X - x) \cap \text{cl}(Y)$ and $\text{cl}^*(X - x) \cap \text{cl}^*(Y)$.

Lemma 2.6 ([3, Lemma 2.11]). *Let (X, Y) be a 3-separation of a 3-connected matroid M . If $X \cap \text{cl}(Y) \neq \emptyset$ and $X \cap \text{cl}^*(Y) \neq \emptyset$, then $|X \cap \text{cl}(Y)| = 1$ and $|X \cap \text{cl}^*(Y)| = 1$.*

Let M be a matroid. A 3-separation (X, Y) of M is a *vertical 3-separation* if $\min\{r(X), r(Y)\} \geq 3$. We say that a partition $(X, \{z\}, Y)$ is a *vertical 3-separation* of M when both $(X \cup \{z\}, Y)$ and $(X, Y \cup \{z\})$ are vertical 3-separations and $z \in \text{cl}(X) \cap \text{cl}(Y)$. We will write (X, z, Y) for $(X, \{z\}, Y)$. If (X, z, Y) is a vertical 3-separation of M , then we say that (X, z, Y) is a *cyclic 3-separation* of M^* .

A *path of 3-separations* of M is a partition (P_1, \dots, P_n) of $E(M)$ such that $(P_1 \cup \dots \cup P_i, P_{i+1} \cup \dots \cup P_n)$ is a 3-separation of M for each $i \in \{1, \dots, n-1\}$. In particular, a vertical 3-separation (X, z, Y) is a path of 3-separations.

Lemma 2.7 ([23, Lemma 3.5]). *Let M be a 3-connected matroid, and $e \in E(M)$. The matroid M has a vertical 3-separation (X, e, Y) if and only if $\text{si}(M/e)$ is not 3-connected.*

Let k be a positive integer, and let (P, Q) be a k -separation. We call the set $\text{cl}(P) \cap \text{cl}(Q)$ the *guts* of (P, Q) , and $\text{cl}^*(P) \cap \text{cl}^*(Q)$ the *coguts* of (P, Q) . We also say that an element $z \in \text{cl}(P) \cap \text{cl}(Q)$ is a *guts element*, and $z \in \text{cl}^*(P) \cap \text{cl}^*(Q)$ is a *coguts element*.

We write “by uncrossing” to refer to an application of the next result.

Lemma 2.8. *Let M be a 3-connected matroid, and let X and Y be 3-separating subsets of $E(M)$. Then the following hold.*

- (i) *If $|X \cap Y| \geq 2$, then $X \cup Y$ is 3-separating.*
- (ii) *If $|E(M) - (X \cup Y)| \geq 2$, then $X \cap Y$ is 3-separating.*

Series classes. We will use the following two results on series classes. We omit the easy proof of the first lemma.

Lemma 2.9. *Let M be a matroid such that $\text{co}(M)$ is 3-connected. If S and S' are distinct series classes of M , then either $S \cup S'$ is independent, or $\text{co}(M) \cong U_{1,3}$.*

When S is a series class of size two, we say S is a *series pair*.

Lemma 2.10. *Let M be a 3-connected matroid, and let $u \in E(M)$ be an element such that $\text{co}(M \setminus u)$ is 3-connected and $\text{co}(M \setminus u) \not\cong U_{1,3}$. Let S be a non-trivial series class of $M \setminus u$. If there is some element $s \in S$ such that $\text{si}(M/s)$ is not 3-connected, then*

- (i) $|S| = 2$;
- (ii) $M \setminus u$ has exactly two distinct non-trivial series classes; and
- (iii) $\text{si}(M/s')$ is 3-connected, where $S = \{s, s'\}$.

Proof. Suppose that there is some element $s \in S$ such that $\text{si}(M/s)$ is not 3-connected. Observe that $r_{M^*}(S \cup u) = 2$. It follows that if $|S| \geq 3$, then $M^* \setminus s$, and hence M/s , is 3-connected for all $s \in S$ by Lemma 2.2; a contradiction. So $|S| = 2$, and (i) holds. Henceforth, we let $S = \{s, s'\}$.

We now consider (ii) and (iii). By Lemma 2.7, M has a vertical 3-separation (A, s, B) . Without loss of generality, we may assume that A is coclosed, and that $u \in A$. Then $(A - u, B)$ is a 2-separation of $M/s \setminus u$, and the matroid $M/s \setminus u$ is 3-connected up to series classes because $\text{co}(M \setminus u)$ is 3-connected. Hence $A - u$ or B is contained in a series class of $M \setminus u$. Since $S \cup u$ is a triad containing s , and (A, s, B) is a vertical 3-separation, it follows from orthogonality (see Lemma 2.1) that $S \cup u$ is not contained in $A \cup s$ or $B \cup s$. So $s' \in B$. If B is contained in a series class of $M \setminus u$, then s is also in this series class; a contradiction. So $A - u$ is contained in a series class, distinct from S . As A is coclosed in M , we have that $A - u$ is a series class in $M \setminus u$. Since $s \in \text{cl}(A)$, there is a circuit C of M such that $s \in C \subseteq A \cup s$. Moreover, $u \in C$ by orthogonality with C and the triad $S \cup u$.

Next we claim that $A - u$ and S are the only series classes of $M \setminus u$. Suppose there is some series pair S' of $M \setminus u$ disjoint from $S \cup (A - u)$. Then $S' \cup u$ is a triad of M that meets the circuit C in the single element u ; a contradiction to orthogonality. Thus S and $A - u$ are the only series classes of $M \setminus u$, so (ii) holds.

Finally, suppose that $\text{si}(M/s')$ is not 3-connected. Then, by Lemma 2.7, M has a vertical 3-separation (A', s', B') . We may assume, without loss of generality, that A' is coclosed and that $u \in A'$. By the same argument as used earlier, $A' - u$ is a series class of $M \setminus u$. By (ii), $A' - u = A - u$, and thus $(A', s', B') = (A, s', (B - s') \cup s)$. Then $s' \in \text{cl}(A)$, so there is some circuit C' of M such that $s' \in C' \subseteq A \cup s'$, and $u \in C'$ by orthogonality. But then we have distinct circuits $C \subseteq A \cup s$ and $C' \subseteq A \cup s'$ such that $u \in C \cap C'$. By circuit elimination, there is a circuit C'' of M such that $C'' \subseteq (A - u) \cup S$. Thus, by Lemma 2.9, $\text{co}(M \setminus u) \cong U_{1,3}$, a contradiction. Therefore (iii) holds. \square

Fans. A subset F of the ground set of a matroid, with $|F| \geq 3$, is a *fan* if there is an ordering (f_1, f_2, \dots, f_k) of the elements of F such that

- (a) $\{f_1, f_2, f_3\}$ is either a triangle or a triad, and
- (b) for all $i \in \{1, 2, \dots, k-3\}$, if $\{f_i, f_{i+1}, f_{i+2}\}$ is a triangle, then $\{f_{i+1}, f_{i+2}, f_{i+3}\}$ is a triad, while if $\{f_i, f_{i+1}, f_{i+2}\}$ is a triad, then $\{f_{i+1}, f_{i+2}, f_{i+3}\}$ is a triangle.

When there is no ambiguity, we also say that the ordering (f_1, f_2, \dots, f_k) is a fan. If F has a fan ordering (f_1, f_2, \dots, f_k) where $k \geq 4$, then f_1 and f_k are the *ends* of F , and f_2, f_3, \dots, f_{k-1} are the *internal elements* of F .

Let F be a fan with ordering (f_1, f_2, \dots, f_k) where $k \geq 4$, and let $i \in \{1, 2, \dots, k\}$ if $k \geq 5$, or $i \in \{1, 4\}$ if $k = 4$. An element f_i is a *spoke element* of F if $\{f_1, f_2, f_3\}$ is a triangle and i is odd, or if $\{f_1, f_2, f_3\}$ is a triad and i is even; otherwise f_i is a *rim element*.

We say that a fan F is *maximal* if there is no fan that properly contains F .

We employ the following results when we encounter fans.

Lemma 2.11 ([3, Lemma 2.12]). *Let M be a 3-connected matroid with $|E(M)| \geq 7$. Suppose that M has a fan F of at least 4 elements, and let f be an end of F .*

- (i) *If f is a spoke element, then $\text{co}(M \setminus f)$ is 3-connected and $\text{si}(M/f)$ is not 3-connected.*
- (ii) *If f is a rim element, then $\text{si}(M/f)$ is 3-connected and $\text{co}(M \setminus f)$ is not 3-connected.*

Lemma 2.12 ([3, Lemma 3.3]). *Let M be a matroid with distinct elements f_1, f_2, f_3, f_4 . If the only triangle containing f_3 is $\{f_1, f_2, f_3\}$ and the only triad containing f_2 is $\{f_2, f_3, f_4\}$, then $\text{si}(M/f_3) \cong \text{co}(M \setminus f_2)$.*

Retaining an N -minor. Let M and N be matroids, and let x be an element of M . If $M \setminus x$ has an N -minor, then x is N -deletable. If M/x has an N -minor, then x is N -contractible. If neither $M \setminus x$ nor M/x has an N -minor, then x is N -essential. If x is both N -deletable and N -contractible, then we say that x is N -flexible. A matroid M is N -fragile if M has an N -minor, and no element of M is N -flexible (note that some authors refer to this as “strictly N -fragile”).

For $X \subseteq E(M)$, we will also say that X is N -deletable (or N -contractible) when $M \setminus X$ (or M/X , respectively) has an N -minor.

The next two results give some conditions for when we can keep an N -minor when dealing with 2-separations.

Lemma 2.13 ([3, Lemma 4.3]). *Let N be a 3-connected matroid such that $|E(N)| \geq 4$. If M has an N -minor, then $\text{si}(M)$ has an N -minor.*

Lemma 2.14 ([14, Lemma 2.7]). *Let (X, Y) be a 2-separation of a connected matroid M and let N be a 3-connected minor of M . Then $\{X, Y\}$ has a member S such that $|S \cap E(N)| \leq 1$. Moreover, when $s \in S$,*

- (i) *if M/s is connected, then M/s has an N -minor; and*
- (ii) *if $M \setminus s$ is connected, then $M \setminus s$ has an N -minor.*

Let (X, z, Y) be a vertical 3-separation of a matroid M . Then (X, Y) is a 2-separation of M/z such that $|X| \geq 3$ and $|Y| \geq 3$. Let N be a 3-connected matroid with $|E(N)| \geq 3$. If M/z has an N -minor, then it follows from Lemma 2.14 that either $|X \cap E(N)| \leq 1$ or $|Y \cap E(N)| \leq 1$. When $|X \cap E(N)| \leq 1$, we refer to X as the *non- N -side* and Y as the *N -side* of (X, z, Y) .

The following result is a routine upgrade of [3, Lemma 4.5] that also covers the case when the N -side of the vertical 3-separation is not closed.

Lemma 2.15. *Let N be a 3-connected minor of a 3-connected matroid M . Let (X, z, Y) be a vertical 3-separation of M such that M/z has an N -minor, where $|X \cap E(N)| \leq 1$. Then, every element of X is either N -deletable or N -contractible in M/z . In particular, letting $Y' = \text{cl}_M(Y) - z$,*

- (i) *every element of $X - Y'$ is N -contractible in M/z , and*
- (ii) *at most one element of X is not N -deletable; moreover, if such an element x exists, then $x \in \text{cl}_M^*(Y') - Y'$ and $z \in \text{cl}_M(X - (Y' \cup x))$.*

Proof. It is immediate from the proof of [3, Lemma 4.5] that the lemma holds when $Y \cup z$ is closed; in particular, (i) holds. We may therefore assume that $Y \cup z$ is not closed. Let $s \in X \cap \text{cl}_M(Y)$. We first show that s is N -deletable. Since (X, Y) is a 2-separation of the connected matroid M/z and $s \in \text{cl}_{M/z}(X) \cap \text{cl}_{M/z}(Y)$, it follows that $M/z \setminus s$ is connected. Then, by Lemma 2.14, $M/z \setminus s$ has an N -minor, so s is N -deletable. Thus any element of X that is not N -deletable belongs to $X - Y'$.

2.15.1. *The partition $(X - s, z, Y \cup s)$ is a vertical 3-separation of M .*

Subproof. By Lemma 2.5(i), $s \in \text{cl}_M^{(*)}(X - s)$. Since $s \in \text{cl}_M(Y)$ it follows from orthogonality that $s \notin \text{cl}_M^*(X - s)$. Therefore $s \in \text{cl}_M(X - s)$ and, by Lemma 2.5(ii), $(X - s, Y \cup \{s, z\})$ is exactly 3-separating. By a similar argument, $((X - s) \cup z, Y \cup s)$ is exactly 3-separating. As $s \in \text{cl}_M(X - s)$, we see that $r(X - s) = r(X) \geq 3$. Hence the partition $(X - s, z, Y \cup s)$ is a vertical 3-separation of M . \square

By repeatedly applying 2.15.1, we see that $(X - Y', z, Y')$ is a vertical 3-separation of M with $|(X - Y') \cap E(N)| \leq 1$. As $Y' \cup z$ is closed, (ii) holds. The fact that each element of X is either N -deletable or N -contractible now follows from Lemma 2.6. \square

The next result is a consequence of Lemma 2.15 and Bixby's Lemma.

Lemma 2.16. *Let N be a 3-connected minor of a 3-connected matroid M . Let (X, z, Y) be a vertical 3-separation of M such that M/z has an N -minor, $|X \cap E(N)| \leq 1$, and $Y \cup z$ is closed. Then there is at most one element of X that is not N -flexible. Moreover, if $s \in X$ is not N -flexible, then s is N -contractible and $\text{si}(M/s)$ is 3-connected.*

Representation Theory. A *partial field* is a pair (R, G) , where R is a commutative ring with unity, and G is a subgroup of the group of units of R such that $-1 \in G$. If $\mathbb{P} = (R, G)$ is a partial field, then we write $p \in \mathbb{P}$ whenever $p \in G \cup \{0\}$.

Let \mathbb{P} be a partial field, and let A be an $X \times Y$ matrix with entries from \mathbb{P} . Then A is a \mathbb{P} -matrix if every subdeterminant of A is contained in \mathbb{P} . If $X' \subseteq X$ and $Y' \subseteq Y$, then we write $A[X', Y']$ to denote the submatrix of A induced by X' and Y' . When X and Y are disjoint, if $Z \subseteq X \cup Y$, then we denote by $A[Z]$ the submatrix induced by $X \cap Z$ and $Y \cap Z$, and we denote by $A - Z$ the submatrix induced by $X - Z$ and $Y - Z$.

Theorem 2.17 ([16, Theorem 2.8]). *Let \mathbb{P} be a partial field, and let A be an $X \times Y$ \mathbb{P} -matrix, where X and Y are disjoint. Let*

$$\mathcal{B} = \{X\} \cup \{X \Delta Z : |X \cap Z| = |Y \cap Z|, \det(A[Z]) \neq 0\}.$$

Then \mathcal{B} is the set of bases of a matroid on $X \cup Y$.

We say that the matroid in Theorem 2.17 is \mathbb{P} -representable, and that A is a \mathbb{P} -representation of M . We write $M = M[I|A]$ if A is a \mathbb{P} -matrix, and M is the matroid whose bases are described in Theorem 2.17.

Let A be an $X \times Y$ \mathbb{P} -matrix, with $X \cap Y = \emptyset$, and let $x \in X$ and $y \in Y$ such that $A_{xy} \neq 0$. Then we define A^{xy} to be the $(X \Delta \{x, y\}) \times (Y \Delta \{x, y\})$

\mathbb{P} -matrix given by

$$(A^{xy})_{uv} = \begin{cases} A_{xy}^{-1} & \text{if } uv = yx \\ A_{xy}^{-1}A_{xv} & \text{if } u = y, v \neq x \\ -A_{xy}^{-1}A_{uy} & \text{if } v = x, u \neq y \\ A_{uv} - A_{xy}^{-1}A_{uy}A_{xv} & \text{otherwise.} \end{cases}$$

We say that A^{xy} is obtained from A by *pivoting* on xy . Note that A^{xy} is a \mathbb{P} -matrix, by [18, Proposition 3.3].

Two \mathbb{P} -matrices are *scaling equivalent* if one can be obtained from the other by repeatedly scaling rows and columns by non-zero elements of \mathbb{P} . Two \mathbb{P} -matrices are *geometrically equivalent* if one can be obtained from the other by a sequence of the following operations: scaling rows and columns by non-zero entries of \mathbb{P} , permuting rows, permuting columns, and pivoting.

Let \mathbb{P} be a partial field, and let M and N be matroids such that N is a minor of M . Suppose that the ground set of N is $X' \cup Y'$, where X' is a basis of N . We say that M is \mathbb{P} -*stabilized by* N if, whenever A_1 and A_2 are $X \times Y$ \mathbb{P} -matrices, with $X' \subseteq X$ and $Y' \subseteq Y$, such that

- (i) $M = M[I|A_1] = M[I|A_2]$,
- (ii) $A_1[X', Y']$ is scaling equivalent to $A_2[X', Y']$, and
- (iii) $N = M[I|A_1[X', Y']] = M[I|A_2[X', Y']]$,

then A_1 is scaling equivalent to A_2 . If M is \mathbb{P} -stabilized by N , and every \mathbb{P} -representation of N extends to a \mathbb{P} -representation of M , then we say M is *strongly \mathbb{P} -stabilized by* N .

Let \mathcal{M} be a class of matroids. We say that N is a \mathbb{P} -*stabilizer for* \mathcal{M} if, for every 3-connected \mathbb{P} -representable matroid $M \in \mathcal{M}$ with an N -minor, M is \mathbb{P} -stabilized by N . We say that N is a *strong \mathbb{P} -stabilizer for* \mathcal{M} if, for every 3-connected \mathbb{P} -representable matroid $M \in \mathcal{M}$ with an N -minor, M is strongly \mathbb{P} -stabilized by N . Usually, we will be interested in the class of \mathbb{P} -representable matroids for some partial field \mathbb{P} . In this case, when it is clear from context, we will simply say “ N is a strong \mathbb{P} -stabilizer”.

Certifying non-representability. Let M be a matroid with a minor N . If M has a pair of elements $\{a, b\}$ such that $M \setminus a, b$ is 3-connected and has an N -minor, then we say $\{a, b\}$ is a *deletion pair with respect to* N . If M has a pair of elements $\{a, b\}$ that are N -deletable, $M \setminus a, b$ is connected, and $\text{co}(M \setminus a)$, $\text{co}(M \setminus b)$, and $\text{co}(M \setminus a, b)$ are 3-connected, then we say $\{a, b\}$ is a *weak deletion pair with respect to* N .

When B is a basis for a matroid M , we write B^* to denote $E(M) - B$. When $Z \subseteq E(M)$, we write $M_B[Z]$ to denote the minor $M/(B-Z) \setminus (B^* - Z)$.

Throughout the rest of this section, we assume that \mathbb{P} is a partial field.

Theorem 2.18 ([11, Theorem 5.5]). *Let M and N be 3-connected matroids. Suppose M has an N -minor, N is a \mathbb{P} -representable matroid that is a strong \mathbb{P} -stabilizer, and $\{a, b\} \subseteq E(M)$ is a weak deletion pair with respect to N such that $M \setminus a$ and $M \setminus b$ are \mathbb{P} -representable. Let D be an $X_N \times Y_N$ \mathbb{P} -matrix such that $N = M[I|D]$. Choose $B, E_N \subseteq E(M) - \{a, b\}$ such that B is a basis of $M \setminus \{a, b\}$, $X_N \subseteq B$, and $M_B[E_N] = N$. Then there exists a $B \times B^*$ matrix A with entries in \mathbb{P} such that*

- (i) $A - a$ and $A - b$ are \mathbb{P} -matrices,

- (ii) $M[I|A - a] = M \setminus a$ and $M[I|A - b] = M \setminus b$, and
- (iii) $A[E_N]$ is scaling equivalent to D .

Moreover, the matrix A is unique up to row and column scaling.

Usually, we will apply Theorem 2.18 to a matroid M that is not \mathbb{P} -representable. We call the matrix A a “companion matrix” for M .

Definition. Let M be a matroid and let $E(M) = X \cup Y$ where X and Y are disjoint. Let A be an $X \times Y$ matrix with entries in \mathbb{P} such that, for some distinct $a, b \in Y$, both $A - a$ and $A - b$ are \mathbb{P} -matrices, $M \setminus a = M[I|A - a]$, and $M \setminus b = M[I|A - b]$. Then A is an $X \times Y$ *companion \mathbb{P} -matrix* for M .

Let M be an excluded minor for the class of \mathbb{P} -representable matroids. Then it is easily seen that M is 3-connected. By Theorem 2.18, given a 3-connected matroid N that is a minor of M and a strong \mathbb{P} -stabilizer, a \mathbb{P} -representation for N , and a weak deletion pair, there is a $B \times B^*$ companion \mathbb{P} -matrix A for M , where B is an appropriately chosen basis of M .

A companion matrix for an excluded minor contains a certificate of non-representability over \mathbb{P} .

Definition. Let B be a basis of M , and let A be a $B \times B^*$ matrix with entries in \mathbb{P} . A subset Z of $E(M)$ *incriminates* the pair (M, A) if $A[Z]$ is square and one of the following holds:

- (i) $\det(A[Z]) \notin \mathbb{P}$,
- (ii) $\det(A[Z]) = 0$ but $B \triangle Z$ is a basis of M , or
- (iii) $\det(A[Z]) \neq 0$ but $B \triangle Z$ is dependent in M .

The next result follows immediately from the definition.

Lemma 2.19. *Let M be a matroid, let A be an $X \times Y$ matrix with entries in \mathbb{P} , where X and Y are disjoint, and $X \cup Y = E(M)$. Exactly one of the following statements is true:*

- (i) A is a \mathbb{P} -matrix and $M = M[I|A]$, or
- (ii) there is some $Z \subseteq X \cup Y$ that incriminates (M, A) .

The next theorem shows that there is some companion matrix A' for M that has a 4-element incriminating set.

Theorem 2.20 ([11, Theorem 5.8]). *Let M be a matroid, let A be an $X \times Y$ companion \mathbb{P} -matrix for M , let $a, b \in Y$, and suppose that $Z \subseteq X \cup Y$ incriminates (M, A) . Then there is some $X' \times Y'$ matrix A' , and $x, y \in X'$, such that*

- (i) $a, b \in Y'$,
- (ii) $A - a$ is geometrically equivalent to $A' - a$,
- (iii) $A - b$ is geometrically equivalent to $A' - b$, and
- (iv) $\{x, y, a, b\}$ incriminates (M, A') .

Let N be a 3-connected non-binary matroid. A matroid M with an N -minor is N -stable if, whenever (X, Y) is a 2-separation of M where X is the non- N -side, then the matroid M_X , corresponding to X in the 1- or 2-sum decomposition of M induced by (X, Y) , is binary.

The following result is proved by Hall, Mayhew, and van Zwam [9, Propositions 3.1 and 3.2].

Lemma 2.21. *Let N be a 3-connected strong \mathbb{P} -stabilizer that is non-binary, and let M be a \mathbb{P} -representable matroid that has an N -minor. If M is N -stable, then M is strongly \mathbb{P} -stabilized by N .*

We next consider how a matroid can lose the property of being N -stable after a single-element extension. We say that a matroid M is 3-connected up to series pairs if $\text{co}(M)$ is 3-connected and every non-trivial series class of M is a series pair.

Lemma 2.22. *Let N be a 3-connected non-binary matroid. Let M be a matroid with an element e such that $M \setminus e$ has an N -minor, where e is not a coloop. Suppose that $M \setminus e$ is 3-connected up to series pairs, and that M is not N -stable. Then M has a 2-separation $(S \cup e, Q)$ where $S \cup e$ is a triangle and a triad, for some series pair S of $M \setminus e$.*

Proof. Suppose $M \setminus e$ is 3-connected up to series pairs and M is not N -stable. If M is 3-connected up to series pairs, then it is trivially N -stable; a contradiction. So there is some 2-separation (A, B) of M with $e \in A$, say. Since $(A - e, B)$ is 2-separating in $M \setminus e$, and $M \setminus e$ is 3-connected up to series pairs, we deduce that $|A| \leq 3$. Let $M = Q_A \oplus_2 Q_B$ be the 2-sum decomposition corresponding to the 2-separation (A, B) of M . Since $M \setminus e$ has an N -minor and $|E(N)| \geq 4$, we have $|A \cap E(N)| \leq 1$. As M is not N -stable, Q_A has a $U_{2,4}$ -minor. But $|A| \leq 3$, so $Q_A \cong U_{2,4}$. The result follows. \square

Definition. Let M be a matroid with a 2-separation (P, Q) where P is a triangle and a triad. Then P is an *unstable triple* of M .

Let N be a 3-connected non-binary matroid, and let M be a connected matroid with $e \in E(M)$ such that $M \setminus e$ has an N -minor. By Lemma 2.22, if $M \setminus e$ is 3-connected up to series pairs, but M is not N -stable, then M has an unstable triple, which contains e .

Observe also that if P is an unstable triple, then $P - p$ is a series pair in $M \setminus p$, for each $p \in P$.

The next lemma gives sufficient conditions for showing a certain minor of M is not \mathbb{P} -representable. It can be proved by a straightforward modification of a result of Mayhew, Whittle, and van Zwam [11, Theorem 5.12]. The conditions (iv) and (v) are changed from “ $M_B[Z_1]$ and $M_B[Z_2]$ are 3-connected up to series-parallel classes” to “ $M_B[Z_1]$ and $M_B[Z_2]$ are N -stable”, using Lemma 2.21.

Lemma 2.23. *Let M be a matroid, let A be a $B \times B^*$ matrix with entries in \mathbb{P} , where $\{x, y, a, b\}$ incriminates (M, A) for $x, y \in B$ and $a, b \in B^*$. Let N be a non-binary strong stabilizer for the class of \mathbb{P} -representable matroids. Suppose that $C \subseteq E(M)$ is such that $M_B[C]$ is N -fragile. If there exist subsets $Z, Z_1, Z_2 \subseteq E(M)$ such that*

- (a) $a \in Z_1 - Z_2$ and $b \in Z_2 - Z_1$,
- (b) $C \cup \{x, y\} \subseteq Z \subseteq Z_1 \cap Z_2$,
- (c) $M_B[Z]$ is connected,
- (d) $M_B[Z_1]$ is N -stable,
- (e) $M_B[Z_2]$ is N -stable, and

(f) $\{x, y, a, b\}$ incriminates $(M_B[Z_1 \cup Z_2], A[Z_1 \cup Z_2])$,
then $M_B[Z_1 \cup Z_2]$ is not strongly \mathbb{P} -stabilized by N .

The following special case of Lemma 2.23 is sufficient for our needs.

Lemma 2.24. *Let M be a matroid, let $E = E(M)$, let A be a $B \times B^*$ matrix with entries in \mathbb{P} , where $\{x, y, a, b\}$ incriminates (M, A) for $x, y \in B$ and $a, b \in B^*$. Let N be a non-binary strong stabilizer for the class of \mathbb{P} -representable matroids. If there exists $u \in E - \{x, y, a, b\}$ such that*

- (a) $M_B[E - \{a, b, u\}]$ is connected and has an N -minor, and
- (b) $M_B[E - \{b, u\}]$ and $M_B[E - \{a, u\}]$ are N -stable,

then $M_B[E - u]$ is not strongly \mathbb{P} -stabilized by N .

We write ‘‘by an allowable pivot’’ to refer to an application of either of the next two results.

Lemma 2.25 ([11, Lemma 5.10]). *Let A be a $B \times B^*$ companion \mathbb{P} -matrix for M . Suppose that $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$. If $p \in \{x, y\}$, $q \in B^* - \{a, b\}$, and $A_{pq} \neq 0$, then $\{x, y, a, b\} \triangle \{p, q\}$ incriminates (M, A^{pq}) .*

Lemma 2.26 ([11, Lemma 5.11]). *Let A be a $B \times B^*$ companion \mathbb{P} -matrix for M . Suppose that $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$. If $p \in B - \{x, y\}$, $q \in B^* - \{a, b\}$, $A_{pq} \neq 0$, and either $A_{pa} = A_{pb} = 0$ or $A_{xq} = A_{yq} = 0$, then $\{x, y, a, b\}$ incriminates (M, A^{pq}) .*

The elements of a set $\{x, y, a, b\}$ that incriminates (M, A) label a 2×2 submatrix $A[\{x, y, a, b\}]$ of A . We will refer to the next result by saying ‘‘the bad submatrix has no zero entries.’’

Lemma 2.27. *Let A be a $B \times B^*$ companion \mathbb{P} -matrix for M . Suppose that $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$. Then $A_{ij} \neq 0$ for $i \in \{x, y\}$ and $j \in \{a, b\}$.*

Proof. Towards a contradiction, suppose that $A_{ij} = 0$ for some $i \in \{x, y\}$ and $j \in \{a, b\}$. We may assume without loss of generality that $A_{xb} = 0$. Then $\det(A[\{x, y, a, b\}]) \in \mathbb{P}$. Since $\{x, y, a, b\}$ incriminates the pair (M, A) , it follows that either

- (i) $\det(A[\{x, y, a, b\}]) = 0$ but $B \triangle \{x, y, a, b\}$ is a basis of M , or
- (ii) $\det(A[\{x, y, a, b\}]) \neq 0$ but $B \triangle \{x, y, a, b\}$ is dependent in M .

Assume that (i) holds. As $\det(A[\{x, y, a, b\}]) = A_{xa} \cdot A_{yb} = 0$ and non-zero elements of \mathbb{P} are units, it follows that $A_{xa} = 0$ or $A_{yb} = 0$. Suppose that $A_{xa} = 0$. Let $B' = B \triangle \{x, y, a, b\}$. Now B and B' are bases of M and $x \in B - B'$, so, by basis exchange, there is some $z \in B' - B = \{a, b\}$ such that $(B - x) \cup z$ is a basis of M . This is a contradiction because $M \setminus b = M[I|A - b]$, $M \setminus a = M[I|A - a]$ and $A_{xa} = A_{xb} = 0$, so both $(B - x) \cup a$ and $(B - x) \cup b$ are dependent in M . Thus $A_{xa} \neq 0$. Similarly, since $a \in B' - B$, it follows that $(B' - a) \cup x$ or $(B' - a) \cup y$ is a basis of $M \setminus a = M[I|A - a]$. Thus $A_{yb} \neq 0$. We deduce that (i) does not hold.

Therefore (ii) holds. Since $\det(A[\{x, y, a, b\}]) = A_{xa} \cdot A_{yb} \neq 0$, it follows that $A_{xa} \neq 0$ and $A_{yb} \neq 0$. Now $M \setminus b = M[I|A - b]$ and $A_{xa} \neq 0$, so $(B - x) \cup a$ is a basis of M . Similarly, $M \setminus a = M[I|A - a]$ and $A_{yb} \neq 0$, so

$(B - y) \cup b$ is also a basis of M . Let $B_1 = (B - x) \cup a$ and $B_2 = (B - y) \cup b$. Then $x \in B_2 - B_1$, so, by basis exchange, there is some $z \in B_1 - B_2$ such that $(B_2 - x) \cup z$ is a basis of M . But $B_1 - B_2 = \{a, y\}$, so either $(B - x) \cup b$ or B' is a basis. In the former case, since $A_{xb} = 0$, it follows that $(B - x) \cup b$ is dependent in $M \setminus a = M[I|A - a]$ and hence in M . Since B' is dependent by assumption, we obtain a contradiction, thus completing the proof. \square

Robust and strong elements. When working with a matroid M and a \mathbb{P} -representation A of M , there is a natural basis B of M that labels the rows of A . We will frequently look to remove elements “relative to B ”; that is, in such a way that we obtain a \mathbb{P} -representation of the minor of M by removing rows and columns of A , without pivoting. This leads to the following definitions.

Let M be a 3-connected matroid, let B be a basis of M , and let N be a 3-connected minor of M . Recall that we write B^* to denote $E(M) - B$. An element $e \in E(M)$ is (N, B) -robust if either

- (i) $e \in B$ and M/e has an N -minor, or
- (ii) $e \in B^*$ and $M \setminus e$ has an N -minor.

Note that an N -flexible element of M is clearly (N, B) -robust for any basis B of M .

An element $e \in E(M)$ is (N, B) -strong if either

- (i) $e \in B$, and $\text{si}(M/e)$ is 3-connected and has an N -minor; or
- (ii) $e \in B^*$, and $\text{co}(M \setminus e)$ is 3-connected and has an N -minor.

Delta-wye exchange. Let M be a matroid with a triangle $T = \{a, b, c\}$. Consider a copy of $M(K_4)$ having T as a triangle with $\{a', b', c'\}$ as the complementary triad labelled such that $\{a, b', c'\}$, $\{a', b, c'\}$ and $\{a', b', c\}$ are triangles. Let $P_T(M, M(K_4))$ denote the generalised parallel connection of M with this copy of $M(K_4)$ along the triangle T . Let M' be the matroid $P_T(M, M(K_4)) \setminus T$ where the elements a', b' and c' are relabelled as a, b and c respectively. The matroid M' is said to be obtained from M by a Δ - Y exchange on the triangle T , and is denoted $\Delta_T(M)$. Dually, M'' is obtained from M by a Y - Δ exchange on the triad $T^* = \{a, b, c\}$ if $(M'')^*$ is obtained from M^* by a Δ - Y exchange on T^* . The matroid M'' is denoted $\nabla_{T^*}(M)$.

We say that a matroid M_1 is Δ - Y -equivalent to a matroid M_0 if M_1 can be obtained from M_0 by a sequence of Δ - Y and Y - Δ exchanges.

Oxley, Semple, and Vertigan proved that excluded minors for the class of \mathbb{P} -representable matroids are closed under Δ - Y exchange.

Proposition 2.28 ([13, Theorem 1.1]). *Let \mathbb{P} be a partial field, and let M be an excluded minor for the class of \mathbb{P} -representable matroids. If M' is Δ - Y -equivalent to M , then M' is an excluded minor for the class of \mathbb{P} -representable matroids.*

Detachable pairs. Let M be a 3-connected matroid, and let N be a 3-connected minor of M . A pair $\{a, b\} \subseteq E(M)$ is N -detachable if either $M \setminus a, b$ or $M/a, b$ is 3-connected and has an N -minor. A 4-element subset of $E(M)$ is a *quad* if it is a circuit and a cocircuit of M . When $P \subseteq E(M)$ is an exactly 3-separating set of M with a partition $\{L_1, \dots, L_t\}$ for $t \geq 3$ such that

- (a) $|L_i| = 2$ for each $i \in \{1, \dots, t\}$,
- (b) $L_i \cup L_j$ is a quad for all distinct $i, j \in \{1, \dots, t\}$, and
- (c) L_i is not contained in a triangle or a triad, for each $i \in \{1, \dots, t\}$,

then P is a *spike-like 3-separator* of M .

Brettell, Whittle, and Williams [4–6] proved that either M has a spike-like 3-separator, or, after performing at most one Δ - Y or Y - Δ exchange on M , we obtain a matroid with a detachable pair. More specifically:

Theorem 2.29 ([4, Theorem 1.1]). *Let M be a 3-connected matroid, and let N be a 3-connected minor of M such that $|E(N)| \geq 4$ and $|E(M)| - |E(N)| \geq 10$. Then either*

- (i) M has an N -detachable pair,
- (ii) there is a matroid M' obtained by performing a single Δ - Y or Y - Δ exchange on M such that M' has an N -detachable pair, or
- (iii) there is a spike-like 3-separator P of M such that at most one element of $E(M) - E(N)$ is not in P .

We note that our definition of a spike-like 3-separator is more restrictive than that which appears in [4], where condition (c) did not appear. However, if M has a spike-like 3-separator for which (c) does not hold, then either (i) or (ii) of Theorem 2.29 holds by [4, Theorem 3.2].

Now let \mathbb{P} be a partial field, let N be a 3-connected strong \mathbb{P} -stabilizer for the class of \mathbb{P} -representable matroids, and let M be an excluded minor for the class of \mathbb{P} -representable matroids. Then M is 3-connected. The results in this paper rely on the existence of a pair of elements $\{a, b\}$ such that $M \setminus a, b$ is 3-connected with an N -minor. By Theorem 2.29, we can guarantee such a pair exists, up to dualising and performing at most one Δ - Y or Y - Δ exchange, unless M has a spike-like 3-separator. We address the possibility of M having a spike-like 3-separator in Section 7.

The main theorems. Let M be an excluded minor for the class of \mathbb{P} -representable matroids for some partial field \mathbb{P} , and let N be a 3-connected strong \mathbb{P} -stabilizer. Let $\{a, b\}$ be a pair of elements of M such that $M \setminus a, b$ is 3-connected with an N -minor. Our first theorem describes, in the case that $M \setminus a, b$ is not N -fragile and $|E(M)| > |E(N)| + 9$, the local structure of $M \setminus a, b$ for any such deletion pair $\{a, b\}$.

Theorem 2.30. *Let M be an excluded minor for the class of \mathbb{P} -representable matroids, and let N be a non-binary 3-connected strong \mathbb{P} -stabilizer for the class of \mathbb{P} -representable matroids. Suppose M has a pair of elements $\{a, b\}$ such that $M \setminus a, b$ is 3-connected with an N -minor. Then either*

- (i) $|E(M)| \leq |E(N)| + 9$, or
- (ii) M has a $B \times B^*$ companion \mathbb{P} -matrix A for which $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$, and either
 - (a) $M \setminus a, b$ is N -fragile, and $M \setminus a, b$ has at most one (N, B) -robust element u outside of $\{x, y\}$, where if such an element u exists, then $u \in B^* - \{a, b\}$ is an (N, B) -strong element of $M \setminus a, b$, and $\{u, x, y\}$ is a coclosed triad of $M \setminus a, b$, or
 - (b) $M \setminus a, b$ is not N -fragile, but there is an element $u \in B^* - \{a, b\}$ that is (N, B) -strong in $M \setminus a, b$; either

- (I) the N -flexible, and (N, B) -robust, elements of $M \setminus a, b$ are contained in $\{u, x, y\}$, or
 - (II) the N -flexible, and (N, B) -robust, elements of $M \setminus a, b$ are contained in $\{u, x, y, z\}$, where $z \in B$, and (z, u, x, y) is a maximal fan of $M \setminus a, b$, or
 - (III) the N -flexible, and (N, B) -robust, elements of $M \setminus a, b$ are contained in $\{u, x, y, z, w\}$, where $z \in B$, $w \in B^*$, and (w, z, x, u, y) is a maximal fan of $M \setminus a, b$;
- the unique triad in $M \setminus a, b$ containing u is $\{u, x, y\}$; and M has a cocircuit $\{x, y, u, a, b\}$ and a triangle $\{d, x, y\}$ for some $d \in \{a, b\}$.

If M is sufficiently larger than N , then up to performing at most one Δ - Y exchange, we can eliminate case (ii)(b) of Theorem 2.30 by choosing a different deletion pair. (Recall that excluded minors for the class of \mathbb{P} -representable matroids are closed under Δ - Y exchange by Proposition 2.28.) This is the second main theorem of this paper, Theorem 2.31. This theorem implies Theorem 1.1, but Theorem 2.31 provides additional information on the existence of (N_0, B) -robust elements in $M_0 \setminus a, b$, and the local structure of $M_0 \setminus a, b$ when an (N_0, B) -robust element exists.

Theorem 2.31. *Let M be an excluded minor for the class of \mathbb{P} -representable matroids, and let N be a non-binary 3-connected strong \mathbb{P} -stabilizer, where M has an N -minor. For some M_1 that is Δ - Y -equivalent to M , and some (M_0, N_0) in $\{(M_1, N), (M_1^*, N^*)\}$, the matroid M_0 is an excluded minor with an N_0 -minor, and at least one of the following holds:*

- (i) $|E(M_0)| \leq |E(N_0)| + 9$;
- (ii) $r(M_0) \leq r(N_0) + 7$; or
- (iii) *there is a pair $\{a, b\} \subseteq E(M)$ such that $M_0 \setminus a, b$ is 3-connected with an N_0 -minor, and $M_0 \setminus a, b$ is N_0 -fragile. Moreover, there is some basis B for M_0 and a $B \times B^*$ companion \mathbb{P} -matrix A for which $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$, $\{a, b\} \subseteq B^*$, and both of the following hold:*
 - (a) $M_0 \setminus a, b$ has at most one (N_0, B) -robust element outside of $\{x, y\}$, and
 - (b) if u is an (N_0, B) -robust element of $M_0 \setminus a, b$, then $u \in B^* - \{a, b\}$, the element u is (N_0, B) -strong in $M_0 \setminus a, b$, and $\{u, x, y\}$ is a triad of $M_0 \setminus a, b$.

The remainder of the paper is structured as follows. In Section 3, we bound the number of (N, B) -strong elements in an excluded minor M with a 3-connected strong stabilizer N and a basis B . In Section 4, we bound $|E(M)|$ relative to $|E(N)|$ in the case where the (N, B) -strong elements are contained in a 4- or 5-element set with particular properties, which we call a “confining set”. In Section 5 we show that elements that are (N, B) -robust but not (N, B) -strong give rise to a structured collection of 3-separations, called a “path of 3-separations”. In Section 6, we use the structure given by the path of 3-separations to bound the number of (N, B) -robust elements and prove Theorem 2.30. In Section 7, we show that $|E(M)|$ is bounded

relative to $|E(N)|$ in the case where the existence of an N -detachable pair cannot be guaranteed. Finally, in Section 8, we prove Theorem 2.31.

3. STRONG ELEMENTS

Let \mathbb{P} be a partial field, and let N be a 3-connected strong \mathbb{P} -stabilizer for the class of \mathbb{P} -representable matroids such that N is non-binary; so, in particular, $|E(N)| \geq 4$. Suppose M is an excluded minor for the class of \mathbb{P} -representable matroids, and M has a pair of elements $\{a, b\}$ such that $M \setminus a, b$ is 3-connected with an N -minor. Let A be a $B \times B^*$ companion \mathbb{P} -matrix of M such that $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$. Let $M' = M \setminus a, b$. We work under these assumptions for the entirety of the section.

Recall that an element $e \in E(M')$ is (N, B) -strong if either

- (i) $e \in B$, and $\text{si}(M'/e)$ is 3-connected and has an N -minor; or
- (ii) $e \in B^*$, and $\text{co}(M' \setminus e)$ is 3-connected and has an N -minor.

In this section, we bound the number of (N, B) -strong elements of M' . The main result is that M' has at most two (N, B) -strong elements outside of $\{x, y\}$, and any such elements are in B^* .

Lemma 3.1. *If u is an (N, B) -strong element of M' such that $u \notin \{x, y\}$, then $u \notin B$.*

Proof. Suppose that u is an (N, B) -strong element of M' such that $u \in B - \{x, y\}$. Since u is (N, B) -strong, M'/u is 3-connected up to parallel classes. Moreover, as $M \setminus a$, $M \setminus b$ and M are 3-connected, it follows that $M \setminus a/u$, $M \setminus b/u$, and M/u are 3-connected up to parallel classes, and hence are N -stable. As $M \setminus a/u$ and $M \setminus b/u$ are N -stable, and M'/u is connected, Lemma 2.24 implies that M/u is not strongly \mathbb{P} -stabilized by N . But, as M/u is N -stable, this contradicts Lemma 2.21. \square

A subset G of $E(M)$ is a *segment* if every 3-element subset of G is a triangle. A *cosegment* is a segment of M^* .

Lemma 3.2. *Suppose u is an (N, B) -strong element of M' such that $u \notin \{x, y\}$. If u is in a cosegment G of M' such that $|G| \geq 4$, then $|G| = 4$ and $G \cap B = \{x, y\}$.*

Proof. Let G be a cosegment of M' with $|G| \geq 4$. Since G is a corank-2 set, $|G \cap B^*| \leq 2$. Hence $|G \cap B| \geq |G| - 2$. Since u is (N, B) -strong, $u \in B^*$ by Lemma 3.1. So $M' \setminus u$ has an N -minor, and hence the elements of the series class $G - u$ of $M' \setminus u$ are N -contractible. Suppose that there is some $c \in G$ that is in $B - \{x, y\}$. Then M'/c is 3-connected by the dual of Lemma 2.2, so c is an (N, B) -strong element, contradicting Lemma 3.1. We deduce that $|G| = 4$ and that $G \cap B = \{x, y\}$. \square

The following lemma applies to an (N, B) -strong element u for which $M' \setminus u$ is not only 3-connected up to series classes, but also 3-connected up to series pairs.

Lemma 3.3. *Suppose $u \in B^* - \{a, b\}$ is an (N, B) -strong element of M' such that $M' \setminus u$ is 3-connected up to series pairs. Then at least one of $M \setminus a, u$ or $M \setminus b, u$ is not N -stable.*

Proof. Towards a contradiction, suppose that both $M \setminus a, u$ and $M \setminus b, u$ are N -stable. Then, as $M \setminus a, b, u$ is connected, Lemma 2.24 implies that $M \setminus u$ is not strongly \mathbb{P} -stabilized by N .

We claim that $M \setminus u$ is N -stable. Suppose that $M \setminus a, u$ is not 3-connected up to series pairs. Then, as $M \setminus a, b, u$ is 3-connected up to series pairs, and $M \setminus a$ is 3-connected, b is in a parallel pair of $\text{co}(M \setminus a, u)$, which does not exist in $M \setminus a$. Hence, there is a triangle $S \cup b$ of M , where S is a series pair of $M \setminus a, u$. Now $S \cup b$ is 2-separating in $M \setminus a, u$. Since $M \setminus a, u$ is N -stable, the $S \cup b$ component in the 2-sum decomposition of $M \setminus a, u$ does not have a $U_{2,4}$ -minor. It follows that b is in the guts of a 2-separation (S, T) where S is a series pair of $M \setminus a, u$. We deduce that either $M \setminus a, u$ is 3-connected up to series pairs, or b is in the guts of some 2-separation (S, T) of $M \setminus a, u$ where S is a series pair of $M \setminus a, u$. By symmetry, either $M \setminus b, u$ is 3-connected up to series pairs, or a is in the guts of some 2-separation (S', T') of $M \setminus b, u$ where S' , say, is a series pair of $M \setminus b, u$. It now follows, by Lemma 2.22, that $M \setminus u$ is N -stable.

By Lemma 2.21, $M \setminus u$ is strongly \mathbb{P} -stabilized by N ; a contradiction. \square

Let M_1 be a minor of M where, for some $e \in E(M_1)$, the matroid $M_1 \setminus e$ has an N -minor and is 3-connected up to series classes, but M_1 is not N -stable. Recall that, by Lemma 2.22, the matroid M_1 has an unstable triple $S \cup e$, where S is a series pair of $M_1 \setminus e$.

If M' has an (N, B) -strong element $u \in B^* - \{a, b\}$ where $M' \setminus u$ is 3-connected up to series pairs, then it follows from Lemma 3.3 that, up to swapping a and b , the matroid $M \setminus a, u$ has an unstable triple containing b .

We now show that the intersection of an unstable triple with B is a non-empty subset of $\{x, y\}$.

Lemma 3.4. *Suppose $u \in B^* - \{a, b\}$ is an (N, B) -strong element of M' such that $M' \setminus u$ is 3-connected up to series pairs. Then $M' \setminus u$ has a series pair S such that $\emptyset \subsetneq S \cap B \subseteq \{x, y\}$. Moreover, $S \cup b$ is an unstable triple of $M \setminus a, u$, up to swapping a and b .*

Proof. By Lemma 3.3, either $M \setminus a, u$ or $M \setminus b, u$ is not N -stable. Without loss of generality, we may assume that $M \setminus b, u$ is not N -stable. Then, by Lemma 2.22, there is a pair S such that $S \cup a$ is an unstable triple in $M \setminus b, u$. Let $S = \{s_1, s_2\}$. Note that, since S is a series pair of $M' \setminus u$, both s_1 and s_2 are N -contractible in M' . We also note that $S \cap B$ is non-empty because, in $M' \setminus u$, the pair S is codependent and $B^* - \{a, b, u\}$ is a cobasis.

Towards a contradiction, suppose that $s_1 \in B - \{x, y\}$. Then s_1 is not (N, B) -strong by Lemma 3.1, so $\text{si}(M'/s_1)$ is not 3-connected. Hence $\text{si}(M'/s_2)$ is 3-connected by Lemma 2.10, so it follows from Lemma 3.1 that either $s_2 \in \{x, y\}$ or $s_2 \in B^* - \{a, b\}$.

3.4.1. *Up to an allowable pivot, we can assume that $s_2 \in \{x, y\}$.*

Subproof. Observe that since S is a series pair of $M' \setminus u$ but M' is 3-connected, $S \cup u$ is a triad in M' . Suppose that $s_2 \in B^* - \{a, b\}$. Then $A_{s_1 s_2} \neq 0$ because $\{s_1, s_2, u\}$ is a triad of M' . If $A_{x s_2} = A_{y s_2} = 0$, then a pivot on $A_{s_1 s_2}$ is allowable, and s_2 is an $(N, B \Delta \{s_1, s_2\})$ -strong element with $s_2 \in (B \Delta \{s_1, s_2\}) - \{x, y\}$, which contradicts Lemma 3.1. Thus we

shall assume that $A_{xs_2} \neq 0$. Then a pivot on A_{xs_2} is an allowable pivot, and s_2 takes the place of x as a member of the set $\{s_2, y, a, b\}$ that incriminates (M, A^{xs_2}) . \square

By 3.4.1 we may assume that $s_2 = x$. Since $\{a, s_1, s_2\}$ is an unstable triple of $M \setminus b, u$, it follows that $a \in \text{cl}_M(\{s_1, s_2\})$ where $\{s_1, s_2\} \subseteq B$. Hence $A_{ja} \neq 0$ if and only if $j \in \{s_1, s_2\}$. But then $A_{ya} = 0$, contradicting that the bad submatrix has no zero entries. This contradiction arose from the assumption that some member of $S \cap B$ was outside of $\{x, y\}$. Therefore $S \cap B \subseteq \{x, y\}$. \square

Lemma 3.5. *Let u and v be distinct (N, B) -strong elements of M' outside of $\{x, y\}$ such that both $M' \setminus u$ and $M' \setminus v$ are 3-connected up to series pairs. Then at least one of $M \setminus a, u$ or $M \setminus a, v$ is N -stable.*

Proof. Suppose that both $M \setminus a, u$ and $M \setminus a, v$ are not N -stable. By Lemma 2.22, there is a series pair S_u of $M' \setminus u$ such that $S_u \cup b$ is an unstable triple of $M \setminus a, u$, and there is a series pair S_v of $M' \setminus v$ such that $S_v \cup b$ is an unstable triple of $M \setminus a, v$.

First, suppose that $S_u \cap S_v = \emptyset$. If $u \notin S_v$, then $S_v \subseteq E(M \setminus a) - (S_u \cup u)$, so $b \in \text{cl}_{M \setminus a}(S_v) \subseteq \text{cl}_{M \setminus a}(E(M \setminus a) - (S_u \cup u))$, implying $b \notin \text{cl}_{M \setminus a}^*(S_u \cup u)$. But $S_u \cup b$ is an unstable triple of $M \setminus a, u$, so $b \in \text{cl}_{M \setminus a, u}^*(S_u) = \text{cl}_{M \setminus a}^*(S_u \cup u)$; a contradiction. We deduce that $u \in S_v$ and, by symmetry, $v \in S_u$. Now, as $S_u \cup u$ and $S_v \cup v$ are triads of M' , the set $S_u \cup S_v$ is a 4-element cosegment of M' that contains $\{u, v\}$. This contradicts that S_u is a series pair of $M' \setminus u$.

Next, suppose that $|S_u \cap S_v| = 1$. Then, as $S_u \cup b$ and $S_v \cup b$ are triangles of $M \setminus a$ and M , it follows that $S_u \cup S_v$ is a triangle of M' , and so $\{u, v\} \cup S_u \cup S_v$ is a 5-element fan of M' with rim ends u and v . But then $\text{co}(M' \setminus v)$ is not 3-connected by Lemma 2.11; a contradiction. Therefore $S_u = S_v$. But now $\{u, v\} \cup S_u$ is a 4-element cosegment of M' , contradicting that S_u is a series pair of $M' \setminus u$. We deduce that either $M \setminus a, u$ or $M \setminus a, v$ is N -stable. \square

Lemma 3.6. *If M' has a 4-element cosegment G such that $G \cap B = \{x, y\}$, then M' has no (N, B) -strong elements outside of G .*

Proof. Towards a contradiction, suppose that M' has an (N, B) -strong element v outside of G . By Lemma 3.1, $v \in B^*$. By Lemma 3.2, if v is in a 4-element cosegment G' of M' , then $G \cup G'$ is a cosegment consisting of more than four elements; a contradiction. So $M' \setminus v$ is 3-connected up to series pairs. Hence $M' \setminus v$ has a series pair S such that $\emptyset \subsetneq S \cap B \subseteq \{x, y\}$, by Lemma 3.4. Now, in M' , the triad $S \cup v$ meets the cosegment G , so $r_{M'}^*(G \cup S \cup v) \leq 3$. It follows that $|(G \cup S \cup v) \cap B^*| = 3$. Thus $S \cup v$ intersects G in two elements, implying $r^*(G \cup S \cup v) = 2$; a contradiction. \square

These results are enough to bound the number of (N, B) -strong elements outside of $\{x, y\}$. The bound on the number of (N, B) -strong elements is a key ingredient in many subsequent arguments.

Proposition 3.7. *M' has at most two (N, B) -strong elements outside of $\{x, y\}$.*

Proof. Let u be an (N, B) -strong element of M' outside of $\{x, y\}$. By Lemma 3.1, $u \in B^*$. Suppose that $M' \setminus u$ has a series class of size at least

three. Then M' has a 4-element cosegment G such that $\{u, x, y\} \subseteq G$ and $G \cap B = \{x, y\}$, by Lemma 3.2. Thus, by Lemma 3.6, M' has at most two (N, B) -strong elements outside of $\{x, y\}$.

We may now assume that $M' \setminus u$ is 3-connected up to series pairs for each (N, B) -strong element u of M' outside of $\{x, y\}$. Suppose there exist distinct (N, B) -strong elements $u, v_1, v_2 \in B^*$ such that $M' \setminus u, M' \setminus v_1$, and $M' \setminus v_2$ are 3-connected up to series pairs. By Lemma 3.3, we may assume without loss of generality that $M' \setminus b, u$ is not N -stable. Now, by Lemma 3.5, both $M' \setminus b, v_1$ and $M' \setminus b, v_2$ are N -stable. By two further applications of Lemma 3.3, both $M' \setminus a, v_1$ and $M' \setminus a, v_2$ are not N -stable. But this contradicts Lemma 3.5. \square

4. CONFINING SETS

In this section, we work under the following setup. Let \mathbb{P} be a partial field, and let M and N be matroids, where N is a non-binary 3-connected strong \mathbb{P} -stabilizer for the class of \mathbb{P} -representable matroids, and M is an excluded minor for the class of \mathbb{P} -representable matroids with a pair of elements $\{a, b\}$ such that $M \setminus a, b$ is 3-connected with an N -minor. Let $M' = M \setminus a, b$.

We say that a subset G of $E(M')$ is a *confining set* if $G \cap B_1 = \{x_1, y_1\}$ for some basis B_1 of M' , and either

- (a) G is a 4-element cosegment, or
- (b) G is the union of two triads T and T' with $|T \cap T'| = 1$, where $G \cap B_1^*$ has at least one (N, B_1) -strong element,

where x_1 and y_1 are elements of B_1 such that $\{x_1, y_1, a, b\}$ incriminates (M, A_1) for some $B_1 \times B_1^*$ companion \mathbb{P} -matrix A_1 of M . In this case, we also say G is a *confining set relative to B_1* . Note that a confining set satisfying (b) has corank 3 in M' . Every confining set G relative to a basis B_1 has the property that $G \cap B_1^*$ cospans G , since $|G \cap B_1^*| = |G| - 2 = r_{M'}^*(G)$.

We first show that M' either has a confining set, or at most one (N, B) -strong element outside of $\{x, y\}$ for some basis B of M such that $\{x, y, a, b\}$ incriminates (M, A) where A is a $B \times B^*$ companion \mathbb{P} -matrix of M with $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$. We then prove the main result of this section: if M' has a confining set, then $|E(M)|$ is bounded relative to $|E(N)|$.

Proposition 4.1. *Suppose M' does not have a confining set. Then there is some basis B_0 of M' , and $B_0 \times B_0^*$ companion \mathbb{P} -matrix A_0 of M such that $\{x_0, y_0, a, b\}$ incriminates (M, A_0) , for some $\{x_0, y_0\} \subseteq B_0$, and either*

- (i) M' has exactly one (N, B_0) -strong element u outside of $\{x_0, y_0\}$, and $\{u, x_0, y_0\}$ is a triad of M' ; or
- (ii) M' has no (N, B_0) -strong elements outside of $\{x_0, y_0\}$ for every choice of basis B_0 with a $B_0 \times B_0^*$ companion \mathbb{P} -matrix A_0 of M such that $\{x_0, y_0, a, b\}$ incriminates (M, A_0) , for some $\{x_0, y_0\} \subseteq B_0$.

Proof. We first prove the following claim.

4.1.1. *Let B_1 be a basis of M' , and let A_1 be a $B_1 \times B_1^*$ companion \mathbb{P} -matrix of M such that $\{x_1, y_1, a, b\}$ incriminates (M, A_1) , for some $\{x_1, y_1\} \subseteq B_1$. If u is an (N, B_1) -strong element of M' outside of $\{x_1, y_1\}$, then $M' \setminus u$ is 3-connected up to series pairs.*

Subproof. By Lemma 3.1, $u \in B_1^*$. If u is in a cosegment G consisting of at least four elements, then, by Lemma 3.2, G is a confining set of M' ; a contradiction. So we may assume that $M' \setminus u$ is 3-connected up to series pairs for each (N, B_1) -strong element u of M' outside of $\{x_1, y_1\}$. \square

If, for every choice of basis B_1 , with corresponding incriminating set $\{x_1, y_1, a, b\}$, the matroid M' has no (N, B_1) -strong elements outside of $\{x_1, y_1\}$, then clearly the proposition holds. So let B_1 be a basis of M' such that u is an (N, B_1) -strong element of M' outside of $\{x_1, y_1\}$.

4.1.2. *Either the proposition holds, or there is a $B_2 \times B_2^*$ companion \mathbb{P} -matrix A_2 such that $\{x_2, y_2, a, b\}$ incriminates (M, A_2) for some $\{x_2, y_2\} \subseteq B_2$, and M' has exactly two (N, B_2) -strong elements outside of $\{x_2, y_2\}$.*

Subproof. By Proposition 3.7, M' has at most two (N, B_1) -strong elements outside of $\{x_1, y_1\}$. Thus if M' has two (N, B_1) -strong elements outside of $\{x_1, y_1\}$, then 4.1.2 holds with $B_2 = B_1$. So suppose that u is the only (N, B_1) -strong element of M' outside of $\{x_1, y_1\}$. Then $M' \setminus u$ is 3-connected up to series pairs, by 4.1.1. By Lemma 3.3 we may assume, up to swapping a and b , that $M \setminus a, u$ is not N -stable, S_u is a series pair of $M' \setminus u$, and $S_u \cup b$ is an unstable triple of $M \setminus a, u$. Since M' is 3-connected, $S_u \cup u$ is a triad of M' . Thus, if $S_u = \{x_1, y_1\}$, then $\{u, x_1, y_1\}$ is a triad, so the proposition holds in this case. Assume that $S_u \neq \{x_1, y_1\}$. Then it follows from Lemma 3.4 that, without loss of generality, $S_u = \{x_1, s\}$ for some $s \in B_1^* - \{a, b, u\}$. Now b is spanned by S_u in M , and $A_{yb} \neq 0$ because the bad submatrix has no zero entries, so it follows that $A_{ys} \neq 0$. Hence a pivot on A_{ys} is allowable. So $\{x_1, s, a, b\}$ incriminates (M, A_1^{ys}) . Let $B_2 = B_1 \triangle \{y_1, s\}$. If y_1 is not (N, B_2) -strong, then the proposition holds, since $\{u, x_1, s\}$ is a triad. Otherwise, u and y_1 are distinct (N, B_2) -strong elements outside of $\{x_1, s\}$, satisfying 4.1.2. \square

By 4.1.2, we may now assume that B_2 is a basis for M' , the matrix A_2 is a $B_2 \times B_2^*$ companion \mathbb{P} -matrix where $\{x_2, y_2, a, b\}$ incriminates (M, A_2) for some $\{x_2, y_2\} \subseteq B_2$, and M' has exactly two (N, B_2) -strong elements, u and v , in B_2^* . By 4.1.1, $M' \setminus u$ and $M' \setminus v$ are 3-connected up to series pairs. We may assume, up to swapping a and b , that $M \setminus a, u$ and $M \setminus b, v$ are not N -stable, but that $M \setminus b, u$ and $M \setminus a, v$ are N -stable, by Lemmas 3.3 and 3.5. Let S_u be a series pair of $M' \setminus u$, and let S_v be a series pair of $M' \setminus v$, where $S_u \cup b$ is an unstable triple of $M \setminus a, u$, and $S_v \cup a$ is an unstable triple of $M \setminus b, v$. Next, we show that $S_u \cup u = S_v \cup v$. In fact, we prove a more general claim that we can apply even after an allowable pivot.

4.1.3. *Let B_3 be a basis of M' such that $\{x_3, y_3, a, b\}$ incriminates (M, A_3) , for some $B_3 \times B_3^*$ companion \mathbb{P} -matrix A_3 of M and $\{x_3, y_3\} \subseteq B_3$. Suppose M' has exactly two (N, B_3) -strong elements $u, v \in B_3^*$, where $M \setminus a, u$ and $M \setminus b, v$ are not N -stable. Let S_u and S_v be pairs such that $S_u \cup b$ and $S_v \cup a$ are unstable triples of $M \setminus a, u$ and $M \setminus b, v$, respectively. Then $S_u \cup u = S_v \cup v$.*

Subproof. Suppose that the triads $S_u \cup u$ and $S_v \cup v$ of M' are disjoint. Then, by Lemma 3.4, we may assume that $S_u = \{s, x_3\}$ and $S_v = \{t, y_3\}$ for some $s, t \in B_3^* - \{a, b, u, v\}$. Then $A_{y_3s} = 0$ because $S_v \cup v$ is a triad of M' . But then s is spanned by $B_3 - y_3$. Since $S_u \cup b$ is a triangle, it follows that b

is spanned by $B_3 - y_3$. Then $A_{y_3b} = 0$; a contradiction because the bad submatrix has no zero entries.

Let $G = S_u \cup S_v \cup \{u, v\}$. Suppose that $|(S_u \cup u) \cap (S_v \cup v)| = 1$. Then G has corank 3 in M' , so $|G \cap B_3^*| \leq 3$. It now follows from Lemma 3.4 that $G \cap B_3 = \{x_3, y_3\}$, so G is a confining set of M' ; a contradiction. If $|(S_u \cup u) \cap (S_v \cup v)| = 2$, then G is a 4-element corank-2 subset of M' , and it follows from Lemma 3.2 that G is a confining set; a contradiction. So $S_u \cup u = S_v \cup v$, completing the proof of 4.1.3. \square

By 4.1.3 we have that $S_u \cup u = S_v \cup v$. Then, by Lemma 3.4, we may assume that $S_u = \{v, x_2\}$ and $S_v = \{u, x_2\}$. Since b is spanned by $S_u = \{v, x_2\}$, and $A_{y_2b} \neq 0$ because the bad submatrix has no zero entries, it follows that $A_{y_2v} \neq 0$. Hence a pivot on A_{y_2v} is allowable. Now $\{x_0, y_0, a, b\}$ incriminates (M, A^{y_2v}) , where $x_0 = x_2$ and $y_0 = v$. Let $B_0 = B_2 \triangle \{y_2, v\}$. Then u is an (N, B_0) -strong element outside of $\{x_0, y_0\}$, and $\{u, x_0, y_0\}$ is a triad. If y_2 is not (N, B_0) -strong, then the proposition holds. So suppose that y_2 is an (N, B_0) -strong element of M' . By 4.1.1, $M' \setminus y_2$ is 3-connected up to series pairs. By Lemma 3.4, $M' \setminus y_2$ has a series pair S_{y_2} , and $S_{y_2} \cup y_2$ is a triad in M' . By 4.1.3, $S_u \cup u = S_{y_2} \cup y_2$. But $y_2 \notin S_u \cup u$, so this is contradictory. \square

Later, we refer to a basis B_0 satisfying Proposition 4.1 as a *strengthened basis*.

In the remainder of this section we show that if $M' = M \setminus a, b$ has a confining set, then $|E(M)| \leq |E(N)| + 9$. Let B be a basis of M , and let A be a $B \times B^*$ companion \mathbb{P} -matrix of M such that $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$.

We begin with the following constraint on the strong elements of M' .

Lemma 4.2. *If M' has a confining set G relative to the basis B , then M' has no (N, B) -strong elements outside of G .*

Proof. Suppose M' has a confining set G relative to the basis B . If G is a 4-element cosegment, then it follows from Lemma 3.6 that M' has no (N, B) -strong elements outside of G .

Assume now that $G = \{u, v, w, x, y\}$ has corank 3 in M' . By the definition of a confining set, $\{u, v, w\} \subseteq B^*$, and $\{u, v, w\}$ contains an (N, B) -strong element. Suppose t is an (N, B) -strong element outside of G . Then $t \in B^*$ by Lemma 3.1. Now $M' \setminus t$ is 3-connected up to series classes; we next show that $M' \setminus t$ is in fact 3-connected up to series pairs.

Suppose that M' has a cosegment G' containing t with $|G'| \geq 4$. Then $G' = \{s, t, x, y\}$ for some $s \in B^*$ by Lemma 3.2. If $s \notin \{u, v, w\}$, then, as there is an (N, B) -strong element in $\{u, v, w\}$, there is some (N, B) -strong element of M' outside of the 4-element cosegment G' , contradicting Lemma 3.6. Thus we may assume, without loss of generality, that $G' = \{u, t, x, y\}$. Then $G \cup G'$ has corank 3; a contradiction, because $G \cup G'$ has a four-element subset $\{t, u, v, w\}$ contained in B^* . Therefore $M' \setminus t$ is 3-connected up to series pairs.

By Lemma 3.4, $M' \setminus t$ has a series pair S_t that meets $\{x, y\}$. Then $T^* = S_t \cup t$ is a triad of M' ; without loss of generality, we may assume that $S_t = \{x, z\}$ for some $z \in E(M') - \{x, t\}$. If $z \in G$, then $G \cup T^*$ has corank

at most 3 but contains a 4-element subset $\{t, u, v, w\}$ of B^* ; a contradiction. Thus $z \notin G$, so $z \in B^*$ by Lemma 3.4. Then $G \cup T^*$ has corank 4 but contains a 5-element subset $\{t, u, v, w, z\}$ of B^* ; a contradiction. \square

The following results consider allowable pivots when M' has a confining set. The routine proof of the first lemma is omitted. The second is a straightforward consequence of the first, using the fact that $\{x, y\} \subseteq \text{cl}_{M'}^*(G \cap B^*)$.

Lemma 4.3. *Let G be a confining set relative to the basis B , and let $p \in B$. Then $A_{pq} = 0$ for all $q \in (B^* - \{a, b\}) - G$ if and only if $p \in \text{cl}_{M'}^*(G \cap B^*)$.*

Lemma 4.4. *Let G be a confining set relative to the basis B . Then $A_{xq} = 0$ and $A_{yq} = 0$ for all $q \in (B^* - \{a, b\}) - G$.*

Lemma 4.5. *Let G be a confining set relative to the basis B . If $A_{pq} \neq 0$ for some $p \in B - \{x, y\}$ and $q \in (B^* - \{a, b\}) - G$, then a pivot on A_{pq} is allowable. Moreover, G is a confining set relative to the basis $B \Delta \{p, q\}$.*

Proof. Suppose that $A_{pq} \neq 0$ for some $p \in B - \{x, y\}$ and $q \in (B^* - \{a, b\}) - G$. Then the pivot on A_{pq} is allowable by Lemma 4.4. Since $G \cap B = G \cap (B \Delta \{p, q\})$ and $G \cap B^* = G \cap (B^* \Delta \{p, q\})$, G is a confining set relative to $B \Delta \{p, q\}$. \square

Due to the existence of these allowable pivots when M' has a confining set, the following restrictions are imposed on elements of M' .

Lemma 4.6. *Let G be a confining set relative to the basis B . For every $z \in E(M')$,*

- (i) *if z is N -contractible and $\text{si}(M'/z)$ is 3-connected, then $z \in G$; and*
- (ii) *if z is N -deletable and $\text{co}(M' \setminus z)$ is 3-connected, then $z \in \text{cl}_{M'}^*(G)$.*

Proof. Suppose there is an element $z \in E(M') - G$ that is N -contractible, and $\text{si}(M'/z)$ is 3-connected. Since $z \notin G$ it follows from Lemma 4.2 that $z \in B^* - G$. Then $A_{xz} = A_{yz} = 0$ by Lemma 4.4, so there is some $p \in B - \{x, y\}$ such that $A_{pz} \neq 0$ because M' has no loops. Let $B' = B \Delta \{p, z\}$. Now, a pivot on A_{pz} is allowable by Lemma 4.5. So M' has an (N, B') -strong element z in $B' - \{x, y\}$; a contradiction of Lemma 3.1. This proves (i).

Now suppose there is an element $z \in E(M') - \text{cl}^*(G)$ that is N -deletable, and $\text{co}(M' \setminus z)$ is 3-connected. Then $z \notin G$, so $z \in B - \{x, y\}$ by Lemma 4.2. It follows from Lemma 4.3 that there is some $q \in (B^* - \{a, b\}) - G$ such that $A_{zq} \neq 0$. Let $B' = B \Delta \{z, q\}$. Now, a pivot on A_{zq} is allowable by Lemma 4.5, and G is a confining set relative to B' . But z is an (N, B') -strong element outside of G ; a contradiction of Lemma 4.2. This proves (ii). \square

When C and D are disjoint subsets of $E(M')$ such that $M'/C \setminus D \cong N$, we say (C, D) is an N -labelling of M' . For the remainder of the section, suppose M' has a confining set G , and let (C, D) be an N -labelling of M' . Recall that if G has corank three, then there is a (N, B) -strong element $u \in G \cap B^*$. In this case, we choose an N -labelling (C, D) such that $u \in D$. Having fixed (C, D) , our goal is to bound the size of $C \cup D$, and thus bound $|E(M)| - |E(N)|$.

We write $r^*(X)$ instead of $r_{M'}^*(X)$, and $\text{cl}^*(X)$ instead of $\text{cl}_{M'}^*(X)$, for the remainder of the section.

Lemma 4.7. *Suppose that the confining set G has corank 3 in M' . If $z', z'' \in C \cup D$ are in $\text{cl}^*(G) - G$, then for every partition (X, Y) of $G \cup \{z', z''\}$, either $r^*(X) \geq 3$ or $r^*(Y) \geq 3$.*

Proof. Suppose that (X, Y) is a partition of $G \cup \{z', z''\}$ such that $\max\{r^*(X), r^*(Y)\} \leq 2$. We claim that either z' or z'' is an element that contradicts Lemma 4.6(i). Since $|G \cup \{z', z''\}| = 7$, we may assume that $|X| \geq 4$. Then X is a cosegment with at least four elements that contains at least one element $z \in \{z', z''\}$, so $\text{si}(M'/z)$ is 3-connected by the dual of Lemma 2.2. Hence z is not N -contractible by Lemma 4.6(i), so $z \in D$.

First suppose that $z', z'' \in X$. Then $z', z'' \in D$, but z' is in a series class $X \cup z'$ of $M' \setminus z''$, so z' is N -contractible in $M' \setminus z''$ and hence in M' ; a contradiction of Lemma 4.6(i).

We may now assume that $z' \in X$ and $z'' \in Y$, so X and $\text{cl}^*(Y)$ are both 4-element cosegments. Hence both $\text{si}(M'/z')$ and $\text{si}(M'/z'')$ are 3-connected by the dual of Lemma 2.2. By the definition of a confining set, there is some element $u \in G - \{x, y\}$ that is (N, B) -strong in M' , and u belongs to either X or Y . Hence, in $M' \setminus u$, either z' or z'' is in a non-trivial series class, so at least one of z' and z'' is N -contractible in M' ; a contradiction of Lemma 4.6(i). \square

Lemma 4.8. *There are at most two elements of D that belong to $\text{cl}^*(G) - G$.*

Proof. Suppose that there are distinct elements $z, z', z'' \in (\text{cl}^*(G) - G) \cap D$. Then $z, z', z'' \in B - \{x, y\}$, since $G \cap B^*$ is a basis for $\text{cl}^*(G)$.

If G is a 4-element cosegment of M' , then $\text{cl}^*(G)$ is a cosegment containing z and z' . Since z' is N -deletable, z is in a non-trivial series class of $M' \setminus z'$, and $|E(N)| \geq 4$, the element z is N -contractible in M' . By the dual of Lemma 2.2, M'/z is 3-connected, so z' is an (N, B) -strong element of $B - \{x, y\}$; a contradiction of Lemma 3.1.

Now we may assume that $\text{cl}^*(G)$ has corank 3 in M' . We first show that z is N -contractible in M' . If $\{z, z', z''\}$ is a triad of M' , then z is N -contractible since it is in a series pair of $M' \setminus z'$, and z' is N -deletable in M' . So suppose $\{z, z', z''\}$ is coindependent in M' . Then $\{z, z', z''\}$ is a cobasis for $\text{cl}^*(G)$. As $M' \setminus z', z''$ has an N -minor, and $G \cup z$ is contained in a series class in this matroid, it follows that $M' \setminus \{z', z''\}/z$ has an N -minor. In particular, z is N -contractible in M' .

Now z is an N -contractible element of M' , so it follows from Lemma 4.6(i) that $\text{si}(M'/z)$ is not 3-connected. Hence there is a vertical 3-separation (X, z, Y) of M' for some X and Y . But then either X or Y cospans $\text{cl}^*(G)$ by Lemma 4.7. Assume X cospans $\text{cl}^*(G)$. Then $z \in \text{cl}^*(X)$, and by the definition of a vertical 3-separation, $z \in \text{cl}(Y)$; a contradiction to orthogonality. \square

Lemma 4.9. *If $c \in E(M')$ is N -flexible, then $c \in \text{cl}^*(G)$.*

Proof. Suppose that c is N -flexible. By Bixby's Lemma, either $\text{si}(M'/c)$ or $\text{co}(M' \setminus c)$ is 3-connected. Lemma 4.6 then implies that $c \in \text{cl}^*(G)$, as required. \square

Lemma 4.10. *If $z \in E(M')$ is N -deletable, then $z \in \text{cl}^*(G)$.*

Proof. Let $z \in E(M') - \text{cl}^*(G)$, and suppose that z is N -deletable. It then follows from Lemma 4.6(ii) that $\text{co}(M' \setminus z)$ is not 3-connected. Thus, by the dual of Lemma 2.16, there is a cyclic 3-separation (X, z, Y) of M' such that at most one element of X is not N -flexible. We claim that $X \subseteq \text{cl}^*(G)$. The claim follows immediately from Lemma 4.9 unless $s \in X$ is the single element of X that is not N -flexible. By the dual of Lemma 2.16, the element s is N -deletable and $\text{co}(M' \setminus s)$ is 3-connected, so, by Lemma 4.6(ii), $s \in \text{cl}^*(G)$. Thus $X \subseteq \text{cl}^*(G)$, as claimed.

Since (X, z, Y) is a cyclic 3-separation, $r^*(X) \geq 3 \geq r^*(G)$. Thus $\text{cl}^*(X) = \text{cl}^*(G)$. But $z \in \text{cl}^*(X)$ because (X, z, Y) is a cyclic 3-separation in M' , so $z \in \text{cl}^*(G)$; a contradiction. \square

Lemma 4.11. *Suppose that the confining set G is a cosegment. Then $|\text{cl}^*(G) - G \cap (C \cup D)| \leq 1$. In particular, no elements of $\text{cl}^*(G) - G$ are N -contractible.*

Proof. As $\text{cl}^*(G)$ has corank two, M'/p is 3-connected for any $p \in \text{cl}^*(G)$, by Lemma 2.2. Thus, for any $p \in \text{cl}^*(G) - G$, Lemma 4.6(i) implies that p is not N -contractible. Let p and q be distinct elements in $C \cup D$ such that $p, q \in \text{cl}^*(G) - G$. Then $p, q \in D$, but p is in a series class in $M' \setminus q$, so p is N -contractible; a contradiction. \square

Lemma 4.12. *Suppose that the confining set G has corank three, and there is an element $p \in \text{cl}^*(G) - G$ that is N -contractible. Then either*

- (i) $\text{cl}^*(G) - G = \{p\}$, or
- (ii) $|E(M)| \leq |E(N)| + 9$.

Proof. Suppose that (i) does not hold. Then there are distinct elements p and q in $\text{cl}^*(G) - G$, where p is N -contractible. By Lemma 4.6(i), $\text{si}(M'/p)$ is not 3-connected. Let (U, p, V) be a vertical 3-separation of M' such that $|U \cap E(N)| \leq 1$ and $V \cup p$ is closed. If U (or V) spans $\text{cl}^*(G)$, then U (or V , respectively) also spans p , as $p \in \text{cl}^*(G)$. But this contradicts that $p \in \text{cl}(U) \cap \text{cl}(V)$. Thus $r^*(\text{cl}^*(G) \cap U) \leq 2$ and $r^*(\text{cl}^*(G) \cap V) \leq 2$. Recall that G is the union of triads T_1^* and T_2^* . It follows that $\text{cl}^*(G) - p$ is the union of two cosegments $G_1 = \text{cl}^*(T_1^*)$ and $G_2 = \text{cl}^*(T_2^*)$. Without loss of generality, we assume that $q \in G_1$, so $|G_1| \geq 4$. By the dual of Lemma 2.2, M'/q is 3-connected, so q is not N -contractible, by Lemma 4.6(i).

If $|G_2| \geq 4$, then, by Lemma 3.2, $|G_2| = 4$ and $G_2 \cap B = \{x, y\}$. So G_2 is also a confining set. But then $q \notin \text{cl}^*(G_2)$, contradicting Lemma 4.10. So we may assume that $|G_2| = 3$.

If $G_1 - q$ contains an element that is N -deletable, then it follows that q is N -contractible; a contradiction. So no element in $G_1 - q$ is N -deletable; in particular, the (N, B) -strong element $u \in G \cap B^*$ is not in G_1 , and no elements in G_1 are N -flexible. Moreover, if $q \in U$, then q is N -contractible by Lemma 2.16; a contradiction. Letting $G_1 \cap G_2 = \{v\}$, we may now assume that $G_1 - v \subseteq V$, and $G_2 - v \subseteq U$.

By Lemma 2.16, each $y \in U$ is either N -flexible, or y is N -contractible and $\text{si}(M'/y)$ is 3-connected. In the former case, $y \in \text{cl}^*(G)$ by Lemma 4.9; in the latter, $y \in G$ by Lemma 4.6(i). So $U \subseteq \text{cl}^*(G)$. Since $G_1 - v \subseteq V$, and $|U| \geq 3$, it now follows that $U = G_2$, where G_2 is the triad containing $\{u, v\}$. Let $G_2 = \{u, v, w\}$. Note that v is not N -deletable, since $v \in G_1 - q$.

It follows, by Lemma 2.15, that $p \in \text{cl}(U - v)$, so $\{u, w, p\}$ is a triangle of M' . This triangle is co-independent, since M' is 3-connected, so it cospans $\text{cl}^*(G)$. Moreover, the only N -flexible elements of M' are $\{u, w, p\}$.

We now bound the elements of $C \cup D$ outside of $\text{cl}^*(G)$. By Lemma 4.10, every element of $C \cup D$ that is not in $\text{cl}^*(G)$ is in C . Let $z \in C - \text{cl}^*(G)$. Then, by Lemma 4.6(i), $\text{si}(M'/z)$ is not 3-connected, so there is a vertical 3-separation (X, z, Y) such that $|X \cap E(N)| \leq 1$ and $Y \cup z$ is closed. Thus, by Lemma 2.16, at most one element of X is not N -flexible, and if there is such an element s , then s is N -contractible and $\text{si}(M'/s)$ is 3-connected. If $X = \{u, w, p\}$, then $z \in \text{cl}(X) - X$, but as $\{u, w, p\}$ cospans G , we then have $|\text{cl}^*(X) - X| > 1$, which contradicts Lemma 2.6. So X contains an element s , where s is N -contractible and $\text{si}(M'/s)$ is 3-connected, so $s \in G$ by Lemma 4.6(i). Note that q is in the coclosure of the co-independent triangle $\{u, w, p\}$, so $\{u, w, p, q\}$ is 3-separating. By uncrossing $\{u, w, p, q\}$ and X , we observe that the set $P = \{u, w, p, q, s\}$ is also 3-separating. Moreover, since $z \in \text{cl}(X)$, we have $z \in \text{cl}(P)$. Let $Q = E(M') - (P \cup z)$. We may assume that $|Q| \geq 3$, otherwise the lemma holds trivially. So $P \cup z$ is exactly 3-separating. As $v \in \text{cl}^*(P \cup z)$, we have $v \notin \text{cl}(Q - v)$, so $r(Q) \geq 3$. Thus, (P, z, Q) is a vertical 3-separation, where $|P \cap E(N)| \leq 1$. Since $q \in \text{cl}^*(P - q)$, we have $q \notin \text{cl}(Q)$. By Lemma 2.15(i), it follows that q is N -contractible; a contradiction.

We deduce that $C - \text{cl}^*(G) = \emptyset$. So $C \cup D \subseteq \text{cl}^*(G)$. As $|(C \cup D) \cap (\text{cl}^*(G) - G)| = 2$, we have $|C \cup D| \leq 7$. Thus, $|E(M)| - |E(N)| = |C \cup D| + |\{a, b\}| \leq 9$, as required. \square

By Lemma 4.10, $D - \text{cl}^*(G) = \emptyset$. We now focus on bounding $|C - \text{cl}^*(G)|$.

Lemma 4.13. *Suppose that there exist distinct $p_1, p_2 \in E(M') - \text{cl}^*(G)$ such that M'/p_i has an N -minor for $i \in \{1, 2\}$. Let (X_1, p_1, Y_1) and (X_2, p_2, Y_2) be vertical 3-separations of M' . Then $|X_1 \cap X_2| \leq 1$ or $|Y_1 \cap Y_2| \leq 1$.*

Proof. Towards a contradiction, suppose that $|X_1 \cap X_2| \geq 2$ and $|Y_1 \cap Y_2| \geq 2$. By uncrossing, the sets $X_1 \cup X_2$, $X_1 \cup X_2 \cup p_1$, $X_1 \cup X_2 \cup p_2$, and $X_1 \cup X_2 \cup \{p_1, p_2\}$ are all 3-separating. Since $|Y_1 \cap Y_2| \geq 2$, the sets $X_1 \cup X_2$, $X_1 \cup X_2 \cup p_1$, $X_1 \cup X_2 \cup p_2$, and $X_1 \cup X_2 \cup \{p_1, p_2\}$ are sides of exact 3-separations of M' and p_1, p_2 are guts elements. In particular, $(X_1 \cup X_2 \cup p_2, p_1, Y_1 \cap Y_2)$ is a vertical 3-separation of M' unless $r(Y_1 \cap Y_2) \leq 2$. But if $r(Y_1 \cap Y_2) \leq 2$, then $(Y_1 \cap Y_2) \cup \{p_1, p_2\}$ is a segment of M' with at least four elements, so p_1 belongs to a non-trivial parallel class of M'/p_2 . Then p_1 is N -deletable in M'/p_2 and hence in M' , so $p_1 \in \text{cl}^*(G)$ by Lemma 4.9; a contradiction. Thus $(X_1 \cup X_2 \cup p_2, p_1, Y_1 \cap Y_2)$ is a vertical 3-separation of M' , and either $|(X_1 \cup X_2 \cup p_2) \cap E(N)| \leq 1$ or $|(Y_1 \cap Y_2) \cap E(N)| \leq 1$.

If $|(X_1 \cup X_2 \cup p_2) \cap E(N)| \leq 1$, then there is an element p_2 in the non- N -side of $(X_1 \cup X_2 \cup p_2, p_1, Y_1 \cap Y_2)$. Since $p_2 \in \text{cl}(Y_1 \cap Y_2)$, it follows from Lemma 2.15(ii) that p_2 is N -deletable. Hence $p_2 \in \text{cl}^*(G)$ by Lemma 4.10; a contradiction. So $|(Y_1 \cap Y_2) \cap E(N)| \leq 1$. But $(X_1 \cup X_2, p_1, (Y_1 \cap Y_2) \cup p_2)$ is also a vertical 3-separation of M' . Moreover, as $|E(N)| \geq 4$, we have $|(X_1 \cup X_2 \cup p_2) \cap E(N)| \geq 3$, so $|(X_1 \cup X_2) \cap E(N)| \geq 2$ and hence, by Lemma 2.14, $|(Y_1 \cap Y_2) \cup p_2| \leq 1$. Again, it follows that $p_2 \in \text{cl}^*(G)$; a contradiction. \square

Lemma 4.14. *Suppose $\text{cl}^*(G)$ has at most six N -contractible elements. Then $|C - \text{cl}^*(G)| \leq 2$.*

Proof. Suppose that $|C - \text{cl}^*(G)| \geq 3$. Let p_1, p_2, p_3 be distinct elements in $C - \text{cl}^*(G)$. It follows from Lemma 4.6(i) that $\text{si}(M'/p_i)$ is not 3-connected, so there is a vertical 3-separation (X_i, p_i, Y_i) of M' , for each $i \in \{1, 2, 3\}$, where $|X_i \cap E(N)| \leq 1$ and $Y_i \cup p_i$ is closed. Then, by Lemma 2.16, each element $x \in X_i$ is either N -flexible, or x is N -contractible and $\text{si}(M'/x)$ is 3-connected. By Lemma 4.9, in the former case, and Lemma 4.6(i), in the latter, $X_i \subseteq \text{cl}^*(G)$. Note that $|X_i| \geq 3$, for each i , and if $|X_i| = 3$, then X_i is a triad.

Let H be the set of N -contractible elements of $\text{cl}^*(G)$. Since, for $i \in \{1, 2, 3\}$, each element in X_i is N -contractible, $X_i \subseteq H$, where $|H| \leq 6$. We claim that $|X_i \cap X_j| \geq 2$ for some distinct $i, j \in \{1, 2, 3\}$. If, for some $\{i, j, k\} = \{1, 2, 3\}$, the sets X_i and X_j are disjoint, then $X_i \cup X_j = H$, so X_k intersects X_i or X_j in two elements, as claimed. Similarly, if $|X_i| \geq 4$, then either $|X_i \cap X_j| \geq 2$, or $X_i \cup X_j = H$, in which case X_k intersects X_i or X_j in two elements. So we may assume that $|X_i| = 3$ for each $i \in \{1, 2, 3\}$, and the pairwise intersection between any two of the three sets has size one. Let $X_2 = \{x_1, x_2, x_3\}$ where $X_1 \cap X_2 = \{x_1\}$ and $X_2 \cap X_3 = \{x_3\}$. Now $X_2 \cup p_2$ contains a circuit, since (X_2, p_2, Y_2) is a vertical 3-separation. By orthogonality, this circuit does not meet the triad X_3 , nor the triad X_1 , so $X_2 \cup p_2$ contains a circuit of size at most two; a contradiction. This proves the claim. Without loss of generality, we may now assume that $|X_1 \cap X_2| \geq 2$.

If $|E(M') - \text{cl}^*(G)| \geq 4$, then $|Y_1 \cap Y_2| \geq |E(M') - (\text{cl}^*(G) \cup \{p_1, p_2\})| \geq 2$, which contradicts Lemma 4.13. So $|E(M') - \text{cl}^*(G)| \leq 3$, in which case $E(M') - \text{cl}^*(G) = \{p_1, p_2, p_3\}$. Now, as $p_1 \notin \text{cl}^*(G)$, we have $p_1 \in \text{cl}(\{p_2, p_3\})$, by orthogonality. Since M' is 3-connected, $\{p_1, p_2, p_3\}$ is a triangle. Then $\{p_2, p_3\}$ is a parallel pair in M'/p_1 , so the element p_2 is N -deletable. As $\text{si}(M/p_2)$ is not 3-connected, $\text{co}(M \setminus p_2)$ is 3-connected by Bixby's Lemma. But $p_2 \notin \text{cl}^*(G)$; contradicting Lemma 4.6(ii). We deduce that $|C - \text{cl}^*(G)| \leq 2$, thus completing the lemma. \square

We handle one more special case.

Lemma 4.15. *Suppose that the confining set G has corank three, $\text{cl}^*(G) - G = \{q\}$ for some $q \in C \cup D$, and $|C - \text{cl}^*(G)| = 2$. Then $|E(M)| \leq |E(N)| + 9$.*

Proof. Let p_1 and p_2 be distinct elements in $C - \text{cl}^*(G)$. By Lemma 4.6(i), $\text{si}(M'/p_i)$ is not 3-connected, so there is a vertical 3-separation (X_i, p_i, Y_i) of M' for $i \in \{1, 2\}$, where $|X_i \cap E(N)| \leq 1$ and $Y_i \cup p_i$ is closed. By Lemma 2.16, each element $x \in X_i$ is either N -flexible, or x is N -contractible and $\text{si}(M'/x)$ is 3-connected. By Lemma 4.9, in the former case, and Lemma 4.6(i), in the latter, $X_i \subseteq \text{cl}^*(G)$. Note that $|X_i| \geq 3$, for each i , and if $|X_i| = 3$, then X_i is a triad. Recall also that G is the union of two triads T_1^* and T_2^* .

Suppose q together with one of the triads, T_1^* say, forms a cosegment. Then $\text{si}(M/q)$ is 3-connected, by the dual of Lemma 2.2, so q is not N -contractible, by Lemma 4.6(i). Now $X_1 \cup X_2 \subseteq G$, so it follows that $\{X_1, X_2\} = \{T_1^*, T_2^*\}$. But then $p_1 \in \text{cl}(T_1^*)$, up to swapping the labels on p_1 and p_2 , so, by orthogonality with T_2^* , we deduce that $\{p_1, s_1, t_1\}$ is a

triangle, where $T_1^* - T_2^* = \{s_1, t_1\}$. Let $T_1^* \cap T_2^* = \{v\}$. Now, as $T_1^* \cup q$ is a cosegment, $\{t_1, v, q\}$ is a triad that intersects the triangle $\{p_1, s_1, t_1\}$ in a single element; a contradiction.

Now suppose $q \notin \text{cl}^*(T_1^*) \cup \text{cl}^*(T_2^*)$. We claim that $|X_1 \cap X_2| \geq 2$. Suppose not. Then $|X_i| = 3$, for some $i \in \{1, 2\}$, so we may assume X_1 , say, is a triad. Either $X_1 \in \{T_1^*, T_2^*\}$, or $q \in X_1$ and X_1 intersects T_1^* and T_2^* in one element each. Let $T_1^* = \{v, s_1, t_1\}$ and $T_2^* = \{v, s_2, t_2\}$.

If $X_1 = T_1^*$, then, as $p_1 \in \text{cl}(X_1)$, by orthogonality with T_2^* we have that $\{p_1, s_1, t_1\}$ is a triangle. But as $\{s_2, t_2, q\}$ cospans $\text{cl}^*(G)$, the element t_1 is in a cocircuit contained in $\{s_2, t_2, q, t_1\}$, which contradicts orthogonality with the triangle $\{p_1, s_1, t_1\}$.

On the other hand, if X_1 is a triad that meets both $\{s_1, t_1\}$ and $\{s_2, t_2\}$, and $q \in X_1$, then, p_1 is in a circuit contained in $X_1 \cup p_1$. But $X_1 \cup p_1$ meets T_1^* and T_2^* in a single element each, so, by orthogonality, p_1 is in a parallel pair; a contradiction. So $|X_1 \cap X_2| \geq 2$ as claimed. Note that the lemma holds trivially if $|E(M)| \leq 11$, since $|E(N)| \geq 4$. So we may assume that $|E(M')| \geq 10$, in which case $|Y_1 \cap Y_2| \geq 2$. But this contradicts Lemma 4.13. \square

Finally, we are in a position to prove the main result of this section.

Proposition 4.16. *Suppose that M' has a confining set. Then $|E(M)| \leq |E(N)| + 9$.*

Proof. First, suppose that G is a cosegment. Then, by Lemma 4.11, $\text{cl}^*(G) - G$ has at most one element of $C \cup D$, and $\text{cl}^*(G)$ consists of at most four N -contractible elements (those elements in G). Therefore, by Lemma 4.14, $|C - \text{cl}^*(G)| \leq 2$. As $D \subseteq \text{cl}^*(G)$, by Lemma 4.10, we have

$$\begin{aligned} |E(M)| - |E(N)| &\leq |\text{cl}^*(G) \cap (C \cup D)| + |C - \text{cl}^*(G)| + |\{a, b\}| \\ &\leq 5 + 2 + 2 = 9. \end{aligned}$$

Now suppose that G has corank three. Consider first the case where $|\text{cl}^*(G) - G| \geq 2$. If $\text{cl}^*(G) - G$ contains an element that is N -contractible, then, by Lemma 4.12, $|E(M)| \leq |E(N)| + 9$, as required. So we may assume that no elements in $\text{cl}^*(G) - G$ are N -contractible. In particular, $\text{cl}^*(G)$ contains at most five N -contractible elements. Now, by Lemma 4.14, $|C - \text{cl}^*(G)| \leq 2$. Suppose there is an element $q \in D \cap (\text{cl}^*(G) - G)$, and let p be an element in $\text{cl}^*(G) - G$, with $q \neq p$. Recall that (C, D) was chosen such that $u \in D$, and note that $r_{M' \setminus u}^*(\text{cl}^*(G) - u) = 2$. Thus p is in a series class in $M' \setminus u \setminus q$, so p is N -contractible; a contradiction. It now follows that $|\text{cl}^*(G) \cap (C \cup D)| \leq 5$, and hence $|E(M)| - |E(N)| \leq 5 + 2 + |\{a, b\}| = 9$, as required.

Now consider the case where $|\text{cl}^*(G) - G| \leq 1$. Since $|\text{cl}^*(G)| \leq 6$, Lemma 4.14 implies that $|C - \text{cl}^*(G)| \leq 2$. If $(\text{cl}^*(G) - G) \cap (C \cup D) = \emptyset$, then $|E(M)| - |E(N)| \leq 5 + 2 + |\{a, b\}| = 9$ as required. So suppose that $\text{cl}^*(G) - G = \{q\}$ for some $q \in C \cup D$. We may also assume that $|C - \text{cl}^*(G)| \geq 2$, otherwise the result holds trivially. Now, by Lemma 4.15, $|E(M)| \leq |E(N)| + 9$, as required. \square

5. ROBUST ELEMENTS

Let M be a 3-connected matroid, let N be a 3-connected minor of M such that $|E(N)| \geq 4$, and let B be a basis of M . In this section, we consider the structure of M that arises from elements that are (N, B) -robust but not (N, B) -strong. Recall that a *path of 3-separations* of M is a partition (P_1, P_2, \dots, P_n) of $E(M)$ such that $(P_1 \cup \dots \cup P_i, P_{i+1} \cup \dots \cup P_n)$ is a 3-separation of M for each $i \in \{1, 2, \dots, n-1\}$. The main result of this section shows that the presence of an element that is (N, B) -robust but not (N, B) -strong gives rise to a particular path of 3-separations.

Let (X, z, Y) be a vertical 3-separation of M . We say that X is *z -closed* if $X = \text{cl}^*(X)$ and $X = \text{cl}(X) - z$. We use z -closure to ensure that the (N, B) -strong elements of M are contained in the non- N -side of a vertical 3-separation of M . A set is *fully closed* if it is both closed and coclosed. Given a subset A of $E(M)$, we use $\text{fcl}_M(A)$ to denote the smallest fully closed set that contains A . Thus, the set X is z -closed if $\text{fcl}_{M/z}(X) = X$.

Dually, given a cyclic 3-separation (X, z, Y) , we say X is *z -coclosed* if X is z -closed in M^* .

Lemma 5.1. *If $z \in B$ and z is (N, B) -robust but not (N, B) -strong, then there is some vertical 3-separation (X, z, Y) of M such that X is z -closed and $|X \cap E(N)| \leq 1$.*

Proof. By Lemma 2.7, M has a vertical 3-separation (X, z, Y) , and we may assume that $|X \cap E(N)| \leq 1$, by Lemma 2.14. The elements of $\text{fcl}_{M/z}(X) - X$ can be ordered (x_1, \dots, x_m) such that $X \cup \{x_1, \dots, x_i\}$ is 2-separating in M/z for all $i \in \{1, \dots, m\}$. Let $X_i = X \cup \{x_1, \dots, x_i\}$ and $Y_i = Y - \{x_1, \dots, x_i\}$ for each $i \in \{1, 2, \dots, m\}$. We also let $(X_0, Y_0) = (X, Y)$. Suppose that $|X_j \cap E(N)| \geq 2$ for some $j \in \{1, 2, \dots, m\}$. We shall assume that j is the smallest index such that $|X_j \cap E(N)| \geq 2$. Then $|X_{j-1} \cap E(N)| \leq 1$, so $|Y_{j-1} \cap E(N)| \geq 3$ because $|E(N)| \geq 4$. Hence $|Y_j \cap E(N)| \geq 2$. But then (X_j, Y_j) is a 2-separation of M/z such that $|Y_j \cap E(N)| \geq 2$ and $|X_j \cap E(N)| \geq 2$; a contradiction. Hence $|X_i \cap E(N)| \leq 1$ for all $i \in \{1, \dots, m\}$. Thus, for each i , the partition (X_i, Y_i) is a 2-separation in M/z such that Y_i is the N -side. It follows that $|Y_i| \geq 3$ for all i . In particular, (X_m, Y_m) is a 2-separation of M/z such that X_m is fully closed. Since M is 3-connected, $z \in \text{cl}_M(X_m) \cap \text{cl}_M(Y_m)$. Finally, Y_m is not a parallel class of M/z because X_m is fully closed, so $r_M(Y_m) \geq 3$. Thus (X_m, z, Y_m) is a z -closed vertical 3-separation of M , as desired. \square

Suppose that F is a 4-element fan of M with ordering (f_1, f_2, f_3, f_4) where $\{f_1, f_2, f_3\}$ is a triangle. We say that (f_1, f_2, f_3, f_4) is a *type-I fan relative to B* if $F \cap B = \{f_1, f_3\}$, and (f_1, f_2, f_3, f_4) is a *type-II fan relative to B* if $F \cap B = \{f_1, f_3, f_4\}$. When there is no ambiguity, we also say, in these cases, that F is a type-I or type-II fan relative to B .

We need the following, which is one of the main results of [3].

Lemma 5.2 ([3, Lemma 4.8]). *Suppose that $z \in B$ is an element that is (N, B) -robust but not (N, B) -strong, and let (X, z, Y) be a vertical 3-separation of M such that $|X \cap E(N)| \leq 1$. Then one of the following holds:*

- (i) *there are distinct (N, B) -strong elements $s_1, s_2 \in X$; or*

- (ii) *there are distinct (N, B) -strong elements $s_1 \in X$ and $s_2 \in \text{cl}^*(X) \cap B$;
or*
- (iii) *there are distinct (N, B) -strong elements $s_1 \in X$ and $s_2, s_3 \in \text{cl}(X) \cap B^*$; or*
- (iv) *M has a type-I or type-II fan relative to B contained in $X \cup z$.*

The next lemma is a consequence of Lemmas 2.15 and 5.2.

Lemma 5.3. *Let $z \in B$ be an element that is (N, B) -robust but not (N, B) -strong, and let (X, z, Y) be a vertical 3-separation of M such that X is z -closed and $|X \cap E(N)| \leq 1$. If there is at most one (N, B) -strong element of M contained in X , then there is a type-I or type-II fan $(\alpha, \beta, \gamma, \delta)$ relative to B that is contained in $X \cup z$ where β, γ, δ are N -contractible, and α, β, γ are N -deletable.*

Proof. Since X is z -closed, it follows from Lemma 5.2 that M has a type-I or type-II fan $(\alpha, \beta, \gamma, \delta)$ relative to B such that $\{\alpha, \beta, \gamma, \delta\} \subseteq X \cup z$. Let T^* be the triad $\{\beta, \gamma, \delta\}$. Note that $z \notin T^*$, since $z \in \text{cl}(Y)$. Since $T^* \subseteq X$, it follows from orthogonality that $\beta, \gamma, \delta \notin \text{cl}_M(Y)$. Hence β, γ, δ are N -contractible by Lemma 2.15. It follows, since $\{\alpha, \beta, \gamma\}$ is a triangle of M and $|E(N)| \geq 4$, that α, β, γ are also N -deletable in M . \square

We will also require the following lemma, which can be proved by making routine modifications to [24, Lemma 5.4] or [3, Lemma 6.3].

Lemma 5.4. *Let M be a 3-connected matroid and let (A, Z, B) a partition of $E(M)$ with $|A|, |B| \geq 2$. If, for all $z \in Z$, there is a path of 3-separations (A_z, z, B_z) such that $A \subseteq A_z$ and $B \subseteq B_z$, then there is an ordering (z_1, \dots, z_n) of the elements of Z such that (A, z_1, \dots, z_n, B) is a path of 3-separations of M .*

For the remainder of this section, we work under the following assumptions. Let \mathbb{P} be a partial field, let N be a non-binary 3-connected strong \mathbb{P} -stabilizer for the class of \mathbb{P} -representable matroids, and let M be an excluded minor for the class of \mathbb{P} -representable matroids. Suppose that M has a pair of elements $\{a, b\}$ such that $M \setminus a, b$ is 3-connected with an N -minor, and let $M' = M \setminus a, b$. Let A be a $B \times B^*$ companion \mathbb{P} -matrix of M such that $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$. We assume that M' has no confining set. We also assume that B is chosen such that either there is one (N, B) -strong element u of M' outside of $\{x, y\}$, and $\{u, x, y\}$ is a triad; or there are no (N, B) -strong elements outside of $\{x, y\}$, and for any $B_1 \times B_1^*$ companion \mathbb{P} -matrix A_1 where $\{x_1, y_1, a, b\}$ incriminates (M, A_1) , with $\{x_1, y_1\} \subseteq B_1$ and $\{a, b\} \subseteq B_1^*$, the matroid $M \setminus a, b$ has no (N, B_1) -strong elements outside of $\{x_1, y_1\}$. Note that such a B exists by Proposition 4.1. Recall that we say that such a basis B is *strengthened*.

Let $S \subseteq E(M')$ be a set containing $\{x, y\}$ and any (N, B) -strong elements of M' , where either $|S| = 2$ or S is a triad. In particular, observe that $S \subseteq \text{cl}_{M'}^*(\{x, y\})$.

For the remainder of the section, all ranks, coranks, closure operators, and coclosure operators are with respect to M' .

If z is an element that is (N, B) -robust but not (N, B) -strong in M' , then there is a vertical (or cyclic) 3-separation (X, z, Y) of M' . We now

prove that if the non- N -side of this vertical 3-separation is z -closed (or z -coclosed, respectively), then it contains S . We first handle the case where $z \in B - \{x, y\}$.

Lemma 5.5. *Let $z \in B - \{x, y\}$ be an element that is (N, B) -robust but not (N, B) -strong in M' , and let (X, z, Y) be a vertical 3-separation of M' such that X is z -closed and $|X \cap E(N)| \leq 1$. Then $S \subseteq X$.*

Proof. Suppose that there are at least two distinct (N, B) -strong elements in X . By definition, the (N, B) -strong elements of M contained in X belong to S . If $|S| = 2$, then it follows immediately that $S \subseteq X$. If $|S| = 3$, then S is a triad, so $S \subseteq X$ because X is coclosed.

We may therefore assume that there is at most one (N, B) -strong element of M' contained in X . Then it follows from Lemma 5.3 that there is a type-I or type-II fan $(\alpha, \beta, \gamma, \delta)$ relative to B contained in $X \cup z$ where β, γ, δ are N -contractible and α, β, γ are N -deletable. Let $F = \{\alpha, \beta, \gamma, \delta\}$.

5.5.1. $\{x, y\} \cap \{\alpha, \gamma\} \neq \emptyset$.

Subproof. Assume that $\{x, y\} \cap \{\alpha, \gamma\} = \emptyset$. Suppose β is an (N, B) -strong element of M' . Then, since $\beta \notin B$, it follows that $S = \{\beta, x, y\}$ is a triad of M' . Since $\{\alpha, \beta, \gamma\}$ is a triangle that meets $\{\beta, x, y\}$, it follows from orthogonality that x or y is in $\{\alpha, \gamma\}$; a contradiction because $\{x, y\} \cap \{\alpha, \gamma\} = \emptyset$. Thus β is not an (N, B) -strong element of M' . Since β is N -flexible, $\text{co}(M' \setminus \beta)$ is not 3-connected. Thus, by Bixby's Lemma, $\text{si}(M' / \beta)$ is 3-connected. Since $\{\alpha, \beta, \gamma\}$ is a triangle of M' the cobasis element β is spanned by the basis elements α and γ , so $A_{i\beta} \neq 0$ if and only if $i \in \{\alpha, \gamma\}$. In particular, since $\{x, y\} \cap \{\alpha, \gamma\} = \emptyset$, this means that $A_{\alpha\beta} \neq 0$ and $A_{x\beta} = A_{y\beta} = 0$. Thus a pivot on $A_{\alpha\beta}$ is an allowable pivot. But then β is an $(N, B \Delta \{\alpha, \beta\})$ -strong element outside of $\{x, y\}$ such that $\beta \in B \Delta \{\alpha, \beta\}$; a contradiction of Lemma 3.1. \square

Now α or γ is a member of $\{x, y\}$. Suppose $\delta \in B$, in which case $(\alpha, \beta, \gamma, \delta)$ is a type-II fan. Since δ is N -contractible and $\text{si}(M / \delta)$ is 3-connected by Lemma 2.11, it follows from Lemma 3.1 that $\delta \in \{x, y\}$. Hence $\{x, y\} \subseteq F - z$, and $S \subseteq \text{cl}^*(F - z) \subseteq \text{cl}^*(X) = X$ as required. Thus we may assume that $\delta \in B^*$, in which case F is a type-I fan.

We first handle the case when $\alpha \in \{x, y\}$.

5.5.2. If $\alpha \in \{x, y\}$, then $S \subseteq X$.

Subproof. Assume that $\alpha = x$. If β is an (N, B) -strong element of M' , then $\{\beta, x, y\}$ is a triad of M' , so $S \subseteq \text{cl}^*(\{\beta, x\}) \subseteq \text{cl}^*(F - z) \subseteq X$, as required. So suppose that β is not an (N, B) -strong element of M' . Consider the entry $A_{\alpha\beta}$. Since $\{\alpha, \beta, \gamma\}$ is a triangle of M' it follows that $A_{\alpha\beta} \neq 0$, so a pivot on $A_{\alpha\beta}$ is an allowable pivot. Then $B' = B \Delta \{\alpha, \beta\}$ is a basis of M' , the set $\{\beta, y, a, b\}$ incriminates $(M, A^{\alpha\beta})$, and α is an (N, B') -strong element outside of $\{\beta, y\}$. Since B is strengthened, there is some element $u \in B^*$ such that u is (N, B) -strong and $\{u, x, y\}$ is a triad. Since β and δ are not (N, B) -strong, it follows that $u \in E(M') - F$. But then, by orthogonality between the triad $\{u, \alpha, y\}$ and the triangle $\{\alpha, \beta, \gamma\}$, we have $y \in \{\beta, \gamma\}$, so $y = \gamma$. Therefore $S \subseteq \text{cl}^*(\{x, y\}) \subseteq \text{cl}^*(F - z) \subseteq X$. \square

We may now assume that $\alpha \notin \{x, y\}$, so $\gamma \in \{x, y\}$. Suppose that $\gamma = x$. If β is (N, B) -strong, then $\{\beta, x, y\}$ is a triad, and $\{\beta, \delta, x, y\}$ is a 4-element cosegment, contradicting that M' has no confining set. We deduce that β is not (N, B) -strong.

Suppose that $\text{co}(M' \setminus x)$ is 3-connected. Since $\{\alpha, \beta, x\}$ is a triangle of M' , we have $A_{x\beta} \neq 0$, so a pivot on $A_{x\beta}$ is allowable. Then $B' = B \triangle \{x, \beta\}$ is a basis such that x is an (N, B) -strong element outside of $\{\beta, y\}$, where $\{\beta, y, a, b\}$ incriminates $(M, A^{x\beta})$. Since B is strengthened, there is some (N, B) -strong element $u \in B^*$ such that $\{u, x, y\}$ is a triad. Since β is not (N, B) -strong, u is not in the triangle $\{\alpha, \beta, x\}$. It then follows from orthogonality that $\alpha = y$; a contradiction. So $\text{co}(M' \setminus x)$ is not 3-connected, and thus $\text{si}(M' / x)$ is 3-connected by Bixby's Lemma.

Since β is not (N, B) -strong, there is a cyclic 3-separation (P, β, Q) of M' . By orthogonality, we may assume that $x \in P$ and $\alpha \in Q$. Consider $(P - x, x, Q \cup \beta)$. Observe that $Q \cup \beta$ and $Q \cup \{\beta, x\}$ are exactly 3-separating, since $x \in \text{cl}(Q \cup \beta)$. But $(P - x, x, Q \cup \beta)$ is not a vertical 3-separation of M' , since $\text{si}(M' / x)$ is 3-connected. Thus $r(P - x) \leq 2$, so P contains a triangle. By orthogonality, P is a triangle and $P = \{x, \delta, \mu\}$ for some $\mu \in E(M')$. Thus M' has a 5-element fan with ordering $(\alpha, \beta, x, \delta, \mu)$. Moreover, $\mu \in \text{cl}(Q)$ or else $\{\beta, x, \delta, \mu\}$ is a 4-element cosegment; a contradiction to orthogonality. Now $\text{co}(M' \setminus \mu)$ is 3-connected by Lemma 2.11, and μ is N -deletable since μ is in a non-trivial parallel class in M' / γ . Suppose $\mu \in B^* - \{a, b\}$. Then μ is (N, B) -strong and outside of $\{x, y\}$, so $\{\mu, x, y\}$ is a triad. By orthogonality, it follows that $\alpha = y$, contradicting the assumption that $\alpha \notin \{x, y\}$. We deduce that $\mu \in B$.

We now repeat this argument, interchanging the roles of x and β . Since $\text{co}(M' \setminus x)$ is not 3-connected, there is a cyclic 3-separation (P', x, Q') of M' . By orthogonality, we may assume that $\beta \in P'$ and $\alpha \in Q'$. Consider $(P' - \beta, \beta, Q' \cup x)$. Observe that $Q' \cup x$ and $Q' \cup \{\beta, x\}$ are exactly 3-separating, since $\beta \in \text{cl}(Q' \cup x)$. But $(P' - \beta, \beta, Q' \cup x)$ is not a vertical 3-separation of M' , since $\text{si}(M' / \beta)$ is 3-connected, by Bixby's Lemma. Thus $r(P' - \beta) = 2$, and it follows by orthogonality that P' is a triangle of M' . By orthogonality between P' and $\{\beta, x, \delta\}$, we have $\delta \in P'$. Since $\beta \notin \text{cl}(P')$, it follows that $\mu \in Q'$. Let $P' = \{\beta, \delta, \varepsilon\}$ for some $\varepsilon \in E(M')$. Now, $\varepsilon \in \text{cl}(Q')$ or else $\{\varepsilon, x, \beta, \delta\}$ is a 4-element cosegment; a contradiction to orthogonality.

Now α, μ, ε are in the closure of the triad $\{\beta, x, \delta\}$, so $\{\alpha, \mu, \varepsilon\}$ is a triangle. But $\alpha, \mu \in B$, so $\varepsilon \in B^*$. We claim that ε is an (N, B) -strong element of M' . That ε is (N, B) -robust follows from the fact that β is N -contractible and $\{\delta, \varepsilon\}$ is a parallel pair in M' / β . Since $(F, \varepsilon, E(M') - F)$ is a vertical 3-separation of M' , Lemma 2.7 and Bixby's Lemma imply that $\text{co}(M' \setminus \varepsilon)$ is 3-connected. As ε is an (N, B) -strong element of M' outside of $\{x, y\}$, we have that $\{\varepsilon, x, y\}$ is a triad of M' . But $\{\varepsilon, x, y\}$ intersects the triangle $\{\beta, \delta, \varepsilon\}$ in a single element; a contradiction to orthogonality. \square

Next we handle the case where the element z , which is (N, B) -robust but not (N, B) -strong, is in B^* .

Lemma 5.6. *Let $z \in B^*$ be an element of M' that is (N, B) -robust but not (N, B) -strong, and let (X, z, Y) be a cyclic 3-separation of M' such that X is z -coclosed and $|X \cap E(N)| \leq 1$. Then $S \subseteq X$.*

Proof. Suppose that there are at least two distinct (N, B) -strong elements in X . The (N, B) -strong elements of M' contained in X must belong to S by the definition of S . If $|S| = 2$, then it follows immediately that $S \subseteq X$. If $|S| = 3$, then S is a triad, so $S \subseteq \text{cl}^*(X) = X \cup z$, as X is z -coclosed, but $z \notin S$, so $S \subseteq X$ as required.

We may therefore assume that there is at most one (N, B) -strong element of M' contained in X . Then it follows from the dual of Lemma 5.3 that there is a type-I or type-II fan $(\alpha, \beta, \gamma, \delta)$ relative to B^* in $(M')^*$ that is contained in $X \cup z$ where β, γ, δ are N -deletable and α, β, γ are N -contractible in M' .

Suppose that F is a type-II fan relative to B^* in $(M')^*$. Then δ is an (N, B) -strong element of M' by Lemma 2.11. Hence M' has a triad $\{\delta, x, y\}$. By orthogonality with the triangle $\{\beta, \gamma, \delta\}$, we have $\{\beta, \gamma\} \cap \{x, y\} \neq \emptyset$; but $\gamma \notin B$, so $\beta \in \{x, y\}$. Since $\{\beta, \delta\} \subseteq F - z \subseteq X$, we have $S \subseteq \text{cl}^*(\{\beta, \delta\}) \subseteq \text{cl}^*(X) = X \cup z$, as X is z -coclosed. But $z \notin S$, so $S \subseteq X$ as required.

We may now assume that F is a type-I fan relative to B^* in $(M')^*$. If γ is an (N, B) -strong element of M' , then M' has a triad $\{\gamma, x, y\}$. By orthogonality with $\{\beta, \gamma, \delta\}$, either $\beta \in \{x, y\}$ or $\delta \in \{x, y\}$. As $z \notin \{\beta, \delta\}$, we have $S \subseteq \text{cl}^*(F - z) \subseteq X$ because X is z -coclosed and $z \notin S$. Therefore we may also assume that $\text{co}(M' \setminus \gamma)$ is not 3-connected, so $\text{si}(M'/\gamma)$ is 3-connected, by Bixby's Lemma.

5.6.1. $\{x, y\} \cap \{\beta, \delta\} \neq \emptyset$.

Subproof. Suppose that $\{x, y\} \cap \{\beta, \delta\} = \emptyset$. Then, since $\{\beta, \gamma, \delta\}$ is a triangle of M' , it follows that $A_{x\gamma} = A_{y\gamma} = 0$ and $A_{\beta\gamma} \neq 0$. Hence a pivot on $A_{\beta\gamma}$ is allowable, and γ is in the basis $B' = B \Delta \{\beta, \gamma\}$ of M' , where $\{x, y, a, b\}$ incriminates $(M, A^{\beta\gamma})$. But then γ is an (N, B') -strong element of M' in $B' - \{x, y\}$; a contradiction of Lemma 3.1. \square

Suppose $\delta \in \{x, y\}$. Then, since $\{\beta, \gamma, \delta\}$ is a triangle of M' , $A_{\delta\gamma} \neq 0$, and a pivot on $A_{\delta\gamma}$ is allowable. Hence M' has a basis $B' = B \Delta \{\delta, \gamma\}$ with an (N, B') -strong element δ in $(B')^*$. Since B is a strengthened basis, there is an (N, B) -strong element $u \in B^*$ such that $S = \{u, x, y\}$ is a triad of M' . By orthogonality, either $\beta \in S$ or $\gamma \in S$. Hence $S \subseteq \text{cl}^*(F - z) \subseteq X$ because X is z -coclosed and $z \notin S$. A similar argument holds if $\beta \in \{x, y\}$ and $\text{co}(M' \setminus \beta)$ is 3-connected.

We may now assume that $\beta \in \{x, y\}$ and that $\text{co}(M' \setminus \beta)$ is not 3-connected. Let (P, β, Q) be a cyclic 3-separation of M' . Since β is in a triangle of M' , we may assume that $\gamma \in P$ and $\delta \in Q$. Consider $(P - \gamma, \gamma, Q \cup \beta)$. Observe that $Q \cup \beta$ and $Q \cup \{\beta, \gamma\}$ are exactly 3-separating, the latter since $\gamma \in \text{cl}(Q \cup \beta)$. But $(P - \gamma, \gamma, Q \cup \beta)$ is not a vertical 3-separation of M' , since $\text{si}(M'/\gamma)$ is 3-connected, so it follows that $r(P) = 2$. By orthogonality with the triad $\{\alpha, \beta, \gamma\}$, it follows that $\alpha \in P$ and P is a triangle of M' . Thus $P = \{\alpha, \gamma, p\}$ for some $p \in E(M') - F$.

Let (P', γ, Q') be a cyclic 3-separation of M' . Since $\{\beta, \gamma, \delta\}$ is a triangle of M' , we may assume that $\beta \in P'$ and $\delta \in Q'$. Now $Q' \cup \gamma$ and $Q' \cup \{\beta, \gamma\}$ are exactly 3-separating, but $(P' - \beta, \beta, Q' \cup \gamma)$ is not a vertical 3-separation of M' , since $\text{si}(M'/\beta)$ is 3-connected, by Bixby's Lemma. It follows, by orthogonality, that $\alpha \in P'$ and P' is a triangle of M' . Therefore $P' =$

$\{\alpha, \beta, p'\}$ for some $p' \in E(M') - F$. Note also that $p \neq p'$, since the triad $\{\alpha, \beta, \gamma\}$ is independent.

Now $\{\alpha, \beta, \gamma, \delta, p, p'\}$ is a rank-3 subset of M' , with $\{\beta, \delta\} \subseteq B$. Hence, at least one of p and p' is in B^* . Suppose $p' \in B^*$. It follows, by Lemma 2.11, that p' is an (N, B) -strong element of M' . But then $S = \{p', x, y\}$ is a triad of M' that meets the triangle $\{\beta, \gamma, \delta\}$, since $\beta \in \{x, y\}$. By orthogonality, and since $p' \notin F$, we have $\{x, y\} \subseteq F - z$. It follows by z -coclosure that $S \subseteq X$. A similar argument applies if $p \in B^*$. \square

In the next lemma, we show that elements on the non- N -side of a vertical 3-separation that are not N -flexible are not (N, B) -robust.

Lemma 5.7. *Let $z \in B - \{x, y\}$ be an element that is (N, B) -robust but not (N, B) -strong in M' , and let (X, z, Y) be a vertical 3-separation of M' such that $|X \cap E(N)| \leq 1$ and $S \subseteq X$. Then for $e \in X - S$, the element e is N -flexible if and only if e is (N, B) -robust. Moreover, at most one element in $X - S$ is not N -flexible in M'/z , and if such an element μ exists, then $(X - \mu, z, Y \cup \mu)$ is a vertical 3-separation of M' .*

Proof. Clearly, if $e \in X - S$ is N -flexible, then e is (N, B) -robust. Suppose $e \in X - S$ is not N -flexible. By Lemma 2.15, either e is N -deletable but not N -contractible, or e is N -contractible but not N -deletable.

First, suppose that e is N -deletable but not N -contractible. Then $e \in \text{cl}(Y)$, by Lemma 2.15(i). It follows that $((X - e) \cup z, e, Y)$ is a vertical 3-separation of M' , so $\text{co}(M' \setminus e)$ is 3-connected by Bixby's Lemma. Since $e \notin S$, it follows that $e \in B - \{x, y\}$, so e is not (N, B) -robust. Moreover, if e and $e' \in X - S$ are N -deletable but not N -contractible, then $\{z, e, e'\} \subseteq \text{cl}(Y) - Y$, so $r(\{z, e, e'\}) = 2$. But $\{z, e, e'\} \subseteq B$, so $e = e'$.

Now suppose that e is N -contractible but not N -deletable. Let $Y' = \text{cl}(Y) - z$. By Lemma 2.15(ii), $e \in \text{cl}^*(Y') - Y'$ and $z \in \text{cl}(X - (Y' \cup e))$, and there is only one such element e . Observe that Y' and $Y' \cup e$ are exactly 3-separating. Moreover, $(X \cup z) - (Y' \cup e)$ contains a circuit, implying $r^*((X \cup z) - (Y' \cup e)) \geq 3$. Now $((X \cup z) - (Y' \cup e), e, Y')$ is a cyclic 3-separation, so $\text{si}(M'/e)$ is 3-connected by Bixby's Lemma. As $e \notin S$, it follows that $e \in B^*$, so e is not (N, B) -robust.

Suppose μ and μ' are distinct elements of $X - S$ that are not N -flexible. Then, by the foregoing, we may assume that μ is not N -deletable, and μ' is not N -contractible. Note that M/z is the two sum of M_X and M_Y with basepoint z' say, where $M_X \setminus z' = (M/z)|_X$ and $M_Y \setminus z' = (M/z)|_Y$. Since μ is not N -deletable and μ' is not N -contractible, $\{z', \mu\}$ is a cocircuit in M_X , and $\{z', \mu'\}$ is a circuit in M_X ; a contradiction to orthogonality. So at most one element in $X - S$ is not N -flexible.

Now let μ be the unique element in $X - S$ that is not N -flexible, and consider $(X - \mu, z, Y \cup \mu)$. If μ is N -deletable but not N -contractible, then, as $\mu \in \text{cl}(Y)$, clearly $(X - \mu, z, Y \cup \mu)$ is a vertical 3-separation of M' . Suppose that μ is N -contractible but not N -deletable. Since μ is the only element in $X - S$ that is not N -flexible, $\text{cl}(Y) = Y \cup z$, so $\mu \in \text{cl}^*(Y)$. Thus, if $(X - \mu, z, Y \cup \mu)$ is not a vertical 3-separation of M' , then $r(X - \mu) \leq 2$. But $X - \mu$ spans z , and $\{x, y\} \subseteq S \subseteq X - \mu$, so $\{x, y, z\}$ is a triangle of M' contained in B ; a contradiction. \square

Note that a similar argument applies when $z \in B^*$ is an element of M' that is (N, B) -robust but not (N, B) -strong; we omit the proof.

Lemma 5.8. *Let $z \in B^*$ be an element of M' that is (N, B) -robust but not (N, B) -strong, and let (X, z, Y) be a cyclic 3-separation of M' such that $|X \cap E(N)| \leq 1$ and $S \subseteq X$. Then for $e \in X - S$, the element e is N -flexible if and only if e is (N, B) -robust. Moreover, at most one element in $X - S$ is not N -flexible in $M' \setminus z$, and if such an element μ exists, then $(X - \mu, z, Y \cup \mu)$ is a cyclic 3-separation of M' .*

We now come to the main result of the section.

Proposition 5.9. *Let $z \in E(M') - \{x, y\}$ be an element that is (N, B) -robust but not (N, B) -strong in M' . Then M' has a path of 3-separations $(S, z_1, z_2, \dots, z_n, z, Y)$ where the elements in $\{z_1, \dots, z_n\}$ are N -flexible, $|(S \cup \{z_1, \dots, z_n, z\}) \cap E(N)| \leq 1$, and $|S \cup \{z_1, \dots, z_n\}| \geq 3$.*

Proof. Let z be an element of M' that is (N, B) -robust but not (N, B) -strong. First, suppose $z \in B - \{x, y\}$. By Lemma 5.1, there exists a vertical 3-separation (X', z, Y') such that X' is z -closed and $|X' \cap E(N)| \leq 1$. By Lemma 5.5, $S \subseteq X'$. By Lemma 5.7, $X' - S$ contains at most one element that is not (N, B) -robust. If such an element μ exists, let $(X, Y) = (X' - \mu, Y' \cup \mu)$; otherwise, let $(X, Y) = (X', Y')$. Now, by Lemma 5.7 again, (X, z, Y) is a vertical 3-separation of M' where $S \subseteq X$, $|X \cap E(N)| \leq 1$, and every element in $X - S$ is N -flexible. We say that (X, z, Y) is a *good separation for z* in M' . Similarly, if $z \in B^* - \{a, b\}$, then, by the dual of Lemma 5.1, and Lemmas 5.6 and 5.8, there is a cyclic 3-separation (X, z, Y) that is a good separation for z in $(M')^*$. Thus, for each (N, B) -robust element z of M' outside of $\{x, y\}$, there is a good separation for z .

We now show that a good separation induces a path of 3-separations in M' . Let (X, z, Y) be a good separation for z , in either M' or $(M')^*$, and let $Z = X - S$. Consider the partition $(S, Z, z \cup Y)$ of $E(M')$, and note that each $z_i \in Z$ is N -flexible, $|(S \cup Z \cup z) \cap E(N)| \leq 1$, and $|S \cup Z| \geq 3$.

We claim that, for each $z_i \in Z$, there is a path of 3-separations (X_i, z_i, Y_i) of M' such that $S \subseteq X_i$ and $z \cup Y \subseteq Y_i$. In what follows, we assume that $z, z_i \in B$, but the argument is similar if one or both of z, z_i are in B^* . Since z_i is N -flexible in M'/z , we can fix an N -minor of $M'/z/z_i$ on ground set E_N . We may assume that $|X \cap E_N| \leq 1$. As z_i is N -flexible, and hence (N, B) -robust, in M' , but z_i is not (N, B) -strong, there is a vertical 3-separation (X'_i, z_i, Y'_i) of M' where $|X'_i \cap E_N| \leq 1$. By Lemma 5.1, we may assume that X'_i is z_i -closed. Hence $S \subseteq X'_i$ by Lemma 5.5 (in the case that $z_i \in B^*$, we can use Lemma 5.6). Since $|E_N| \geq 4$, it now follows that $|Y \cap Y'_i| \geq |E_N| - 2 \geq 2$. Therefore, by uncrossing $Y \cup z$ and Y'_i , the set $Y \cup Y'_i \cup z$ is 3-separating. Similarly, by uncrossing $Y \cup z$ and $Y'_i \cup z_i$, the set $Y \cup Y'_i \cup \{z, z_i\}$ is 3-separating. Now $(X_i, z_i, Y_i) = (X \cap X'_i, z_i, Y \cup Y'_i \cup z)$ is a path of 3-separations of M' that satisfies the claim.

It now follows from Lemma 5.4 that there is an ordering (z_1, \dots, z_n) of Z such that $(S, z_1, \dots, z_n, z, Y)$ is a path of 3-separations of M' , satisfying the proposition. \square

6. PROOF OF THEOREM 2.30

Let \mathbb{P} be a partial field, let N be a non-binary 3-connected strong \mathbb{P} -stabilizer for the class of \mathbb{P} -representable matroids, and let M be an excluded minor for the class of \mathbb{P} -representable matroids with a pair of elements $\{a, b\}$ such that $M \setminus a, b$ is 3-connected with an N -minor. Let A be a $B \times B^*$ companion \mathbb{P} -matrix of M such that $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$, for some basis B of M . Let $M' = M \setminus a, b$. We assume that M' has no confining set, and that B is a strengthened basis.

Let $S \subseteq E(M')$ be a set containing $\{x, y\}$ and any (N, B) -strong elements of M' , where either $|S| = 2$ or S is a triad. If M' has an element z that is (N, B) -robust but not (N, B) -strong, then, by Proposition 5.9, M' has a path of 3-separations of the form $(S, z'_1, \dots, z'_n, z, Y)$ where each element in $\{z'_1, \dots, z'_n\}$ is N -flexible. In this section, we study such paths of 3-separations, in order to prove Theorem 2.30.

It is convenient to write such a path of 3-separations as $(\{x, y\}, z_1, \dots, z_n, z, Y)$. (In the case where $|S| = 3$, z_1 labels the (N, B) -strong element outside of $\{x, y\}$.) We say that $(\{x, y\}, z_1, \dots, z_n, z, Y)$ is a *good path of 3-separations* for z . Note that $n \geq 1$, since $|S \cup \{z'_1, \dots, z'_n\}| \geq 3$, and observe that z_i is (N, B) -robust for each $i \in \{1, 2, \dots, n\}$.

Lemma 6.1. *Suppose M' has an element z that is (N, B) -robust but not (N, B) -strong, and let $(\{x, y\}, z_1, \dots, z_n, z, Y)$ be a good path of 3-separations for z . Then*

- (i) $\{x, y, z_1\}$ is a triad of M' , and
- (ii) $z_1 \in B^*$.

Proof. First we prove (i). Clearly (i) holds when $z_1 \in S$, so we may assume that $z_1 \notin S$. It suffices to show that $\{x, y, z_1\}$ is not a triangle of M' . Towards a contradiction, suppose $\{x, y, z_1\}$ is a triangle. Then $z_1 \in B^*$, since $\{x, y\} \subseteq B$. Thus $\text{co}(M' \setminus z_1)$ is not 3-connected, as z_1 is (N, B) -robust but not (N, B) -strong.

Let (P, z_1, Q) be a cyclic 3-separation of M' . Since $\{x, y, z_1\}$ is a triangle, it follows from orthogonality that $|P \cap \{x, y\}| = |Q \cap \{x, y\}| = 1$. We shall therefore assume that $x \in P$ and $y \in Q$. Since z_1 is N -flexible, it follows that x and y are N -deletable in M' . Suppose $\text{co}(M' \setminus x)$ is 3-connected. Due to the triangle $\{x, y, z_1\}$, $A_{xz_1} \neq 0$, so a pivot on A_{xz_1} is allowable. Thus $B' = B \triangle \{x, z_1\}$ is a basis, $\{z_1, y, a, b\}$ incriminates (M, A^{xz_1}) , and x is an (N, B') -strong element outside of $\{z_1, y\}$, contradicting that B is a strengthened basis. Hence $\text{co}(M' \setminus x)$ and, by symmetry, $\text{co}(M' \setminus y)$ are not 3-connected.

Now $P \cup z_1$ is exactly 3-separating, and $y \in \text{cl}(P \cup z_1)$, so $(P \cup z_1, y, Q - y)$ is a path of 3-separations of M' , and $y \in \text{cl}(Q - y)$. Similarly, $(P - x, x, Q \cup z_1)$ is a path of 3-separations of M' . If $r(P \cup z_1) \geq 3$ and $r(Q - y) \geq 3$, then $(P \cup z_1, y, Q - y)$ is a vertical 3-separation of M' , in which case $\text{si}(M'/y)$ is not 3-connected, contradicting Bixby's Lemma. Therefore $r(P \cup z_1) \leq 2$ or $r(Q - y) \leq 2$. But if $r(P \cup z_1) \leq 2$, then $M' \setminus z_1$ is 3-connected by Lemma 2.2; a contradiction. Thus $r(Q - y) \leq 2$, and hence $r(Q) \leq 2$. Similarly, it follows that $r(P - x) \leq 2$, and hence $r(P) \leq 2$. Since $x \in P$, $y \in Q$, and $\text{co}(M' \setminus x)$

and $\text{co}(M' \setminus y)$ are not 3-connected, it follows from Lemma 2.2 that $|P| = 3$ and $|Q| = 3$. But now $|E(M')| = 7$, so $n = 1$, and it is readily checked that the (N, B) -robust element z_2 is (N, B) -strong; a contradiction.

We now prove (ii). When z_1 is an (N, B) -strong element of M' , (ii) holds by Lemma 3.1. So we may assume that M' has no (N, B) -strong elements outside of $\{x, y\}$. Towards a contradiction, suppose that $z_1 \in B$. Then $\text{si}(M'/z_1)$ is not 3-connected. Now M' has an (N, B) -robust element $z' \in \{z_2, z_3, \dots, z_n, z\}$ that is either in the closure or coclosure of the triad $\{x, y, z_1\}$. If z' is in the coclosure of $\{x, y, z_1\}$, then $\{x, y, z_1, z'\}$ is a 4-element cosegment of M' , so $\text{si}(M'/z_1)$ is 3-connected by the dual of Lemma 2.2; a contradiction. Thus z' is in the closure of $\{x, y, z_1\}$. Then $(\{x, y, z_1\}, z', E(M') - \{x, y, z_1, z'\})$ is a vertical 3-separation of M' , so $\text{si}(M'/z')$ is not 3-connected. Hence $\text{co}(M' \setminus z')$ is 3-connected by Bixby's Lemma. But, since $\{x, y, z_1, z'\}$ contains a circuit of M' , it follows that $z' \in B^*$. Thus z' is an (N, B) -strong element outside of $\{x, y\}$; a contradiction. \square

Let $(\{x, y\}, z_1, \dots, z_n, z, Y)$ be a good path of 3-separations for some element $z \in E(M')$ that is (N, B) -robust but not (N, B) -strong. Recall that an element $z_i \in \{z_1, z_2, \dots, z_n\}$ is a guts or coguts element according to whether z_i is in the guts or coguts of the 3-separation $(\{x, y, z_1, \dots, z_{i-1}\}, \{z_i, \dots, z_n, z\} \cup Y)$. Similarly, z is a guts or coguts element depending on whether z is in the guts or coguts of the 3-separation $(\{x, y, z_1, \dots, z_n\}, z \cup Y)$.

Lemma 6.2. *Suppose M' has an element z that is (N, B) -robust but not (N, B) -strong, and let $(\{x, y\}, z_1, \dots, z_n, z, Y)$ be a good path of 3-separations for z . Let $z' \in \{z_1, \dots, z_n, z\}$. Then z' is a guts element if and only if $z' \in B$.*

Proof. Suppose that z' is a guts element. Then, by Lemma 6.1(i), $z' \neq z_1$. If z' is not N -deletable, then $z' \in B$ because z' is an (N, B) -robust element. Thus we may assume that z' is N -deletable. Since z' is in the guts of a vertical 3-separation, $\text{co}(M' \setminus z')$ is 3-connected by Bixby's Lemma, so $z' \in B$ because no element in $\{z_2, \dots, z_n, z\}$ is (N, B) -strong in M' .

Conversely, suppose that z' is a coguts element. If $z' = z_1$, then $z' \in B^*$ by Lemma 6.1(ii). So we may assume that $z' \in \{z_2, \dots, z_n, z\}$. If z' is not N -contractible, then $z' \in B^*$ because z' is an (N, B) -robust element. Thus we may assume that z' is N -contractible. Now z' is in the coguts of a cyclic 3-separation of M' , so Bixby's Lemma implies that $\text{si}(M'/z')$ is 3-connected. Thus $z' \in B^*$ because no element in $\{z_2, \dots, z_n, z\}$ is (N, B) -strong in M' . \square

Lemma 6.3. *Suppose M' has elements z and z' that are (N, B) -robust but not (N, B) -strong, and let $(\{x, y\}, z_1, \dots, z_n, z, Y)$ and $(\{x, y\}, z'_1, \dots, z'_{n'}, z', Y')$ be good paths of 3-separations for z and z' respectively. Let $z_{n+1} = z$ and $z'_{n'+1} = z'$. Then*

- (i) $z_1 = z'_1$, and
- (ii) $z_2 = z'_2$, where $z_2 \in B$.

Moreover, $\{x, y, z_1, z_2\}$ is closed in M' .

Proof. By Lemma 6.1, $\{x, y, z_1\}$ and $\{x, y, z'_1\}$ are triads of M' , and $z_1, z'_1 \in B^*$. Since M' has no confining sets, $z_1 = z'_1$.

Consider the element z_2 . Suppose $z_2 \notin B$. Then z_2 is a coguts element by Lemma 6.2, so $\{x, y, z_1, z_2\}$ is a 4-element cosegment, contradicting that M' has no confining set. Thus $z_2 \in B$. Similarly, $z'_2 \in B$.

By Lemma 6.2, z_2 and z'_2 are spanned by the triad $\{x, y, z_1\}$. Now it suffices to show that $\text{cl}(\{x, y, z_1\}) = \{x, y, z_1, z_2\}$. Towards a contradiction, suppose there is some $z'' \in \text{cl}_{M'}(\{x, y, z_1\}) - \{x, y, z_1, z_2\}$. Then, as $\{x, y, z_1\}$ and $\{x, y, z_1, z''\}$ are exactly 3-separating, it follows that $z'' \notin \text{cl}_{M'}^*(\{x, y, z_1\})$, by Lemmas 2.1 and 2.5. Since $z_2 \in B$ is (N, B) -robust but not (N, B) -strong, $r(E(M') - \{x, y, z_1, z''\}) \geq 3$. Thus $(\{x, y, z_1\}, z'', E(M') - \{x, y, z_1, z''\})$ is a vertical 3-separation of M' , so $\text{si}(M'/z'')$ is not 3-connected. Thus $\text{co}(M' \setminus z'')$ is 3-connected by Bixby's Lemma. Moreover, $(\{x, y, z_1, z''\}, z_2, E(M') - \{x, y, z_1, z_2, z''\})$ is a vertical 3-separation of M' , so Lemma 2.15(ii) implies that z'' is N -deletable. Now, if $z'' \in B^*$, then z'' is an (N, B) -strong element; a contradiction. Thus $z'' \in B$. But then the rank-3 set $\text{cl}_{M'}(\{x, y, z_1\})$ contains $\{x, y, z_2, z''\}$, and $\{x, y, z_2, z''\} \subseteq B$; a contradiction. \square

We need the following result of Whittle and Williams.

Lemma 6.4 ([24, Lemma 2.13]). *Let M_0 be a 3-connected matroid with a triad $\{c, d, e\}$ and circuit $\{c, d, e, f\}$. Then at least one of the following holds:*

- (i) *either $\text{co}(M_0 \setminus c)$ or $\text{co}(M_0 \setminus e)$ is 3-connected,*
- (ii) *there exist $c', e' \in E(M_0)$ such that both $\{c, c', d\}$ and $\{d, e, e'\}$ are triangles, or*
- (iii) *there exists $g \in E(M_0)$ such that $\{c, d, e, g\}$ is a 4-element cosegment.*

We require one more definition. Let B be a strengthened basis, and let A be a $B \times B^*$ companion \mathbb{P} -matrix of M such that $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$. Suppose that M' has no (N, B) -strong elements outside of $\{x, y\}$. We say that B is *bolstered* if for any $B_1 \times B_1^*$ companion \mathbb{P} -matrix A_1 where $\{x_1, y_1, a, b\}$ incriminates (M, A_1) , with $\{x_1, y_1\} \subseteq B_1$ and $\{a, b\} \subseteq B_1^*$, the number of (N, B) -robust elements of M' outside of $\{x, y\}$ is at least the number of (N, B_1) -robust elements of M' outside of $\{x_1, y_1\}$. When M' has an (N, B) -strong element u of M' outside of $\{x, y\}$, where $u \in B^*$ by Lemma 3.1, then we say that B is *bolstered* if for any $B_1 \times B_1^*$ companion \mathbb{P} -matrix A_1 where $\{x, y, a, b\}$ incriminates (M, A_1) , with $\{x, y\} \subseteq B_1$ and $\{u, a, b\} \subseteq B_1^*$, the number of (N, B) -robust elements of M' is at least the number of (N, B_1) -robust elements of M' . (Loosely speaking, a basis B is bolstered if no allowable pivot increases the number of (N, B) -robust elements.) Note that for any strengthened basis B , we can perform allowable pivots in order to obtain a bolstered basis B' , where B' is also strengthened.

We can now show that either M is bounded relative to N , or M' has at most two elements outside of $\{x, y\}$ that are (N, B) -robust but not (N, B) -strong. Recall that when F is a 4-element fan with ordering (f_1, f_2, f_3, f_4) such that $\{f_1, f_2, f_3\}$ is a triangle, and B' is a basis, we say that (f_1, f_2, f_3, f_4) is a *type-II fan relative to B'* if $F \cap B' = \{f_1, f_3, f_4\}$.

Lemma 6.5. *Suppose B is a bolstered basis. Then there are at most two elements outside of $\{x, y\}$ that are (N, B) -robust but not (N, B) -strong in M' . Moreover, either*

- (i) M' has a maximal type-II fan (z, u, x, y) relative to B , where u is (N, B) -strong, and z is the only element outside of $\{x, y\}$ that is (N, B) -robust but not (N, B) -strong;
- (ii) M' has a maximal fan (w, z, x, u, y) , where u is (N, B) -strong, $z \in B$ is N -flexible, $w \in B^*$, and the elements outside of $\{x, y\}$ that are (N, B) -robust but not (N, B) -strong are contained in $\{z, w\}$; or
- (iii) $|E(M)| \leq |E(N)| + 7$.

Proof. Suppose that z is (N, B) -robust but not (N, B) -strong, let $(\{x, y\}, z_1, z_2, \dots, z_n, z, Y)$ be a good path of 3-separations for z , where $n \geq 1$, and let $z_{n+1} = z$. By Lemmas 6.1 to 6.3, $\{x, y, z_1\}$ is a triad with $z_1 \in B^*$, and $z_2 \in B$ is a guts element. Observe that $\{x, y, z_1, z_2\}$ is either a 4-element fan or a circuit of M' . In the case that $\{x, y, z_1, z_2\}$ is a 4-element fan, $\{z_2, x, y\}$ is not a triangle, since $z_2 \in B$, so we may assume, up to swapping x and y , that this fan has ordering (z_2, z_1, x, y) , where $\{z_2, z_1, x\}$ is a triangle.

6.5.1. *Suppose (z_2, z_1, x, y) is a maximal fan and $\text{co}(M' \setminus x)$ is not 3-connected. Then case (i) of the lemma holds, with $z = z_2$ and $u = z_1$.*

Subproof. By Lemma 2.12, $\text{si}(M'/z_1) \cong \text{co}(M' \setminus x)$, so $\text{si}(M'/z_1)$ is not 3-connected. As z_1 is (N, B) -robust, Bixby's Lemma implies that z_1 is (N, B) -strong. We claim that z_2 is the only element outside of $\{x, y\}$ that is (N, B) -robust but not (N, B) -strong. Suppose $z' \in E(M') - \{x, y, z_1, z_2\}$ is (N, B) -robust but not (N, B) -strong, and let $(\{x, y\}, z'_1, \dots, z'_n, z', Y')$ be a good path of 3-separations for z' . Let $z' = z'_{n'+1}$. By Lemma 6.3, $z'_1 = z_1$, $z'_2 = z_2$, $n' \geq 2$, and z'_3 is an (N, B) -robust coguts element. By Lemma 6.2, $z'_3 \in B^*$. Now z'_3 is in a cocircuit C^* contained in $\{x, y, z_1, z_2, z'_3\}$. Since $\{x, y, z_1\}$ is a triad, $\{x, y, z_1\} \not\subseteq C^*$. Moreover, if $y \in C^*$, then by cocircuit elimination with $\{x, y, z_1\}$, there is a cocircuit contained in $\{x, z_1, z_2, z'_3\}$, and this cocircuit also contains z'_3 . So we may assume that $y \notin C^*$. Since M' has no confining sets, $z_2 \in C^*$. Since (z_2, z_1, x, y) is maximal, neither $\{x, z_2, z'_3\}$ nor $\{z_1, z_2, z'_3\}$ is a triad. So $\{x, z_1, z_2, z'_3\}$ is a cocircuit of M' . Now $\{z_2, z_1, x\}$ is not contained in a 4-element segment by orthogonality, and $\{x, z_2\}$ is not contained in a triad because (z_2, z_1, x, y) is maximal. Therefore, by the dual of Lemma 6.4, either $\text{si}(M'/z_2)$ or $\text{si}(M'/z_1)$ is 3-connected. But $\text{si}(M'/z_2)$ is not 3-connected because z_2 is not (N, B) -strong, and $\text{si}(M'/z_1) \cong \text{co}(M' \setminus x)$ is not 3-connected; a contradiction. We deduce that z_2 is the only element outside of $\{x, y\}$ that is (N, B) -robust but not (N, B) -strong. \square

6.5.2. *Suppose $\{z_2, z_1, x, y\}$ is contained in a fan (w, z_2, x, z_1, y) , for some $w \in E(M') - \{x, y, z_1, z_2\}$, and $\text{co}(M' \setminus x)$ is not 3-connected. Then case (ii) of the lemma holds, with $z = z_2$ and $u = z_1$.*

Subproof. As $z_1 \in B^*$ is (N, B) -robust, and $\text{co}(M' \setminus z_1)$ is 3-connected by Lemma 2.11, z_1 is (N, B) -strong. Since z_1 is N -deletable, and $\{x, y\}$ is a series pair in $M' \setminus z_1$, it follows that x is N -contractible. Similarly, since x

is N -contractible, z_2 is N -deletable. As $z_2 \in B$ is (N, B) -robust, z_2 is N -flexible. Moreover, as z_2 is N -deletable, w is N -contractible. Now, if $w \in B$, then w is (N, B) -strong by Lemma 2.11; so $w \in B^*$.

Next we show that the fan (w, z_2, x, z_1, y) is maximal. First, observe that $\{z_1, y\}$ is not contained in a triangle by Lemma 6.3. Suppose $\{z_2, w\}$ is contained in a triangle $\{z_2, w, z'\}$ say. Since z_2 is N -contractible, it follows that z' is N -deletable. By Lemma 2.11, $\text{co}(M \setminus z')$ is 3-connected, so, as the (N, B) -strong elements are contained in $\{x, y, z_1\}$, we have $z' \in B$. Now, as $\{z_2, w, z'\}$ is a triangle, $A_{xw} = A_{yw} = 0$ and $A_{z_2w} \neq 0$, so a pivot on A_{z_2w} is allowable. But then $B' = B \triangle \{z_2, w\}$ is a basis, and z_2 is an (N, B') -strong element in $B' - \{x, y\}$, contradicting Lemma 3.1.

Now, if the elements outside of $\{x, y\}$ that are (N, B) -robust but not (N, B) -strong are contained in $\{z_2, w\}$, then 6.5.2 holds.

Suppose there is some N -contractible element $w' \in \text{cl}^*(\{x, y, z_1, z_2\}) - \{x, y, z_1, z_2, w\}$. Then $\{y, w, w'\}$ is in the coclosure of the 3-separating triangle $\{x, z_1, z_2\}$, so, as M' is 3-connected, $\{y, w, w'\}$ is a triad. Recall that $w \in B^*$. By Bixby's Lemma, $\text{si}(M'/w')$ is 3-connected. Since w' is N -contractible, and the (N, B) -strong elements of M' are contained in $\{x, y, z_1\}$, it follows that $w' \in B^*$. Now $\{x, y, z_1, w, w'\}$ is a confining set; a contradiction.

Suppose there is some $w' \in E(M') - \{x, y, z_1, z_2, w\}$ that is (N, B) -robust. Let $(\{x, y\}, z'_1, z'_2, \dots, z'_{n'}, w', Y')$ be a good path of 3-separations for w' , and let $z'_{n'+1} = w'$. It follows from Lemmas 6.2 and 6.3 and the preceding paragraph that $z'_1 = z_1$, $z'_2 = z_2$, $z'_3 = w$, and z'_4 is a guts element, so $z'_4 \in B$. We work towards a contradiction.

We first claim that $\{y, z_1, z_2, w, z'_4\}$ is a circuit of M' . Certainly, z'_4 is in a circuit contained in $\{x, y, z_1, z_2, w, z'_4\}$. If this circuit contains x , then, by circuit elimination with the triangle $\{z_2, x, z_1\}$, there is a circuit contained in $\{y, z_1, z_2, w, z'_4\}$. So we may assume that there is a circuit C contained in $\{y, z_1, z_2, w, z'_4\}$, which may or may not contain z'_4 . By orthogonality with the triad $\{w, z_2, x\}$, either C contains $\{w, z_2\}$ or $C \cap \{w, z_2\} = \emptyset$. But in the latter case, $\{z_1, y, z'_4\}$ is a triangle of M' , contradicting the maximality of the fan (w, z_2, x, z_1, y) . So C contains $\{w, z_2\}$ and, similarly, $\{z_1, y\}$. Finally, if $z'_4 \notin C$, then $\{w, z_2, x, z_1, y\}$ is 2-separating; a contradiction. This proves the claim.

By orthogonality, the only triads containing x are $\{w, z_2, x\}$ and $\{x, z_1, y\}$, so $\text{co}(M' \setminus x) \cong M' \setminus x / z_1, z_2$. Let $M'' = M' \setminus x / z_1, z_2$. As $\{w, z_2, z_1, y, z'_4\}$ is a circuit of M' , the set $T = \{w, z'_4, y\}$ is a triangle of M'' . Let (P, Q) be a 2-separation of M'' , where $|P \cap T| \geq 2$. Now $(\text{fcl}_{M''}(P), Q - \text{fcl}_{M''}(P))$ is also a 2-separation of M'' , so we may also assume, without loss of generality, that P is fully closed. In particular, $T \subseteq P$. Since $\{z_1, z_2\} \subseteq \text{cl}^*_{M' \setminus x}(P)$, we have that $(P \cup \{z_1, z_2\}, Q)$ is a 2-separation in $M' \setminus x$. As $x \in \text{cl}(P \cup \{z_1, z_2\})$, it follows that $(P \cup \{z_1, z_2, x\}, Q)$ is a 2-separation of M' ; a contradiction.

We deduce that no $w' \in E(M') - \{x, y, z_1, z_2, w\}$ is (N, B) -robust, so the elements of M' that are (N, B) -robust but not (N, B) -strong are contained in $\{z_2, w\}$, as required. \square

Suppose that (z_2, z_1, x, y) is a 4-element fan, and $\text{co}(M' \setminus x)$ is not 3-connected. If $\{z_2, z_1\}$ is contained in a triad, then x is a spoke end of

a 4-element fan, contradicting Lemma 2.11; whereas if y is in a triangle, then, by orthogonality, this contradicts Lemma 6.3. Thus either the fan (z_2, z_1, x, y) is maximal, in which case (i) holds by 6.5.1; or it is contained in a fan (w, z_2, x, z_1, y) for some $w \in E(M') - \{x, y, z_1, z_2\}$, in which case (ii) holds by 6.5.2. So we may assume that when (z_2, z_1, x, y) is a 4-element fan, $\text{co}(M' \setminus x)$ is 3-connected.

Now consider the case where $\{x, y, z_1, z_2\}$ is a circuit. Suppose $\{x, y, z_1\}$ is contained in a 4-element cosegment $\{x, y, z_1, f\}$. Then, as M' has no confining sets, $f \in B$. Since $z_1 \in B^*$ is (N, B) -robust, it follows that f is N -contractible. But $\text{si}(M'/f)$ is 3-connected by the dual of Lemma 2.2, so f is (N, B) -strong; a contradiction. Now, since $\{x, y, z_1, z_2\}$ is closed by Lemma 6.3, it follows from Lemma 6.4 that either $\text{co}(M' \setminus x)$ or $\text{co}(M' \setminus y)$ is 3-connected. Thus, when $\{x, y, z_1, z_2\}$ is a circuit, we may assume without loss of generality that $\text{co}(M' \setminus x)$ is 3-connected.

Now, in either case, we may assume that $\text{co}(M' \setminus x)$ is 3-connected.

6.5.3. $A_{pa} = A_{pb} = 0$ for all $p \in B - \{x, y, z_2\}$, and z_1 is (N, B) -strong.

Subproof. Either $\{x, y, z_1, z_2\}$ is a circuit, or this set is a 4-element fan containing the triad $\{x, z_1, z_2\}$. If x is in a 4-element cosegment of M' , then, by orthogonality, this cosegment intersects the circuit $\{x, y, z_1, z_2\}$ or $\{x, z_1, z_2\}$ in three elements. But this implies that $\{x, y, z_1\}$ is contained in a 4-element cosegment; a contradiction. So $M' \setminus x$ is 3-connected up to series pairs. Similarly, z_1 is not in a 4-element cosegment of M' .

Next we claim that x is N -deletable. Observe that $(\{x, y, z_1\}, z_2, E(M') - \{x, y, z_1, z_2\})$ is a vertical 3-separation, where M'/z_2 has an N -minor since $z_2 \in B$ is (N, B) -robust. Since $\{x, y, z_1\}$ is a triad, $E(M') - \{x, y, z_1\}$ is closed. Moreover, $x \in \text{cl}(\{y, z_1, z_2\})$, so $x \notin \text{cl}_{M'}^*(E(M') - \{x, y, z_1, z_2\})$. Thus, by Lemma 2.15(ii), the element x is N -deletable.

We work towards showing that $a, b \in \text{cl}_M(\{x, y, z_2\})$. Observe that $A_{xz_1} \neq 0$ because $\{x, y, z_1, z_2\}$ is a circuit with $\{x, y, z_2\} \subseteq B$. So a pivot on A_{xz_1} is allowable. Now $\{z_1, y, a, b\}$ incriminates (M, A^{xz_1}) . Let $B' = B \Delta \{x, z_1\}$. Then x is an (N, B') -strong element outside of $\{z_1, y\}$. By Lemma 3.4, $M' \setminus x$ has a series pair that meets $\{z_1, y\}$ and is contained in an unstable triple of $M \setminus a, x$ or $M \setminus b, x$. Since $\{z_1, y\}$ is a series class of $M' \setminus x$, the series pair $\{z_1, y\}$ is contained in an unstable triple $\{z_1, y, b\}$ of $M \setminus a, x$, up to swapping labels on a and b . So $b \in \text{cl}_M(\{z_1, y\})$. Since $z_1 \in \text{cl}(\{x, y, z_2\})$, it follows that $b \in \text{cl}_M(\{x, y, z_2\})$.

As B is a strengthened basis and x is an (N, B') -strong element outside of $\{z_1, y\}$, it follows that z_1 is an (N, B) -strong element outside of $\{x, y\}$. Since z_1 is not in a 4-element cosegment of M' , the matroid $M' \setminus z_1$ is 3-connected up to series pairs. Thus, by Lemma 3.4 again, $M' \setminus z_1$ has a series pair that meets $\{x, y\}$ and is contained in an unstable triple of $M \setminus a, z_1$ or $M \setminus b, z_1$. It now follows that this series pair is $\{x, y\}$, so either $a \in \text{cl}_M(\{x, y\})$ or $b \in \text{cl}_M(\{x, y\})$. But in the latter case, $\{x, y, z_1\}$ is a triangle, contradicting Lemma 6.1(i). So $a \in \text{cl}_M(\{x, y\})$. Now $a, b \in \text{cl}_M(\{x, y, z_2\})$, so $A_{pa} = A_{pb} = 0$ for all $p \in B - \{x, y, z_2\}$. Thus 6.5.3 holds. \square

6.5.4. *There are no N -contractible elements of M' outside of $\{x, y, z_1, z_2\}$.*

Subproof. Suppose that M' has an N -flexible element $q \in B^* - z_1$. Then, since q is not (N, B) -strong in M' , the matroid $\text{co}(M' \setminus q)$ is not 3-connected. Hence $\text{si}(M'/q)$ is 3-connected by Bixby's Lemma. By Lemma 6.3, $q \notin \text{cl}(\{x, y, z_1\}) = \text{cl}(\{x, y, z_2\})$, so $A_{pq} \neq 0$ for some $p \in B - \{x, y, z_2\}$. Since $A_{pa} = A_{pb} = 0$, by 6.5.3, a pivot on A_{pq} is allowable. But then $B' = B \Delta \{p, q\}$ has an (N, B') -strong element q in $B' - \{x, y\}$, contradicting Lemma 3.1. Thus M' has no N -flexible elements in $B^* - z_1$.

Suppose that M' has an (N, B) -robust element $p \in B - \{x, y, z_2\}$. Let $(\{x, y\}, z'_1, \dots, z'_{n'}, p, Y')$ be a good path of 3-separations for p . By Lemma 6.3, $z'_1 = z_1$, $z'_2 = z_2$, and $\{x, y, z_1, z_2\}$ is closed. Hence, as p is a guts element by Lemma 6.2, $n' \geq 3$. But then $z'_3 \in B^* - z_1$ is N -flexible; a contradiction. Therefore no element in $B - \{x, y, z_2\}$ is N -contractible.

For each element $q \in B^* - z_1$, there is some $p \in B - \{x, y, z_2\}$ such that A_{pq} is non-zero, by Lemma 6.3, so a pivot on A_{pq} is allowable. Since B is bolstered, there are at least as many (N, B) -robust elements as $(N, B \Delta \{p, q\})$ -robust elements, so M' has no N -contractible elements in $B^* - z_1$. \square

6.5.5. *There is at most one element outside of $\{x, y\}$ that is (N, B) -robust but not (N, B) -strong.*

Subproof. By 6.5.4, each (N, B) -robust element of M' outside of $\{x, y, z_1, z_2\}$ is in B^* , so any such element is a coguts element by Lemma 6.2. Suppose q and q' are distinct (N, B) -robust elements of M' in $B^* - z_1$. Then $(\{x, y\}, z_1, z_2, q, Y)$ and $(\{x, y\}, z_1, z_2, q', Y')$ are the good paths of 3-separations for q and q' respectively, otherwise there is an N -contractible element in $B^* - z_1$. Now $(\{x, y, z_1, z_2, q\}, q', Y - q')$ is a cyclic 3-separation, and $q \in \text{cl}_{M'}^*(Y - q')$, so $q \in B^* - z_1$ is N -contractible in M' by the dual of Lemma 2.15(ii); a contradiction. Hence there is at most one (N, B) -robust element of M' outside of $\{x, y, z_1, z_2\}$. \square

By 6.5.5, it now suffices to show that $|E(M')| \leq |E(N)| + 5$. Towards a contradiction, suppose that $|E(M')| \geq |E(N)| + 6$. Let R be the set consisting of $\{x, y, z_1, z_2\}$ and the (N, B) -robust element of M' outside of $\{x, y, z_1, z_2\}$, if such an element exists. So the set of (N, B) -robust elements of M' is contained in R , where $|R| \leq 5$. Since $|E(M')| \geq |E(N)| + 6$, there is an element p outside of R that is either N -deletable or N -contractible in M' , but is not (N, B) -robust in M' . By 6.5.4, $p \in B - R$ and p is N -deletable. Since z_1 is in a circuit of M' contained in $\{x, y, z_1, z_2\}$, it follows from orthogonality that $p \notin \text{cl}_{M'}^*(\{z_1, z_2\}) = \text{cl}_{M'}^*(R \cap B^*)$. Thus there is some $q \in B^* - R$ such that $A_{pq} \neq 0$. Since $A_{pa} = A_{pb} = 0$, by 6.5.3, a pivot on A_{pq} is allowable. Again letting $B' = B \Delta \{p, q\}$, we see there are more (N, B') -robust elements than (N, B) -robust elements, so B is not bolstered; a contradiction. We deduce that $|E(M')| \leq |E(N)| + 5$, as required. \square

Lemma 6.6. *Suppose B is a bolstered basis. If M' has no (N, B) -robust elements outside of $\{x, y\}$, then $M \setminus a, b$ is N -fragile.*

Proof. Suppose M' has no (N, B) -robust elements outside of $\{x, y\}$. Since the elements outside of $\{x, y\}$ are not (N, B) -robust, it suffices, by symmetry, to show that $M' \setminus x$ has no N -minor. Towards a contradiction, suppose that x is N -deletable. There is some $x' \in B^* - \{a, b\}$ such that $A_{xx'} \neq 0$ because x is

not a coloop of M' , so a pivot on $A_{xx'}$ is allowable. Let $B' = B \Delta \{x, x'\}$. Now x is an (N, B') -robust element of M' outside of $\{x', y\}$, contradicting the fact that B is bolstered. We deduce that x is not N -deletable, as required. \square

We now prove Theorem 2.30, which we restate here for ease of reference.

Theorem 6.7. *Let M be an excluded minor for the class of \mathbb{P} -representable matroids, and let N be a non-binary 3-connected strong \mathbb{P} -stabilizer for the class of \mathbb{P} -representable matroids. Suppose M has a pair of elements $\{a, b\}$ such that $M \setminus a, b$ is 3-connected with an N -minor. Then either*

- (i) $|E(M)| \leq |E(N)| + 9$, or
- (ii) M has a $B \times B^*$ companion \mathbb{P} -matrix A for which $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$, and either
 - (a) $M \setminus a, b$ is N -fragile, and $M \setminus a, b$ has at most one (N, B) -robust element outside of $\{x, y\}$, where if such an element u exists, then $u \in B^* - \{a, b\}$ is an (N, B) -strong element of $M \setminus a, b$, and $\{u, x, y\}$ is a coclosed triad of $M \setminus a, b$, or
 - (b) $M \setminus a, b$ is not N -fragile, but there is an element $u \in B^* - \{a, b\}$ that is (N, B) -strong in $M \setminus a, b$; either
 - (I) the N -flexible, and (N, B) -robust, elements of $M \setminus a, b$ are contained in $\{u, x, y\}$, or
 - (II) the N -flexible, and (N, B) -robust, elements of $M \setminus a, b$ are contained in $\{u, x, y, z\}$, where $z \in B$, and (z, u, x, y) is a maximal fan of $M \setminus a, b$, or
 - (III) the N -flexible, and (N, B) -robust, elements of $M \setminus a, b$ are contained in $\{u, x, y, z, w\}$, where $z \in B$, $w \in B^*$, and (w, z, x, u, y) is a maximal fan of $M \setminus a, b$; the unique triad in $M \setminus a, b$ containing u is $\{u, x, y\}$; and M has a cocircuit $\{x, y, u, a, b\}$ and a triangle $\{d, x, y\}$ for some $d \in \{a, b\}$.

Moreover, B is a bolstered basis.

Proof. It follows from Proposition 4.1 that M' has either a confining set or a strengthened basis B . If M' has a confining set, then (i) holds by Proposition 4.16. Assume that M' has a strengthened basis B and that (i) does not hold, so $|E(M)| \geq |E(N)| + 10$ and M' has no confining sets. We may assume that the strengthened basis B is chosen to be bolstered. If M' has no (N, B) -robust elements outside of $\{x, y\}$, then (ii)(a) holds by Lemma 6.6. We shall therefore assume M' has an (N, B) -robust element outside of $\{x, y\}$.

We distinguish two cases. First, suppose that all (N, B) -robust elements of M' outside of $\{x, y\}$ are (N, B) -strong. Then M' has exactly one (N, B) -strong element u , and $\{u, x, y\}$ is a triad of M' by Proposition 4.1. Since M' has no confining sets, Lemma 3.2 implies that $M' \setminus u$ is 3-connected up to series pairs; in particular, the triad $\{u, x, y\}$ is coclosed. If M' is N -fragile, then (ii)(a) holds. Suppose then that M' is not N -fragile. Since N -flexible elements are (N, B) -robust, it follows that the N -flexible elements of M' are contained in $\{u, x, y\}$. To show that (ii)(b)(I) holds, it remains to prove that $\{d, x, y\}$ is a triangle of M for some $d \in \{a, b\}$, the unique triad in $M \setminus a, b$ containing u is $\{u, x, y\}$, and $\{x, y, u, a, b\}$ is a cocircuit of M' . Since $M' \setminus u$

is 3-connected up to series pairs, the former follows from Lemma 3.4. We return to the latter two claims momentarily.

Second, suppose that some (N, B) -robust element of M' outside of $\{x, y\}$ is not (N, B) -strong. Since $|E(M)| \geq |E(N)| + 10$, Lemma 6.5 implies that u is (N, B) -strong, and either M' has a maximal type-II fan (z, u, x, y) relative to B , or M' has a maximal fan (w, z, x, u, y) such that $z \in B$ and $w \in B^*$, where $\{u, x, y, z\}$ or $\{u, x, y, z, w\}$, respectively, contains all of the (N, B) -robust elements of M' . Hence M' is not N -fragile, and $\{u, x, y, z\}$, or $\{u, x, y, z, w\}$, contains all of the N -flexible elements of M' . Lemma 3.4 implies that $\{d, x, y\}$ is a triangle of M for some $d \in \{a, b\}$.

Now, in either of the two cases, M' has an (N, B) -strong element u .

6.7.1. $\{x, y, u, a, b\}$ is a cocircuit of M .

Subproof. By orthogonality, $\{x, y\}$ is not contained in a 4-element segment of M' , so there is at most one element in B^* that is in a triangle of M' with $\{x, y\}$. Thus, as $|E(M)| \geq |E(N)| + 10$, there is either some $p \in B - \{x, y\}$ that is N -deletable but not (N, B) -robust, or some $q \in B^* - \{a, b, u\}$ that is N -contractible but not (N, B) -robust such that $\{q, x, y\}$ is not a triangle. In the former case, as $\{p, a, b\}$ is not a triad of M since $M \setminus a, b$ is 3-connected, we can choose $q \in B^* - \{a, b, u\}$ such that the entry A_{pq} is non-zero. In the latter case, we can choose $p \in B - \{x, y\}$ so that the entry A_{pq} is non-zero, since $\{q, x, y\}$ is not a triangle. Now, if $A_{xq} = 0$ and $A_{yq} = 0$, then the pivot on A_{pq} is allowable, in which case $B \Delta \{p, q\}$ is a basis, and there are more $(N, B \Delta \{p, q\})$ -robust elements than (N, B) -robust elements in M' , contradicting the fact that B is bolstered. So we may assume that $A_{yq} \neq 0$. Now a pivot on A_{yq} is allowable, so A^{yq} is a companion \mathbb{P} -matrix where $\{x, q, a, b\}$ incriminates (M, A^{yq}) . If $\{b, u, x, y\}$ is a cocircuit of M , then $(A^{yq})_{xa} = 0$ because $\{b, y, u\}$ cospans x , contradicting that the bad submatrix $A^{yq}[\{x, q, a, b\}]$ has no zero entries. So $\{b, u, x, y\}$, and similarly $\{a, u, x, y\}$, are not cocircuits of M . Therefore $\{x, y, u, a, b\}$ is a cocircuit of M . \square

6.7.2. $\{x, y\}$ is the only series pair of $M' \setminus u$.

Subproof. Suppose $\{p, q\}$ is a series pair of $M' \setminus u$ that is distinct from $\{x, y\}$. Since $\{u, x, y\}$ is a coclosed triad of M' , the pairs $\{x, y\}$ and $\{p, q\}$ are not contained in the same series class of $M' \setminus u$; in particular, they are disjoint. As $\{u, p, q\}$ is a triad of M' and u is an N -deletable element in B^* , both p and q are N -contractible in M' , and at least one of p and q is in $B - \{x, y\}$. So p , say, is an (N, B) -robust element in $B - \{x, y\}$. Since $\{u, p, q\}$ is a triad of M' , for some $q \in E(M') - \{u, p, x, y\}$, it now follows that we are in the case where (ii)(b)(III) holds. Now M' has a 5-element fan F with ordering (w, p, x, u, y) , where $q \notin F$ and q is N -contractible. Since q is not (N, B) -robust, $q \in B^*$. Moreover, as $\{y, w, q\} \subseteq \text{cl}_{M'}^*(\{u, x, p\}) - \{u, x, p\}$, where $\{u, x, p\}$ is 3-separating, it follows that $\{y, w, q\}$ is a triad of M' . Now $\{x, y, u, p, q\}$ is a confining set; a contradiction. \square

Finally, either (I), (II), or (III) of (ii)(b) holds, by 6.7.1 and 6.7.2. \square

7. SPIKE-LIKE 3-SEPARATORS

Suppose that M is an excluded minor for the class of \mathbb{P} -representable matroids, for some partial field \mathbb{P} , with a minor N where N is a 3-connected strong \mathbb{P} -stabilizer. By Theorem 2.29, if M has no spike-like 3-separator, then, after replacing M by a Δ - Y -equivalent matroid, and possibly dualising, we obtain a matroid with a deletion pair with respect to N or N^* . In this section, we show that in the case that M has a spike-like 3-separator, $|E(M)|$ is bounded relative to $|E(N)|$.

We require the following lemma which shows, in particular, that an element that is in a quad but not in a triangle (or, dually, a triad) can be contracted (or deleted, respectively) without destroying 3-connectivity.

Lemma 7.1 ([23, Lemma 3.8]). *Let C^* be a rank-3 cocircuit of a 3-connected matroid M . If $x \in C^*$ has the property that $\text{cl}_M(C^*) - x$ contains a triangle of M/x , then $\text{si}(M/x)$ is 3-connected.*

Lemma 7.2. *Let \mathbb{P} be a partial field, let N be a non-binary 3-connected strong stabilizer for the class of \mathbb{P} -representable matroids, and let M be an excluded minor for the class of \mathbb{P} -representable matroids, where M has an N -minor. If M has a spike-like 3-separator P such that at most one element of $E(M) - E(N)$ is not in P , then $|E(M)| \leq |E(N)| + 5$.*

Proof. Towards a contradiction, suppose that $|E(M)| \geq |E(N)| + 6$. By the definition of a spike-like 3-separator, there is a partition $\{L_1, \dots, L_t\}$ of P such that $|L_i| = 2$ for each $i \in \{1, \dots, t\}$, and $L_i \cup L_j$ is a quad for all distinct $i, j \in \{1, \dots, t\}$, where $t \geq 3$. Since at most one element of $E(M) - E(N)$ is not in P , we have $|P - E(N)| \geq 5$.

Up to possibly replacing (M, N) with (M^*, N^*) , there are distinct elements $a, b \in P$ such that $\{a, b\}$ is N -deletable, $a \in L_i$, and $b \in L_j$, with $i \neq j$. It follows from orthogonality, and the fact that $i \neq j$ and $t \geq 3$, that if $\{a, b\}$ is contained in a triad, then this triad meets $L_{i'}$ for each $i' \in \{1, \dots, t\}$. But then $t = 3$ and $r^*(P) = 3$, implying $\lambda(P) = 1$; a contradiction. Thus, by the dual of Lemma 7.1, $M \setminus a$ and $M \setminus b$ are 3-connected, and $M \setminus a, b$ is 3-connected up to series classes. Thus $\{a, b\}$ is a weak deletion pair. By Theorems 2.18 and 2.20, there exists a $B \times B^*$ companion \mathbb{P} -matrix A with $\{x, y\} \subseteq B$ and $\{a, b\} \subseteq B^*$ such that $\{x, y, a, b\}$ incriminates (M, A) .

Since $L_i \cup L_j$ is a cocircuit, there is some $u \in (L_i \cup L_j) \cap B$. As u is in a series pair of $M \setminus a, b$, the element u is N -contractible in $M \setminus a, b$, and $M \setminus a, b/u$ is 3-connected up to series classes. Without loss of generality, we may assume that $u \in L_i$. By the definition of a spike-like 3-separator, $L_i = \{a, u\}$ is not contained in a triangle. Thus, if u is in a triangle, then, by orthogonality with the cocircuits $L_i \cup L_{j'}$ for $j' \in \{1, \dots, t\} - i$, this triangle meets each $L_{j'}$. But then $t = 3$ and $r(P) = 3$, implying $\lambda(P) = 1$; a contradiction. So M/u is 3-connected by Lemma 7.1. It now follows that $\text{co}(M \setminus a/u)$ and $\text{co}(M \setminus b/u)$ are 3-connected. In particular, $M \setminus a/u$ and $M \setminus b/u$ are N -stable, and $M \setminus a, b/u$ is connected. Thus, by Lemma 2.24, M/u is not strongly \mathbb{P} -stabilized by N . But, as M/u is 3-connected, and hence N -stable, this contradicts Lemma 2.21. \square

The following is a consequence of Lemma 7.2 and Theorem 2.29.

Corollary 7.3. *Let \mathbb{P} be a partial field, let M be an excluded minor for the class of \mathbb{P} -representable matroids, and let N be a non-binary 3-connected strong stabilizer for the class of \mathbb{P} -representable matroids, where M has an N -minor. Suppose that $|E(M)| \geq |E(N)| + 10$. Then, there exists a matroid M_0 , where M_0 is obtained from M by at most one Δ - Y or Y - Δ exchange, and $(M_1, N_1) \in \{(M_0, N), (M_0^*, N^*)\}$ such that M_1 has a pair of elements $\{a, b\}$ for which $M_1 \setminus a, b$ is 3-connected and has an N_1 -minor.*

8. PROOF OF THEOREM 2.31

Let M be an excluded minor for the class of \mathbb{P} -representable matroids, for some partial field \mathbb{P} , and let N be a non-binary 3-connected strong \mathbb{P} -stabilizer for the class of \mathbb{P} -representable matroids.

In this section we prove Theorem 2.31. We first address a few more cases where we can bound $|E(M)|$ relative to $|E(N)|$.

Lemma 8.1. *Suppose M has a pair of elements $\{a, b\}$ such that $M \setminus a, b$ is 3-connected with an N -minor. If (ii)(b) of Theorem 6.7 holds, and $\{a, b\} \subseteq \text{cl}_M(\{x, y\})$, then $|E(M)| \leq |E(N)| + 7$.*

Proof. Suppose that (ii)(b) of Theorem 6.7 holds, and $\{a, b\} \subseteq \text{cl}_M(\{x, y\})$, but $|E(M)| \geq |E(N)| + 8$. Then there is at least one element in $E(M) - \{x, y\}$ that is N -deletable or N -contractible in M' but not (N, B) -robust, where B is a bolstered basis.

Suppose that p is N -deletable but not (N, B) -robust. Then $p \in B - \{x, y\}$. Now $A_{pa} = A_{pb} = 0$ because $\{a, b\} \subseteq \text{cl}_M(\{x, y\})$. We claim that there is some element $q \in B^* - \{a, b\}$ that is not (N, B) -robust and $A_{pq} \neq 0$. By Lemma 6.5, there is a single element $u \in B^*$ that is (N, B) -strong in M' , and at most one element in $B^* - \{u, a, b\}$ that is (N, B) -robust. First consider the case where no element in $B^* - \{u, a, b\}$ is (N, B) -robust. Then there is some $q \in B^* - \{u, a, b\}$ such that $A_{pq} \neq 0$, because M' has no coloops or series pairs, and q is not (N, B) -robust. Now consider the case where there is an element $w \in B^* - \{u, a, b\}$ that is (N, B) -robust. Then (w, z, x, u, y) is a 5-element fan by Lemma 6.5, and it follows that $\{u, w\}$ is not contained in a triad. Hence, there is some $q \in B^* - \{u, w, a, b\}$ such that $A_{pq} \neq 0$ and q is not (N, B) -robust. Now, in either case, a pivot on A_{pq} is allowable, and $B' = B \Delta \{p, q\}$ is a basis of M' for which there are more (N, B') -robust elements than (N, B) -robust elements, contradicting that B is a bolstered basis.

We may now assume that there is an element q that is N -contractible but not (N, B) -robust in M' , so $q \in B^*$. Since x is in a triad with the (N, B) -strong element u , it follows that x is N -contractible in M' . If $q \in \text{cl}(\{x, y\})$, then, since $\{q, y\}$ is a parallel pair in M'/x , it follows that q is N -deletable, and hence (N, B) -robust, in M' ; a contradiction. Thus $q \notin \text{cl}(\{x, y\})$. Moreover, in the case that there is an (N, B) -robust element $z \in B$, as z is not N -deletable, it follows that $q \notin \text{cl}(\{x, y, z\})$. So $A_{pq} \neq 0$ for some element $p \in B - \{x, y\}$ that is not (N, B) -robust. Now a pivot on A_{pq} is allowable, and $B' = B \Delta \{p, q\}$ is a basis for M' such that there are more (N, B') -robust elements than (N, B) -robust elements, contradicting that B is a bolstered basis. \square

Lemma 8.2. *Suppose M has a pair of elements $\{a, b\}$ such that $M \setminus a, b$ is 3-connected with an N -minor. If (ii)(b) of Theorem 6.7 holds, and there is some $p \in (B - \{x, y\}) \cap \text{cl}(\{u, x, y\})$ such that $\{a, b\} \subseteq \text{cl}_M(\{p, x, y\})$, then $|E(M)| \leq |E(N)| + 7$.*

Proof. Let R be the set consisting of $\{p, x, y, a, b\}$ and the (N, B) -robust elements of M' outside of $\{x, y\}$. Consider the case where (ii)(b)(II) or (ii)(b)(III) of Theorem 6.7 holds. Then $\{u, x, y\}$ is contained in a (not necessarily maximal) 4-element fan (z, u, x, y) , where $z \in B$. Since $\{p, x, y, z\} \subseteq B$, but $r(\text{cl}(\{u, x, y\})) = 3$, we deduce that $p = z$. Thus $|R| \leq 7$. Towards a contradiction, suppose that $|E(M)| \geq |E(N)| + 8$. Then M has at least one element outside of R that is either N -deletable or N -contractible, but not (N, B) -robust.

Suppose first that there is some $p' \in B - \{x, y, p\}$ that is N -deletable. Then there is an element $q \in B^* - R$ such that $A_{p'q} \neq 0$, because M' is 3-connected and, in the case that Theorem 6.7(ii)(b)(III) holds, $\{z, w, u\}$ is not a triad. Since $\{a, b\} \subseteq \text{cl}_M(\{p, x, y\})$, it follows that $A_{p'a} = A_{p'b} = 0$, so a pivot on $A_{p'q}$ is allowable. But, with $B' = B \Delta \{p', q\}$, there are more (N, B') -robust elements than there are (N, B) -robust elements, contradicting that B is bolstered.

So M' has an N -contractible element $q \in B^* - R$. Suppose that $q \in \text{cl}(\{x, y, p\})$. Then, as $\{u, x, y\}$ is a triad of M' , it follows that $(\{q, u, x, y\}, p, E(M') - \{p, q, u, x, y\})$ is a vertical 3-separation of M' . But then q is N -deletable by Lemma 2.15(ii), contradicting that q is not (N, B) -robust.

Thus we may assume that $q \notin \text{cl}(\{x, y, p\})$, so there is some $p' \in B - \{x, y, p\}$ such that $A_{p'q} \neq 0$. Then $A_{p'a} = A_{p'b} = 0$, so a pivot on $A_{p'q}$ is allowable. But with $B' = B \Delta \{p', q\}$, there are more (N, B') -robust elements than there are (N, B) -robust elements, contradicting that B is bolstered. \square

We also use the following, which is proved in [3].

Lemma 8.3 ([3, Lemma 3.1]). *Let M_0 be a 3-connected matroid with $r(M_0) \geq 4$. Suppose that C^* is a rank-3 cocircuit of M_0 . If there exists some $x \in C^*$ such that $x \in \text{cl}(C^* - x)$, then $\text{co}(M_0 \setminus x)$ is 3-connected.*

We now prove our second main result, Theorem 2.31, first restating it.

Theorem 8.4. *Let M be an excluded minor for the class of \mathbb{P} -representable matroids, and let N be a non-binary 3-connected strong \mathbb{P} -stabilizer, where M has an N -minor. For some M_1 that is Δ - Y -equivalent to M , and some (M_0, N_0) in $\{(M_1, N), (M_1^*, N^*)\}$, the matroid M_0 is an excluded minor with an N_0 -minor, and at least one of the following holds:*

- (i) $|E(M_0)| \leq |E(N_0)| + 9$;
- (ii) $r(M_0) \leq r(N_0) + 7$; or
- (iii) *there is a pair $\{a, b\} \subseteq E(M)$ such that $M_0 \setminus a, b$ is 3-connected with an N_0 -minor, and $M_0 \setminus a, b$ is N_0 -fragile. Moreover, there is some bolstered basis B for M_0 and a $B \times B^*$ companion \mathbb{P} -matrix A for which $\{x, y, a, b\}$ incriminates (M, A) , where $\{x, y\} \subseteq B$, $\{a, b\} \subseteq B^*$, and both of the following hold:*

- (a) $M_0 \setminus a, b$ has at most one (N_0, B) -robust element outside of $\{x, y\}$, and
- (b) if u is an (N_0, B) -robust element of $M_0 \setminus a, b$, then $u \in B^* - \{a, b\}$, the element u is (N_0, B) -strong in $M_0 \setminus a, b$, and $\{u, x, y\}$ is a triad of $M_0 \setminus a, b$.

Proof. Suppose that neither (i) nor (ii) holds; in particular, $|E(M)| \geq |E(N)| + 10$ and $r^*(M) \geq r^*(N) + 8$. By Corollary 7.3, there exists a matroid M_0 , where M_0 is obtained from M by at most one Δ - Y or Y - Δ exchange, and $(M_1, N_1) \in \{(M_0, N), (M_0^*, N^*)\}$ such that M_1 has a pair of elements $\{a, b\}$ for which $M_1 \setminus a, b$ is 3-connected and has an N_1 -minor. By Proposition 2.28, M_1 is an excluded minor for the class of \mathbb{P} -representable matroids. We relabel (M_1, N_1) as (M, N) and apply Theorem 6.7. If (ii)(a) of Theorem 6.7 holds, then (iii) holds. We may therefore assume that (ii)(b) of Theorem 6.7 holds. Without loss of generality, we may assume that $\{b, x, y\}$ is a triangle of M .

Note that $M' = M \setminus a, b$ has an element $u \in B^*$ that is (N, B) -strong, where $\{u, x, y\}$ is a triad.

8.4.1. *The element u is N -contractible in M' .*

Subproof. As M' is not N -fragile, M' has at least one N -flexible element. If x is N -deletable, then, as u is in a series pair of $M' \setminus x$, the element u is N -contractible. Similarly, if y is N -deletable, then u is N -contractible. Thus, if the N -flexible elements of M' are contained in the triad $\{u, x, y\}$, then, since M' has at least one N -flexible element, it follows that u is N -contractible in M' . Next, suppose that (z, u, x, y) is a fan of M' , and z is N -flexible. As x is in a parallel pair of M' / z , the element x is N -deletable, so u is N -contractible. Finally, we may assume that (w, z, x, u, y) is a fan of M' , and w is N -flexible. As z is in a series pair of $M' \setminus w$, the element z is N -contractible, and it follows that x is N -deletable, so u is N -contractible. \square

Next, we show that, up to duality and replacing M by a Δ - Y -equivalent matroid, there is some deletion pair that is contained in a triangle. This triangle will provide additional leverage in later orthogonality arguments.

8.4.2. *For some $M_2 \in \{M, \nabla_T(M^*)\}$, where $T = \{b, x, y\}$, there is a pair $\{a', b'\} \subseteq E(M_2)$ such that $M_2 \setminus a', b'$ is 3-connected with an N -minor, and $\{a', b'\}$ is contained in a triangle of M_2 .*

Subproof. We first consider the case where $\{a, u\}$ is contained in a triangle with either x or y . If (ii)(b)(II) or (ii)(b)(III) of Theorem 6.7 holds, then $z \in (B - \{x, y\}) \cap \text{cl}(\{u, x, y\})$, and $\{a, b\} \subseteq \text{cl}_M(\{x, y, z\})$, so $|E(M)| \leq |E(N)| + 7$ by Lemma 8.2; a contradiction. So we may assume that (ii)(b)(I) of Theorem 6.7 holds. Now we have symmetry between x and y , so we may assume that $\{a, u, x\}$ is a triangle.

We claim that $\{b, x\}$ is a deletion pair with the desired properties. Clearly $M \setminus b$ is 3-connected and has an N -minor. By 8.4.1, u is N -contractible in $M \setminus b$. But $\{a, x\}$ is a parallel pair in $M \setminus b / u$, so $M \setminus b, x / u$, and hence $M \setminus b, x$, has an N -minor. As $\{a, u, x, y\}$ is a rank-3 cocircuit of $M \setminus b$, the matroid $\text{co}(M \setminus b, x)$ is 3-connected by Lemma 8.3. Thus, if $M \setminus b, x$ is not 3-connected, then there is a triad T^* of $M \setminus b$ that contains x . By orthogonality with the

triangle $\{a, u, x\}$, the triad T^* meets $\{a, u\}$. But $a \notin T^*$ because $M \setminus a, b$ is 3-connected. Thus T^* contains $\{x, u\}$. But since $M \setminus a, b$ is 3-connected, T^* is also a triad of $M \setminus a, b$, so $T^* \cup y$ is a 4-element cosegment of $M \setminus a, b$. Let $T^* - \{x, u\} = \{q\}$. Now $q \in B^*$, since q is N -contractible but not (N, B) -robust. But then $T^* \cup y$ is a confining set, so Proposition 4.16 implies that $|E(M)| \leq |E(N)| + 9$; a contradiction. Thus $M \setminus b, x$ is 3-connected with an N -minor, and $\{b, x\}$ is contained in a triangle of M .

We may now assume that neither $\{a, u, x\}$ nor $\{a, u, y\}$ is a triangle of M . Suppose that (ii)(b)(II) or (ii)(b)(III) of Theorem 6.7 holds. Consider the matroid $\Delta_T(M)$ obtained by a Δ - Y exchange on $T = \{b, x, y\}$. Observe that $\Delta_T(M)/b \cong M \setminus b$, where the labels on x and y are swapped. Thus, if $\Delta_T(M)/b, x \cong M \setminus b/y$ is 3-connected with an N -minor, then $\{b, x\}$ is a deletion pair of $\nabla_T(M^*)$ with the desired properties. Since y is a rim end of a maximal fan in $M \setminus a, b$, the matroid $M \setminus a, b/y$ is 3-connected by [15, Lemma 1.5]. Moreover, as $M \setminus a, b, u$ has an N -minor, and y is in a series pair in this matroid, $M \setminus b/y$ has an N -minor. If $M \setminus b/y$ is 3-connected, then $\{b, x\}$ is a deletion pair of $\nabla_T(M^*)$ as desired.

So we may assume that $M \setminus b/y$ is not 3-connected; then a is in a parallel pair of $M \setminus b/y$. Since $M \setminus b$ is 3-connected, $\{a, y, q'\}$ is a triangle of $M \setminus b$ for some $q' \in E(M) - \{a, y, u\}$. Note also that $q' \neq x$, by Lemma 8.1. If $q' \in B$, then $\{a, b\} \subseteq \text{cl}_M(\{q', x, y\})$, so (i) holds by Lemma 8.2; a contradiction. So $q' \in B^*$. Moreover, q' is N -deletable because x is N -contractible in $M \setminus b$ and q' is in a parallel pair of $M \setminus b/x$. So q' is (N, B) -robust, implying that (ii)(b)(III) holds and (q', z, x, u, y) is a maximal fan in $M \setminus a, b$. We will show that $M \setminus a, y$ is 3-connected with an N -minor. Since $\{x, y, b\}$ is a triangle and $\{x, y, u, b\}$ is a rank-3 cocircuit of $M \setminus a$, the matroid $\text{co}(M \setminus a, y)$ is 3-connected by Lemma 8.3. Suppose y is in a triad T^* of $M \setminus a$. By orthogonality, T^* meets $\{x, b\}$. But $b \notin T^*$, since $M \setminus a, b$ is 3-connected, so $x \in T^*$. Now, by orthogonality with the triangle $\{u, x, z\}$, either $T^* = \{y, x, u\}$ or $T^* = \{y, x, z\}$. Since $\{x, y, u, b\}$ is a cocircuit of $M \setminus a$, we deduce $T^* = \{y, x, z\}$. But then $\{q', z, x, y\}$ is a cosegment of $M \setminus a, b$, contradicting orthogonality with the triangle $\{z, x, u\}$. Hence $M \setminus a, y$ is 3-connected. Since $M \setminus a/x$ has an N -minor, and $\{b, y\}$ is a parallel pair in this matroid, $M \setminus a, y$ has an N -minor. So $\{a, y\}$ is a deletion pair of M that meets the requirements.

We may now assume that (ii)(b)(I) of Theorem 6.7 holds. Again, consider the matroid $\Delta_T(M)$, where $T = \{b, x, y\}$. We claim that either $\Delta_T(M)/b, x$ or $\Delta_T(M)/b, y$ is 3-connected with an N -minor, so either $\{b, x\}$ or $\{b, y\}$ is a deletion pair of $\nabla_T(M^*)$ with the desired properties. Observe that $\Delta_T(M)/b \cong M \setminus b$, so $\Delta_T(M)/b$ is 3-connected and has an N -minor. Now $\Delta_T(M)/b, x \cong M \setminus b/y$ and $\Delta_T(M)/b, y \cong M \setminus b/x$. Since u is N -deletable in M' , the elements x and y are N -contractible, so $M \setminus b/x$ and $M \setminus b/y$ have N -minors. Thus $\Delta_T(M)/b, x$ and $\Delta_T(M)/b, y$ have N -minors.

Suppose that $\text{si}(M \setminus b/x)$ is not 3-connected. Then there is a vertical 3-separation (P, x, Q) of $M \setminus b$. Recall that $\{x, y\}$ is the only series pair of $M' \setminus u$. Now, as $\text{co}(M' \setminus u) = M \setminus b/x \setminus a, u$ is 3-connected, it follows that $Q = \{a, u, q\}$ for some $q \in E(M') - \{u, x\}$, up to swapping P and Q . Since Q is 3-separating and $r(Q) \geq 3$, the set Q is a triad of $M \setminus b$. But then

$\{u, q\}$ is a series pair in $M \setminus a, b$; a contradiction. Thus $M \setminus b/x$, and hence $\Delta_T(M)/b, x$, is 3-connected up to parallel pairs. The same argument shows that $\Delta_T(M)/b, y$ is 3-connected up to parallel pairs.

Now, if $\Delta_T(M)/b, x$ or $\Delta_T(M)/b, y$ is 3-connected, then 8.4.2 holds. Thus we may assume that x and y are in triangles T_x and T_y of $M \setminus b$. If $\{a, x, y\}$ is a triangle, then $|E(M)| \leq |E(N)| + 7$ by Lemma 8.1; a contradiction. Suppose that $\{p, x, y\}$ is a triangle of M' for some $p \in E(M') - \{x, y\}$. Then p is not (N, B) -robust. Since u is N -deletable in M' , it follows that x is N -contractible in M' . Since $\{p, y\}$ is a parallel pair of M'/x , the element p is N -deletable in M' . Moreover, $p \in B^*$, since $\{x, y\} \subseteq B$ and $\{p, x, y\}$ is a triangle of M' . Therefore p is an (N, B) -robust element of M' ; a contradiction. We deduce that $\{x, y\}$ is not contained in a triangle of $M \setminus b$.

By orthogonality, T_x meets $\{a, y, u\}$, and T_y meets $\{a, x, u\}$. So either $T_x = \{x, a, q\}$ or $T_x = \{x, u, q\}$ for some $q \in E(M') - \{u, x, y\}$. Now q is N -deletable because x is N -contractible in $M \setminus b$ and q is in a parallel pair of $M \setminus b/x$. But q is not (N, B) -robust, since $q \notin \{u, x, y\}$, so $q \in B$. If $T_x = \{x, a, q\}$, then, as $q \in B - \{x, y\}$, we have $\{a, b\} \subseteq \text{cl}_M(\{q, x, y\})$, and so (i) holds by Lemma 8.2; a contradiction. So $T_x = \{x, u, q\}$. Likewise, arguing with y in the place of x , we deduce that $T_y = \{y, u, q'\}$ for some $q' \in E(M') - \{u, x, y\}$ where q' is N -deletable.

Now $T_x = \{x, u, q\}$ and $T_y = \{y, u, q'\}$ for some N -deletable elements $q, q' \in E(M') - \{u, x, y\}$. Moreover, $q \neq q'$, since $\{x, y, u\}$ is not a triangle. Since $\{q, q', u, x, y\}$ is a rank-3 set, and $\{x, y\} \subseteq B$, at most one of q and q' is in B . Without loss of generality, say $q \in B^*$. Then q is (N, B) -robust; a contradiction. \square

Let M_2 and $\{a', b'\}$ be as given in 8.4.2. We again apply Theorem 6.7, this time on the matroid M_2 with minor N and deletion pair $\{a', b'\}$; we may assume that (ii)(b) holds. We relabel M_2 as M and $\{a', b'\}$ as $\{a, b\}$. Now $M' = M \setminus a, b$ has an (N, B) -strong element $u \in B^*$, there is a 5-element cocircuit $\{x, y, u, a, b\}$ of M , and the only (N, B) -robust elements of M' are contained in a set R where $\{u, x, y\} \subseteq R$, and R is either a triad, a maximal type-II fan (z, u, x, y) relative to B , or a maximal 5-element fan (w, z, x, u, y) . Up to switching the labels on a and b , we may assume that $\{b, x, y\}$ is a triangle of M .

Additionally, now $\{a, b\}$ is contained in a triangle of M ; let $\{a, b, p\}$ be this triangle. Note that if $p \in \{x, y\}$, then $\{a, b\} \subseteq \text{cl}_M(\{x, y\})$, contradicting Lemma 8.1. So $p \notin \{x, y\}$.

8.4.3. *Let q be an N -deletable element of M' such that $q \notin \text{cl}_{M'}^*(R \cup p)$. Either*

- (I) $M \setminus b, q$ is 3-connected with an N -minor, or
- (II) there exists $t \in E(M') - q$ such that $t \notin \text{cl}_{M'}^*(R \cup p)$, the matroid $M \setminus b, t$ is 3-connected with an N -minor, $t \in B^*$, and, for some $s \in \{x, y\}$, the matroid $M \setminus b$ has a triangle $T = \{s, t, a\}$ and a 4-element cocircuit $T \cup q$.

Subproof. Note that $q \in B$, since $q \notin R$ and q is N -deletable in M' .

Suppose that $\text{co}(M \setminus b, q)$ is 3-connected, but $M \setminus b, q$ is not 3-connected. Then $M \setminus b$ has a triad $\{q, s, t\}$. Hence either $\{q, s, t\}$ or $\{b, q, s, t\}$ is a cocircuit of M . Since M' is 3-connected, it follows that $a \notin \{q, s, t\}$ and that $\{q, s, t\}$ is a triad of M' . As q is N -deletable in M' , the elements s and t are N -contractible in M' . We claim that $(R - \{x, y\}) \cap \{s, t\} = \emptyset$. To begin with, $u \notin \{s, t\}$ since $\{x, y\}$ is the only series pair of $M' \setminus u$. If R is a maximal type-II fan (z, u, x, y) of M' , then $z \notin \{s, t\}$ since the fan is maximal. Finally, if R is a 5-element fan (w, z, x, u, y) of M' with $w \in \{s, t\}$, then w is N -deletable, implying q is N -contractible and hence N -flexible in M' ; a contradiction. Thus, as s and t are N -contractible but not in $R - \{x, y\}$, either $\{s, t\} \subseteq B^* - \{a, b, u\}$, or $\{s, t\}$ meets $\{x, y\}$.

Suppose that $\{s, t\} \subseteq B^* - \{a, b, u\}$. If $\{q, s, t\}$ is a triad of M , then $A_{qa} = A_{qb} = 0$, so there is an allowable pivot on A_{qs} or A_{qt} that gives a basis for M' with more robust elements, contradicting that B is a bolstered basis. On the other hand, if $\{b, q, s, t\}$ is a cocircuit of M , then it intersects the triangle $\{b, x, y\}$ in a single element; a contradiction to orthogonality.

Therefore $\{s, t\}$ meets $\{x, y\}$. If $\{s, t\} = \{x, y\}$, then $q \in \text{cl}_{M'}^*(\{x, y\})$; a contradiction. So we may assume that $s \in \{x, y\}$ and $t \notin \{x, y\}$. If $\{q, s, t\}$ is a triad of M , then this triad intersects $\{b, x, y\}$ in a single element; a contradiction. On the other hand, if $\{q, s, t, b\}$ is a cocircuit of M , then by orthogonality with the triangle $\{a, b, p\}$, we have $t = p$, in which case $q \in \text{cl}_{M'}^*(\{x, y, p\})$; a contradiction.

We may now assume that $\text{co}(M \setminus b, q)$ is not 3-connected. We first show $\text{co}(M \setminus a, b, q)$ is 3-connected. Suppose not. Then there is a cyclic 3-separation (X, q, Y) of M' such that $|X \cap E(N)| \leq 1$ and $Y \cup q$ is coclosed in M' . By the dual of Lemma 2.15, at most one element of X is not N -flexible in M' , and if such an element v exists, then $q \in \text{cl}_{M'}^*(X - v)$. But $X - v \subseteq R$, so $q \in \text{cl}_{M'}^*(R)$; a contradiction. So $\text{co}(M \setminus a, b, q)$ is 3-connected.

Since $\text{co}(M \setminus b, q)$ is not 3-connected, there is a cyclic 3-separation (P, q, Q) of $M \setminus b$ with $a \in Q$. Since $\text{co}(M \setminus a, b, q)$ is 3-connected, $(P, Q - a)$ is not a cyclic 2-separation of $M \setminus a, b, q$, so $Q - a$ is a series class of $M \setminus a, b, q$. Hence $(Q - a) \cup q$ is a cosegment of M' . Suppose that $|Q - a| \geq 3$. Then $Q - a$ meets B , and, since q is N -deletable in M' , the elements of $Q - a$ are N -contractible. By the dual of Lemma 2.2, the elements of $(Q - a) \cap B$ are (N, B) -strong, so Lemma 3.1 implies that $(Q - a) \cap B \subseteq \{x, y\}$. Since q is not cospanned by $\{x, y\}$ in M' , we have $|(Q - a) \cap B| = 1$, and thus $|Q - a| = 3$. But then $\{x, y, u\} \cup (Q - a)$ is a corank-3 confining set of $M \setminus a, b$, contradicting Proposition 4.16. Therefore $|Q - a| = 2$. Since Q is a 3-separating set of $M \setminus b$ that contains a circuit, $Q = \{s, t, a\}$ is a triangle.

Since M' is 3-connected, either $\{q, s, t\}$ or $\{q, s, t, a\}$ is a cocircuit of $M \setminus b$. By orthogonality between the triangle $\{s, t, a\}$ and the cocircuit $\{x, y, u, a\}$ of $M \setminus b$, we have that $\{x, y, u\}$ meets $\{s, t\}$. Moreover, $u \notin \{s, t\}$ because the only triad containing u in M' is $\{u, x, y\}$. Thus $\{x, y\}$ meets $\{s, t\}$. However, $\{x, y\} \neq \{s, t\}$, otherwise $\{x, y\}$ spans $\{a, b\}$, contradicting Lemma 8.1. Without loss of generality, let $s \in \{x, y\}$ and $t \notin \{x, y\}$.

Suppose $\{q, s, t\}$ is a triad of $M \setminus b$. If $\{q, s, t\}$ is a triad of M , then, by orthogonality with the triangle $\{b, x, y\}$, we have $t \in \{x, y\}$; a contradiction. On the other hand, if $\{q, s, t, b\}$ is a triad of M , then, by orthogonality,

the triangle $\{a, b, p\}$ meets $\{s, t\}$. But then $p = t$, so $q \in \text{cl}_{M'}^*(\{x, y, p\})$; a contradiction. So $\{q, s, t, a\}$ is a cocircuit of $M \setminus b$.

We claim that t satisfies (II). Recall $s \in \{x, y\}$, and pick s' such that $\{s, s'\} = \{x, y\}$. Since s is N -contractible in $M \setminus b$ and $\{a, t\}$ is a parallel pair of $M \setminus b/s$, it follows that $M \setminus b, t$ has an N -minor. If $t \in \text{cl}_{M'}^*(R \cup p)$, then $q \in \text{cl}_{M'}^*(R \cup p)$, since $s \in R$. Thus $t \notin \text{cl}_{M'}^*(R \cup p)$. Now, t is N -contractible in M' , since t is in a series pair in $M' \setminus q$. Thus, if $t \in B$, then t is (N, B) -robust, and $q \in \text{cl}_{M'}^*(R)$; a contradiction. So $t \in B^*$.

It remains to prove that $M \setminus b, t$ is 3-connected. Since $\{q, s, t, a\}$ is a rank-3 cocircuit in $M \setminus b$, the matroid $\text{co}(M \setminus b, t)$ is 3-connected by Lemma 8.3. Suppose $M \setminus b, t$ is not 3-connected. Then t is in a triad T^* of $M \setminus b$. By orthogonality with the triangle $\{s, t, a\}$, the triad T^* meets $\{s, a\}$. But $a \notin T^*$ because $M \setminus a, b$ is 3-connected. Thus $\{s, t\} \subseteq T^*$. If $p \in T^*$, then $t \in \text{cl}_{M \setminus b}^*(\{s, p\})$, so $t \in \text{cl}_{M'}^*(\{x, y, p\})$; a contradiction. So $p \notin T^*$. Now T^* or $T^* \cup b$ is a cocircuit of M . But $\{b, x, y\}$ is a triangle of M that meets T^* in a single element, so T^* is not a cocircuit; and $\{a, b, p\}$ is a triangle of M that meets $T^* \cup b$ in a single element, so $T^* \cup b$ is not a cocircuit. We deduce that $M \setminus b, t$ is 3-connected, and 8.4.3 follows. \square

8.4.4. *There are distinct elements $q', q'' \in E(M)$ such that, for $q \in \{q', q''\}$, both of the following hold:*

- (I) $M \setminus b, q$ is 3-connected with an N -minor, where $q \notin \text{cl}_{M'}^*(R \cup p)$; and
- (II) either
 - (a) $q \in B$, the element q is N -deletable in $M \setminus a, b$, and neither $\{x, u, q\}$ nor $\{y, u, q\}$ is a triangle; or
 - (b) $q \in B^*$ and, for some $s \in \{x, y\}$, the set $T = \{s, q, a\}$ is a triangle that is contained in a 4-element cocircuit of $M \setminus b$.

Subproof. Suppose R is either a 4- or 5-element fan. Then $r_{M'}^*(R) = 3$, the set $\{x, u\}$ is contained in a triangle with an element $z \in R$, and $\{y, u\}$ is not in a triangle, since R is a maximal fan. Now $r_M^*(R \cup \{a, b, p\}) \leq 6$. Since $r^*(M) \geq r^*(N) + 8$, there are distinct N -deletable elements q', q'' outside of $\text{cl}_M^*(R \cup \{a, b, p\})$, neither of which is in a triangle with $\{x, u\}$ or $\{y, u\}$.

Now suppose $R = \{u, x, y\}$. Then $r_{M'}^*(R) = 2$, so $r_M^*(R \cup \{a, b, p\}) \leq 5$. Since $r^*(M) \geq r^*(N) + 8$, there are at least three N -deletable elements outside of $\text{cl}_M^*(R \cup \{a, b, p\})$. Since these N -deletable elements are not (N, B) -robust, they belong to $B - \{x, y\}$. As $r_{M'}(\{x, y, u\}) = 3$, and $\{x, y\} \subseteq B$, at most one of these elements is in a triangle with $\{x, u\}$ or $\{y, u\}$. Thus there exist distinct elements q', q'' outside of $\text{cl}_M^*(R \cup \{a, b, p\})$, neither of which is in a triangle with $\{x, u\}$ or $\{y, u\}$.

Now $q', q'' \notin \text{cl}_M^*(R \cup \{a, b, p\}) = \text{cl}_{M'}^*(R \cup p)$. By 8.4.3, either q' satisfies (I) and (II)(a), or there exists an element t' that satisfies (I) and (II)(b). Likewise, either q'' satisfies (I) and (II)(a), or there exists an element t'' that satisfies (I) and (II)(b). Suppose that neither q' nor q'' satisfies (II)(a), and $t' = t''$. Then $\{s', t', a\}$ and $\{s'', t', a\}$ are triangles of $M \setminus b$ where $s', s'' \in \{x, y\}$. If $\{s', s''\} = \{x, y\}$, then $\{x, y, t'\}$ is a triangle of M' , but then $\text{co}(M' \setminus u) \cong M' \setminus u/x$ is not 3-connected; a contradiction. So $s' = s''$. Now $\{q', s', t', a\}$ and $\{q'', s', t', a\}$ are distinct cocircuits of $M \setminus b$, so $\{q', q'', s', t'\}$

is a cosegment of M' . But q' is N -deletable in M' , implying q'' is N -contractible and hence (N, B) -robust; a contradiction. \square

Let q' and q'' be elements as in 8.4.4. Suppose that $\{q', q''\} \subseteq B^*$. Then, by 8.4.4(II)(b), $M \setminus b$ has a triangle $T' = \{s', q', a\}$ that is contained in a 4-element cocircuit C^* , and a triangle $T'' = \{s'', q'', a\}$, for some $s', s'' \in \{x, y\}$. Observe that $\{x, y\} \not\subseteq C^*$, since $q' \notin \text{cl}_{M'}^*(\{x, y\})$ by 8.4.4(I). Thus $C^* \cup b$ is a cocircuit of M , by orthogonality with the triangle $\{b, x, y\}$. If $s' = s''$, then $\{q', q'', a\}$ is a triangle that intersects the cocircuit $\{x, y, u, a, b\}$ in a single element; a contradiction. Thus we may assume that $T' = \{x, q', a\}$ and $T'' = \{y, q'', a\}$. By orthogonality between $C^* \cup b$ and T'' , we deduce that $q'' \in C^* \cup b$, since $y \notin C^*$. Now $\{x, q', q''\}$ is a triad of M' with $\{q', q''\} \subseteq B^*$, so $\{u, x, y, q', q''\}$ is a corank-3 confining set, contradicting Proposition 4.16.

Without loss of generality, we may now assume that $q' \in B$ and q' is N -deletable in M' . Towards a contradiction, assume that (iii) does not hold for M and the deletion pair $\{b, q'\}$. Then, after applying Theorem 6.7, (ii)(b) holds. Let A' be the $B' \times (B')^*$ companion \mathbb{P} -matrix where $\{x', y', b, q'\}$ incriminates (M, A') for $\{x', y'\} \subseteq B'$ and $\{b, q'\} \subseteq (B')^*$. Then M has a 5-element cocircuit $D' = \{x', y', u', b, q'\}$, where $M \setminus b, q'$ has an (N, B') -strong element u' outside of $\{x', y'\}$, and either $\{b, x', y'\}$ or $\{q', x', y'\}$ is a triangle.

Suppose that $\{b, x', y'\}$ is a triangle of M . By orthogonality between the cocircuit D' of M and the triangles $\{b, x, y\}$ and $\{a, b, p\}$, and using the fact that $q' \notin \{x, y, a, p\}$, we deduce that $\{x, y\}$ and $\{a, p\}$ meet $\{x', y', u'\}$. If $\{x, y\}$ or $\{a, p\}$ intersects $\{x', y'\}$ in a single element, then $\{b, x, y\}$ or $\{a, b, p\}$ is in the span of $\{x', y'\}$, so $\{x', y'\}$ spans a 4-element segment in M . Thus $\{x', y'\}$ spans a triangle in $M \setminus b, q'$. But then $\text{co}(M \setminus b, q', u')$ is not 3-connected by Lemma 2.11, contradicting that u' is (N, B') -strong in $M \setminus b, q'$. We deduce that $\{x', y', u'\} \subseteq \{x, y, a, p\}$. But $q' \in \text{cl}_{M'}^*(\{x', y', u'\}) \subseteq \text{cl}_{M'}^*(\{x, y, p\})$, contradicting 8.4.4(I).

We may now assume that $\{q', x', y'\}$ is a triangle of M . The triangles $\{b, x, y\}$ and $\{a, b, p\}$ meet the cocircuit D' in the element b . Thus, by orthogonality, $\{x, y\}$ and $\{a, p\}$ meet $\{x', y', u'\}$. Let $D = \{x, y, u, a, b\}$, and recall that D is a cocircuit of M .

First suppose that $D \cap \{x', y'\} = \emptyset$. Then $u' \in \{x, y\}$ and $p \notin D$, so $p \in \{x', y'\}$. Since q' is N -deletable in M' , the element a is N -deletable in $M \setminus b, q'$. If $a \in B'$, then $\{a, b, p\}$ is a triangle of M with $a, p \in B'$, so $A'_{vb} = 0$ for $v \in \{x', y'\} - p$, contradicting that the bad submatrix $A'[\{x', y', b, q'\}]$ has no zero entries. So $a \in (B')^*$, hence a is (N, B') -robust in $M \setminus b, q'$. It follows that $M \setminus b, q'$ has a 5-element fan (a, z', x', u', y') for some z' , where the (N, B') -robust elements of $M \setminus b, q'$ are contained in $\{x', y', u', z', a\}$. Since $M \setminus b, q', a$ has an N -minor, and x' is in a series pair in this matroid, the element x' is N -contractible in $M \setminus a, b$. Moreover, $\{u', z'\}$ and $\{q', y'\}$ are parallel pairs in $M \setminus a, b/x'$, so $M \setminus a, b, u', y'$ has an N -minor. But $\{x', q'\}$ is a series pair in this matroid, so q' is also N -contractible in $M \setminus a, b$. Now q' is (N, B) -robust; a contradiction.

Now we may assume that $D \cap \{x', y'\} \neq \emptyset$. By orthogonality with the triangle $\{q', x', y'\}$, we have $\{x', y'\} \subseteq D$. If $u' \in D$, then $q' \in \text{cl}_M^*(D) \subseteq \text{cl}_M^*(R)$; a contradiction. By orthogonality between the cocircuit D' and triangles $\{b, x, y\}$ and $\{a, b, p\}$, one of $\{x', y'\}$ is in $\{x, y\}$ and the other in

$\{a, p\} \cap D$. By 8.4.4(II)(a), neither $\{x, u, q'\}$ nor $\{y, u, q'\}$ is a triangle, so $\{s, a, q'\}$ is a triangle for some $s \in \{x, y\}$. But $\{s, q'\} \subseteq B$, so either $A_{xa} = 0$ or $A_{ya} = 0$, contradicting that the bad submatrix has no zero entries. We deduce that (iii) holds for M and the pair $\{b, q'\}$. \square

ACKNOWLEDGEMENTS

We thank the anonymous referees of an earlier version of this manuscript, whose valuable suggestions lead to vast improvements in this version.

REFERENCES

- [1] BIXBY, R. E. On Reid's characterization of the ternary matroids. *Journal of Combinatorial Theory, Series B* 26, 2 (1979), 174–204.
- [2] BIXBY, R. E. A simple theorem on 3-connectivity. *Linear Algebra and its Applications* 45 (1982), 123–126.
- [3] BRETTELL, N., AND SEMPLE, C. A splitter theorem relative to a fixed basis. *Annals of Combinatorics* 18, 1 (2014), 1–20.
- [4] BRETTELL, N., WHITTLE, G., AND WILLIAMS, A. N -detachable pairs in 3-connected matroids I: unveiling X . Preprint, arXiv:1804.05637, 2018.
- [5] BRETTELL, N., WHITTLE, G., AND WILLIAMS, A. N -detachable pairs in 3-connected matroids II: life in X . Preprint, arXiv:1804.06029, 2018.
- [6] BRETTELL, N., WHITTLE, G., AND WILLIAMS, A. N -detachable pairs in 3-connected matroids III: the theorem. Preprint, arXiv:1804.06588, 2018.
- [7] CLARK, B., MAYHEW, D., VAN ZWAM, S. H. M., AND WHITTLE, G. The structure of $\{U_{2,5}, U_{3,5}\}$ -fragile matroids. *SIAM Journal on Discrete Mathematics* 30, 3 (2016), 1480–1508.
- [8] GEELEN, J. F., GERARDS, A. M. H., AND KAPOOR, A. The excluded minors for $GF(4)$ -representable matroids. *Journal of Combinatorial Theory, Series B* 79, 2 (2000), 247–299.
- [9] HALL, R., MAYHEW, D., AND VAN ZWAM, S. H. M. The excluded minors for near-regular matroids. *European Journal of Combinatorics* (2011).
- [10] MAYHEW, D., AND ROYLE, G. F. Matroids with nine elements. *Journal of Combinatorial Theory, Series B* 98, 2 (2008), 415–431.
- [11] MAYHEW, D., VAN ZWAM, S. H. M., AND WHITTLE, G. Stability, fragility, and Rota's Conjecture. *Journal of Combinatorial Theory, Series B* 102, 3 (2012), 760–783.
- [12] OXLEY, J. *Matroid theory, Second Edition*. Oxford University Press, New York, 2011.
- [13] OXLEY, J., SEMPLE, C., AND VERTIGAN, D. Generalized Delta-Y Exchange and k -Regular Matroids. *Journal of Combinatorial Theory, Series B* 79, 1 (2000), 1–65.
- [14] OXLEY, J., SEMPLE, C., AND WHITTLE, G. Maintaining 3-connectivity relative to a fixed basis. *Advances in Applied Mathematics* 41, 1 (2008), 1–9.
- [15] OXLEY, J., AND WU, H. On the structure of 3-connected matroids and graphs. *European Journal of Combinatorics* 21, 5 (2000), 667–688.
- [16] PENDAVINGH, R. A., AND VAN ZWAM, S. H. M. Confinement of matroid representations to subsets of partial fields. *Journal of Combinatorial Theory, Series B* 100, 6 (2010), 510–545.
- [17] PENDAVINGH, R. A., AND VAN ZWAM, S. H. M. Lifts of matroid representations over partial fields. *Journal of Combinatorial Theory, Series B* 100, 1 (2010), 36–67.
- [18] SEMPLE, C., AND WHITTLE, G. Partial fields and matroid representation. *Advances in Applied Mathematics* 17, 2 (1996), 184–208.
- [19] SEMPLE, C. A. *k -regular matroids*. PhD thesis, Victoria University of Wellington, 1998.
- [20] SEYMOUR, P. D. Matroid representation over $GF(3)$. *Journal of Combinatorial Theory, Series B* 26, 2 (1979), 159–173.
- [21] TUTTE, W. T. A homotopy theorem for matroids. I, II. *Transactions of the American Mathematical Society* 88, 1 (1958), 144–174.

- [22] WHITTLE, G. On matroids representable over $GF(3)$ and other fields. *Transactions of the American Mathematical Society* 349, 2 (1997), 579–603.
- [23] WHITTLE, G. Stabilizers of classes of representable matroids. *Journal of Combinatorial Theory, Series B* 77, 1 (1999), 39–72.
- [24] WHITTLE, G., AND WILLIAMS, A. On preserving matroid 3-connectivity relative to a fixed basis. *European Journal of Combinatorics* 34, 6 (2013), 957–967.
- [25] YU, Y. More forbidden minors for wye-delta-wye reducibility. *The Electronic Journal of Combinatorics* 13, 1 (2006), 7.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, EINDHOVEN UNIVERSITY OF TECHNOLOGY, THE NETHERLANDS

E-mail address: `nbrettell@gmail.com`

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA, USA

E-mail address: `bclark@lsu.edu`

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA, USA

E-mail address: `oxley@math.lsu.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF CANTERBURY, NEW ZEALAND

E-mail address: `charles.semple@canterbury.ac.nz`

SCHOOL OF MATHEMATICS AND STATISTICS, VICTORIA UNIVERSITY OF WELLINGTON, NEW ZEALAND

E-mail address: `geoff.whittle@vuw.ac.nz`