

2013

Epidemiology and molecular characterization of human and canine hookworm

Ntombi B. Mudenda

Louisiana State University and Agricultural and Mechanical College

Follow this and additional works at: https://digitalcommons.lsu.edu/gradschool_dissertations



Part of the [Veterinary Pathology and Pathobiology Commons](#)

Recommended Citation

Mudenda, Ntombi B., "Epidemiology and molecular characterization of human and canine hookworm" (2013). *LSU Doctoral Dissertations*. 1209.

https://digitalcommons.lsu.edu/gradschool_dissertations/1209

This Dissertation is brought to you for free and open access by the Graduate School at LSU Digital Commons. It has been accepted for inclusion in LSU Doctoral Dissertations by an authorized graduate school editor of LSU Digital Commons. For more information, please contact gradetd@lsu.edu.

EPIDEMIOLOGY AND MOLECULAR CHARACTERIZATION OF HUMAN AND CANINE HOOKWORM

A Dissertation

Submitted to the Graduate Faculty of the
Louisiana State University and
Agricultural and Mechanical College
in partial fulfillment of the
requirements for the degree of
Doctor of Philosophy

in

Veterinary Medical Sciences

by

Ntombi B. Mudenda
BVM, University of Zambia, 2002
MS, Royal Veterinary College, 2004
December 2013

Dedicated to my sons,
Penjani, Tabiso and Aiden

ACKNOWLEDGEMENTS

I would like to thank my mentor, Dr. John B. Malone for the guidance, encouragement and understanding. Many thanks to my graduate committee members, Dr. James Miller and Dr. Christopher Mores for their invaluable contribution to this work and for allowing me to use resources from their labs. I am also grateful to my Dean's Rep, Dr. Lewis Gaston for being available and for his critic.

Thanks to Dr. Wendy Wolfson, the staff and students in the shelter medicine program/rotation for the sample collection and the fruitful discussions we had. I am also grateful to the animal shelter managers for agreeing to be a part of this work.

Special thanks to our collaborator at Case Western Reserve University, Dr. Ronald Blanton for helping me prepare for and organize the work in Brazil. I am very grateful to Dr. Mitermayer Reis and his laboratory staff at Oswaldo Cruz (FIOCRUZ), Salvador, Brazil for allowing me to work in their laboratory and facilitating my field work. Special mention goes to Dr. Jackson Costa for helping me organize the field work in Mutuípe, Bahia and Manuela Costa for being my very able Portuguese translator. I am grateful to the Mutuípe municipality Health Secretary for permission to work in the Central Laboratory and for the great welcome and pleasant work atmosphere from the staff in the lab. I am also grateful to Dr. Mariah Bavia for the facilitating my stay in Brazil.

Thanks to Dr. Dwight Bowman and Janice Liotta for the donation of *A. braziliense* eggs and DNA which allowed me to complete my molecular work on canine hookworm.

I am forever grateful to Dan Chisenhall for patiently teaching me PCR and giving me a better understanding of molecular techniques, for always being available to discuss my

work and providing guidance. I am also grateful to Mr Michael Kearney for the guidance with statistics.

To my fellow lab mates, past and present, Brooke Delcambre, Moara Martins, Cassan Pulaski, Ryan Avery, Paula Mischler and Patricia Smith (Miss Pat), I say thank you all for your input into my work and for creating a pleasant work environment. I also thank my fellow graduate students for willingly helping me whenever I needed it, both academically and otherwise. I could not have asked for better colleagues.

This dissertation would not have been possible without funding from Pan American Health Organization, the LSU-SVM CORP grant as well as my personal sponsorship from the Fulbright Foreign Student grant and for that I am grateful.

Last but definitely not least, I am eternally grateful to my supportive husband, Chew Nkonde, my patient children, my mother for providing the American experience to our boys, my siblings, father, my whole family and friends across the globe for the encouragement, support and prayers. May God bless you all.

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	iii
LIST OF TABLES	vii
LIST OF FIGURES	ix
ABSTRACT	xi
1 BACKGROUND	1
1.1 Introduction	1
1.2 Ecological Niche Modeling	1
1.3 Soil-Transmitted Helminths	5
1.3.1 Biology of STHs	5
1.3.2 Public Health Significance.....	11
1.3.3 Prevalence	13
1.3.4 Diagnosis.....	14
1.3.5 Treatment and Control	19
1.4 Human Hookworm Infection	23
1.4.1 Hookworm in Brazil	25
1.5 Common Intestinal Parasites of Dogs.....	26
1.5.1 Biology and infection.....	27
1.5.2 Epidemiology of Canine Intestinal Parasites.	35
1.5.3 Diagnosis of Canine Gastrointestinal Parasites	38
1.5.4 Treatment and control	40
1.6 References	41
2 MODELING THE ECOLOGICAL NICHE OF HOOKWORM IN BRAZIL BASED ON CLIMATE	59
2.1 Introduction	59
2.2 Materials and Methods	62
2.2.1 GDD	63
2.2.2 Moisture Adjusted GDD Model (MA-GDD)	64
2.2.3 GDD-Water Budget Gradient Model (GDD X WB)	64
2.2.4 Calculation of Mean Transmission Time and Potential Transmission Cycles per Year	64
2.2.5 Model Validation	65
2.2.6 Statistical Analysis.....	65
2.3 Results	66
2.4 Discussion	70
2.4.1 MA-GDD Index and Transmission Cycles.....	71
2.4.2 GDDxWB Index	73
2.4.3 Data Validation	74

2.4.4 Other Environmental Factors	75
2.5 References	76
3 HOOKWORM AND OTHER SOIL TRANSMITTED HELMINTHS AMONG PATIENTS IN MUTUÍPE, BRAZIL	83
3.1 Introduction	83
3.2 Materials and Methods	86
3.2.1 Study Area	86
3.2.2 Parasitological Survey	86
3.2.3 Larvae Culture and Examination	87
3.2.4 Preparation and Storage of Fecal Samples	88
3.2.5 DNA Extraction from Infective Larvae and Fecal Samples	89
3.2.6 PCR for <i>N. americanus</i> and <i>A. duodenale</i> Differentiation	89
3.3 Results	91
3.3.1 Hookworm Infection	93
3.4 Discussion	97
3.4.1 Conclusions	100
3.5 References	100
4 GASTROINTESTINAL PARASITE SURVEY OF SHELTER DOGS IN SOUTHERN LOUISIANA	106
4.1 Introduction	106
4.2 Methods	107
4.2.1 Parasitological Examination	108
4.2.2 PCR and Restriction Fragment Length Polymorphism (RFLP) for Hookworm Speciation	109
4.3 Results	111
4.3.1 Distribution of Parasites by Parish	111
4.3.2 Effect of Age on Prevalence	112
4.3.3 Effect of Sex on Prevalence	116
4.3.4 Deworming	116
4.3.5 Speciation of Hookworm by PCR	119
4.4 Discussion	122
4.4.1 Conclusion	126
4.5 References	127
APPENDIX	
A	129
B	132
C	133
THE VITA	134

LIST OF TABLES

1.1	Morphological differences between <i>A. duodenale</i> and <i>N. americanus</i>	11
1.2	Classes of intensity of infection	20
1.3	Prevalence of common dog intestinal parasites from national and regional surveys in shelter and pet dogs	37
2.1	Distribution of hookworm prevalence among the 975 municipalities surveyed with corresponding MA-GDD and GDDxWB values	66
2.2	Important biological and climatic variables for the transmission of <i>N. americanus</i> used as baseline values for calculations based on the GDD-WB concept	73
3.1	Prevalence of parasites in out-patients from health posts of Mutuípe municipality	91
3.2	STH egg detection by number of fecal samples submitted by patients.....	92
3.3	Occurrence of single and multi-parasite infections of STH in out-patients.....	92
3.4	Fecal egg counts of STH in patients.....	92
3.5	Morphological characteristics of hookworm infective larvae in Mutuípe.....	96
4.1	Prevalence of parasites found in dogs from shelters in nine parishes of south Louisiana	114
4.2	Parasite prevalence in dogs less than 6 months old and more than 6 months old	116
4.3	Occurrence of single species or combinations of parasite species found in dogs over 6 months and less than 6 months of age	117
4.4	Parasite prevalence in male and female dogs	118
4.5	Parasitism in shelter dogs based on history of deworming.....	118
4.6	Prevalence of parasites in dogs with a history of anthelmintic treatment at shelters	119
4.7	Occurrence of hookworm and <i>Trichuris</i> in dogs treated with pyrantel	119

4.8	Comparison of PCR-RFLP protocols for speciation of <i>A. caninum</i> from 88 samples by size of PCR amplicon and number of fragments generated from endonuclease digestion	120
-----	--	-----

LIST OF FIGURES

1.1	Lifecycle of <i>A. lumbricoides</i>	7
1.2	Lifecycle of <i>T. trichiura</i>	8
1.3	Hookworm lifecycle.....	10
1.4	Global distribution of Soil-transmitted helminths	14
1.5	Schematic representation of part of the rDNA transcriptional unit and relative locations of primers NC5, NC1 and NC2 commonly used for hookworm species differential diagnosis	19
1.6	Climatic suitability for hookworm transmission defined by land surface temperature and aridity	24
1.7	Zoonotic Hookworms.....	28
1.8	<i>Toxocara canis</i> life cycle.....	30
2.1	Risk Model for Hookworm using Moisture Adjusted GDD.....	68
2.2	Hookworm prevalence risk model using GDDxWB index.....	69
2.3	Scatter plot of hookworm prevalence from municipality data points against transmission cycles per year	69
2.4	Scatter plot of Hookworm Prevalence against GDD-WB index.....	70
3.1	Harada-Mori culture.....	88
3.2	Proportions of parasitized and non-parasitized patients in urban and rural Mutuípe.....	93
3.3	Prevalence (A) and Mean hookworm EPG (B) among age groups	94
3.4	Distribution of hookworm positive patients among health posts	95
3.5	<i>N. americanus</i> infective larvae	96
4.1	Map of Louisiana with the location of parishes in the study area	108
4.2	PCR-RFLP results for the <i>A. braziliense</i> using primer sets NC1/NC2 and NC5/NC2	121

4.3	PCR-RFLP results for the <i>A. caninum</i> using primer sets NC1/NC2 and NC5/NC2.....	121
4.4	PCR-RFLP results for <i>A. braziliense</i> showing non-specific amplification	122

ABSTRACT

Among the soil-transmitted helminths (STH), hookworms are a worldwide problem in both humans and animals. They cause non-specific gastrointestinal symptoms, and in young children and animals, they can cause stunting, malnutrition and anemia. Canine hookworms have significant zoonotic potential as a cause of cutaneous larvae migrans and eosinophilic enteritis in humans.

To determine the ecological niche of human hookworm in Brazil, two risk models were developed based on the Growing Degree Day-Water Budget (GDD-WB) concept, one based on accumulation of monthly temperatures above a base temperature of 15°C and threshold WB value >0.4. The second was based on a 'gradient index' of the product of monthly accumulated GDD and WB values. It was determined that both environmental temperature and moisture are important in the distribution of hookworm. This study supports the validity of the GDD-WB concept for mapping risk of hookworm at a national scale.

A cross-sectional survey was conducted in human outpatients in Mutuípe municipality, Brazil, to determine prevalence of soil-transmitted helminths, including hookworm. Mutuípe falls within the permissive zone for the transmission of hookworm. A sucrose double centrifugation flotation technique was used for the concentration of helminth eggs in fecal samples. Hookworm infection was the most prevalent of the STH and the prevalence was highest in adults and males. PCR was then used to determine the species of hookworm present. *Necator americanus* was confirmed by PCR as the predominant hookworm species. A single case of *Ancylostoma ceylanicum* was identified.

A study on the prevalence of hookworms and other gastrointestinal parasites in shelter dogs in south Louisiana and the anthelmintic protocols used in the shelters was conducted. Fecal samples examined by direct smear, flotation and sedimentation methods revealed that hookworm had the highest prevalence (53.6%) followed by *Trichuris vulpis* (28.7), *Cystoisospora ohioensis* (17.2%), *Giardia duodenalis* (12.0%), *C. canis* (7.7), *Toxocara canis* (6.2%), *Dipylidium caninum*, *Alaria* spp and *Capillaria* sp. A PCR-RFLP developed to differentiate *A. caninum* and *A. braziliense* revealed *A. caninum* as the only species found. Evaluation of the anthelmintic protocols used in nine shelters showed current methods were inadequate for control of hookworms in shelter dogs.

1 BACKGROUND

1.1 Introduction

Hookworm occurs worldwide and usually concurrently with other parasites. Hookworm was the focus of this dissertation; however, in surveying populations for hookworm, it is important to describe the occurrence of other parasites as the morbidity associated is confounded by co-infections. Hookworm surveys normally involve examination of fecal samples for eggs microscopically and sometimes the more modern molecular methods. Spatial modeling is useful as a tool in planning surveys as it provides some insight into the possible occurrence of parasites in hard to reach areas or where surveys have never been done. Canine hookworm in Louisiana was studied in preparation for work with human hookworm in Brazil and for its zoonotic capability. This chapter describes background information on ecological niche modeling, human and canine hookworm as well as parasites that normally occur with hookworm.

1.2 Ecological Niche Modeling

A geographic information system (GIS) is a computerized system which is used to capture, store, manipulate and analyze georeferenced data (Bergquist and Rinaldi, 2010; Jones, 2012). It is widely used in environmental epidemiology and public health as it enables a visual presentation of results and disease forecasts (Jones, 2012). Remote sensing is defined as obtaining information about an object without physical contact with it (Campbell and Wynne, 2011). Earth orbiting satellites are used in remote sensing to collect information about the earth's surface by electromagnetic radiation reflected or emitted by the earth's surface. The Landsat satellite program was launched in 1972 by the US government to capture satellite images of the earth and continues to provide

information about the surface of the earth with a spatial resolution of 15-60m (Campbell and Wynne, 2011). The Moderate-resolution Imaging Spectroradiometer (MODIS) is another sensor that orbits the earth onboard the Terra and Aqua Earth Orbiting System (EOS) satellites and captures data at 250m to 1km spatial resolutions. Other satellite sensors, often used in geospatial health applications include the Advanced Very- High Resolution Radiometer (AVHRR) and Advanced Space-borne Thermal Emission and reflection Radiometer (ASTER) (Campbell and Wynne, 2011). These sensors provide data on land and ocean surface temperature, snow coverage, land use, land cover, elevation, topography, soil moisture, rainfall, and other environmental features.

Normalized Difference Vegetative Index (NDVI) is a measure of the greenness or chlorophyll activity of vegetation. It is derived from the red and near infrared radiation readings corresponding to different levels of vegetation cover. Surface temperature is influenced by the presence of vegetation, as plants absorb visible red light during photosynthesis and reflect solar radiation in the near-infrared spectrum which results in a high NDVI value. NDVI is calculated as follows:

$$NDVI = \frac{NIR - R}{NIR + R}$$

Where *NIR* is *near infrared radiation* and *R* is in the *red* spectral range (Campbell and Wynne, 2011).

GIS provides the ability to bring remotely sensed and other geospatial data into one analytical framework, thereby enhancing the range of analytical products (Campbell and

Wynne, 2011). These products include mapping of urban infrastructure, floodplain mapping, and disease distribution mapping.

Maps for human and animal parasites and as well as prediction/risk maps for the diseases they cause have been developed in many regions of the world. This has been possible using the biological/ecological niche concept for parasites that have part or all their life cycle affected by environmental conditions. Environmental properties that contribute to the ecological niche of a particular parasite or vector have to be determined in order to be able to predict the distribution of the organism (Malone, 2005). In pursuing ecological analysis, the following steps should be followed (Bergquist and Rinaldi, 2010):

- Construction of a GIS for the area of interest using data layers delineating environmental and climate features;
- Construction of a frame of georeferenced areas;
- Construction of a database with environmental and parasitological data;
- Statistical analysis to detect environmental risk factors and/or develop prediction models.

Risk maps based on known biological requirements and climatic variables have been employed for a number of diseases including malaria (Thomson et al., 1997; Yang et al., 2010), schistosomiasis (Lindsay and Thomas, 1999; Malone et al., 2001), filariasis (Gyapong et al., 2002; Lindsay and Thomas, 1999), leishmaniasis (Nieto et al., 2006; Salahi-Moghaddam et al., 2010), Chagas disease (Mischler et al., 2012) *Opisthorchis viverrini* (Suwannatrai et al., 2011), STHs (Pullan and Brooker, 2012) and others

(Bergquist and Rinaldi, 2010). Models such as these are useful in developing and managing intervention programs (Daash et al., 2009; Dongus et al., 2011).

There are two main approaches to developing GIS models for disease (Malone, 2005). The statistically-based approach is used when knowledge of the biological requirements of the organism of interest is limited and only survey data is available (Lindsay and Thomas, 1999; Pullan and Brooker, 2012; Thomson et al., 1997). The second approach, the biology-based model, is initiated with known environmental parameters that are critical to the survival of the organism, that have been determined from laboratory or field studies (Nieto et al., 2006; Valencia-López et al., 2012). Statistical analysis is however required for the model validation process using independent field data (Malone, 2005).

A parameter that has become useful in biology-based modeling is that of heat units, also known as Growing Degree Days (GDD). It was initially developed for and used in crop farming to determine timing of phenotypic events (McMaster and Wilhelm, 1997). It is calculated using climatic data for the study area of interest by obtaining the average daily temperature and subtracting the base temperature (T_{base}), below which the organism cannot develop in the environment (McMaster and Wilhelm, 1997).

$$GDD = \frac{(T_{max} - T_{min})}{2} - T_{base}$$

When applied to parasitology, GDD is used to determine the period of time an organism needs to develop from one stage to another in the environment, and to assess if a given environment is suitable enough to support such development and therefore the

propagation and transmission of the parasite (Malone, 2005). In addition to GDD, water budget (WB) has also proved important in risk modeling. WB is the ratio of precipitation (P) and evapotranspiration (ET), i.e. P/ET . It is a measure of soil moisture saturation, with 0 representing aridity and 1 representing full soil saturation. Values >1 represent water run-off (Shuttleworth, 1983). The concept of GDD and WB has been used successfully to generate predictive models of fascioliasis in the United States, Ethiopia, Ecuador and southern Brazil (Dutra et al., 2010; Fuentes et al., 2005; Malone et al., 1987; Malone et al., 2001; Valencia-López et al., 2012), schistosomiasis in China and East Africa (Malone, 2005; Zhou et al., 2001) and visceral leishmaniasis in Bahia, Brazil (Nieto et al., 2006).

1.3 Soil-Transmitted Helminths

Soil-transmitted helminths (STH) are those nematodes that have a soil-based or environmental developmental stage in soil in order for them to become infective. These include hookworms (*Necator americanus* and *Ancylostoma duodenale*), *Ascaris lumbricoides* and *Trichuris trichiura*. STH occur worldwide and are particularly important in tropical and subtropical countries that have high poverty levels in Asia, sub-Saharan Africa, and the Americas (Bethony et al., 2006; Brooker et al., 2006a; de Silva et al., 2003; Horton, 2003). As a group they are considered among the neglected tropical diseases (NTDs) and have received considerable attention since the United Nations millennium declaration in 2000 (Hotez et al., 2008). The association between STH and poverty is widely accepted (de Silva et al., 2003; Hotez, 2008) and has been shown to be significant and complex, with each component perpetuating the other (de Silva et al., 2003). Reduced productivity of chronically infected people perpetuates poverty and the

poor economic growth in such countries means poor sanitation and resultant high of STH (de Silva et al., 2003; King, 2010).

1.3.1 Biology of STHs

Soil transmitted helminths belong to the Kingdom Animalia, Phylum Nematoda and Class Secernentea.

1.3.1.1 *Ascaris lumbricoides*

Order: Ascaridida

Superfamily: Ascaridoidea

Family: Ascarididae

Subfamily: Ascaridinae

A. lumbricoides eggs are released into the environment in feces and develop to L₂ infective larvae contained in the egg after 18 or more days, depending on environmental moisture and temperature (Aaron, 1950; Anderson, 2000; Brown, 1927). When the infective eggs are ingested, the larvae develop into L₃ before they hatch in the small intestine and penetrate the mucosa. They are then transported via the venous system to the heart and lungs from about six days post infection (Dold and Holland, 2011; Geenen et al., 1999). The larvae develop further in the lungs for approximately 10-14 days. They then penetrate the alveolar wall and travel up the bronchial tree and trachea and are swallowed and then develop to adults in the small intestine (Figure 1.1) (Dold and Holland, 2011). The adult worms are the largest intestinal nematodes infecting the human intestine and measure between 15 to 35cm (Anderson, 2000) and can live up to 23 weeks (Olsen et al., 1958). Adult worms mate and females lay up to 200,000 eggs per day. Eggs measure 65-75 x 35-50 µm and are passed in feces unembryonated (Anderson, 2000).

Infertile and undeveloped eggs are not infective until fully embryonated eggs, which can remain viable in the environment for up to five years under favorable conditions (Anderson, 2000).




Figure 1.1: Lifecycle of *A. lumbricoides* (C.D.C., 2013a)

1.3.1.2 *Trichuris trichiura* (whipworm)

Order: Enoplida

Superfamily: Trichinelloidea

Family: Trichuridae

Subfamily: Trichurinae

Unembryonated eggs of *T. trichiura* are passed in feces and develop to infective L₃ larvae within the egg in 11 or more days depending on environmental temperature and moisture (Anderson, 2000; Brown, 1927). *T. trichiura* has a direct lifecycle, i.e. no tissue

migratory phase. When the infective eggs are ingested they hatch in the small intestine and develop further into adult worms (Figure 1.2). The adult worms reside in the large intestine (cecum and colon) and thread their anterior ends into the mucosa, causing significant inflammation. Female worms start laying eggs 60-70 days after establishment and the unembryonated eggs are passed in feces. They can oviposit between 3,000 to 20,000 eggs per day (Anderson, 2000). Adult worms are thought to live an average of 1-2 years in the cecum/colon (Anderson, 2000).




Figure 1.2: Lifecycle of *T. trichiura* (C.D.C., 2013c)

1.3.1.3 Hookworm

Order: Strongylida

Superfamily: Ancylostomatoidea

Family: Ancylostomatidae

Subfamily: Ancylostomatinae Bunostominae

Genus: *Ancylostoma* *Necator*

Ancylostoma duodenale and *Necator americanus* are the two most important hookworms infecting humans (Brooker et al., 2004). Their two to eight cell stage eggs, measuring 50-80 x 36-42 µm, are passed in feces into the environment and develop to L₁ larvae within 1-2 days (Chandler, 1929) (Figure 1.3). The eggs do not develop in temperatures less than 15°C (Udonsi and Atata, 1987) and the eggs of *N. americanus* die at temperatures less than 7°C while those of *A. duodenale* survive (Anderson, 2000). The L₁ hatch and develop to L₂ larvae which feed on bacteria in the soil and feces. The L₂ larvae develop to L₃ which do not feed but retain the L₂ cuticle till they infect humans by the oral route (*A. duodenale*) or skin penetration (both species) (Anderson, 2000; Looss, 1911). Infective larvae (L₃) develop in five or more days depending on environmental temperature and moisture (Chandler, 1929) and can survive in moist soil for up to six weeks (Svensson, 1925; Udonsi and Atata, 1987). The optimum temperature for development of the free living stages ranges from 21-27°C for *A. duodenale* and 25-28 for *N. americanus* (Anderson, 2000; Crompton and Whitehead, 1993). On skin penetration, intense pruritus is experienced at the site and is known as ground itch or hookworm dermatitis (Anderson, 2000; Schad and Warren, 1990). Larvae travel via the lymphatic and circulatory system to the lungs (Looss, 1911) and, after penetrating into the alveoli, travel up the bronchial tree to the pharynx and are swallowed. The pulmonary phase may induce episodes of coughing and pharyngitis 7-14 days post infection (Nawalinski and Schad, 1974; Schad and Warren, 1990). The larvae take up residence in the small intestine and develop into

L₄ which feed on mucosa and blood and then into adults which attach to the mucosa and feed on blood (Pawlowski et al., 1991). *A. duodenale* females can produce 10,000 to 25,000 eggs per day and *N. americanus* 6,000 to 11,000 eggs per day (Anderson, 2000; Crompton and Whitehead, 1993). The prepatent period for *N. americanus* is 44 or more days (Crompton and Whitehead, 1993; Looss, 1911). *A. duodenale* is capable of hypobiosis, an arrested development stage, and the prepatent period can therefore be as little as 31 days up to eight months (Nawalinski and Schad, 1974). Life expectancy of the adult hookworms is 2-7 years (Anderson, 2000; Brooker et al., 2006b; Crompton and Whitehead, 1993). The eggs of the two common species of hookworm cannot be differentiated by microscopy; however, the infective larvae and adults can (Table 1.1).




Figure 1.3: Hookworm lifecycle (C.D.C., 2013b)

Table 1.1. Morphological differences between *A. duodenale* and *N. americanus* (Chandler, 1929; Crompton and Whitehead, 1993; Looss, 1911; Nichols, 1956; Pawlowski et al., 1991; Svensson and Kessel, 1926; Wu and Peng, 1965).

Stage	Features	<i>A. duodenale</i>	<i>N. americanus</i>
Infective larvae	Buccal cavity		More prominent
	Genital primordium		More anterior and smaller
	Esophagus	Narrower Continuous with intestine	Broader Constriction (gap) at the junction with intestine
	Intestine	Narrower than esophageal bulb at junction with esophagus	As wide as esophageal bulb at junction
	sheath	Striations inconspicuous	Conspicuous striations in sheath
	Body length (µm)	600-700	500-600
	Length of sheath (µm)	720	660
Adult	Buccal cavity	Two pairs of teeth	Two cutting plates
	Anterior end	Less distinct curve Less tapered	Distinct curve More slender and more finely tapered
	Size (mm)	Male: 8-11 Female: 10-13	Male: 7-9 Female: 9-11

1.3.2 Public Health Significance

STH infections tend to cause chronic infections with low mortality rates. Symptoms associated with STH include diarrhea, reduced appetite and weight gain, impaired cognitive abilities, malnutrition, anemia, abdominal pain and stunted growth (Beaver, 1988; Bethony et al., 2006; Eppig et al., 2010; Ezeamama et al., 2005; Haque et al., 2010;

Jardim-Botelho et al., 2008; Stephenson et al., 1993; Yap et al., 2012). *A. lumbricoides* can cause life threatening intestinal obstruction, when the adult parasite is present in large numbers particularly in young children (Akgun, 1996; Villamizar et al., 1996) and rarely airway obstruction (Bailey and Warner, 2010). *A. lumbricoides* has also been associated with Vitamin A deficiency due to malabsorption. (Mahalanabis et al., 1976). In addition to the symptoms of STH infection above, *T. trichiura* causes dysentery and occasionally rectal prolapse in severe cases (Cooper et al., 1992; Haque et al., 2010). Trichuris may also cause anemia, although hookworms cause greater blood loss with resultant iron deficiency anemia (Stoltzfus et al., 1997). The iron deficiency anemia related to hookworm infection is more pronounced in moderate to high infections, particularly in children, but even low intensity infections cause lower hemoglobin levels in adults (Brooker, 2010).

The WHO uses Disability-adjusted life years (DALYs) to estimate the global burden of disease caused by any disease. DALYs are an estimate of the number of healthy life years lost due to disability or premature death caused by a particular disease and are derived by assigning disability ‘weights’ to different conditions and multiplying that weight by the number of years that an infected person is affected by that condition (Brown et al., 2009). Among the DALYs calculated for NTDs in 2010, STHs rank second with 5,184,000 DALYs after malaria (82,685,000 DALYs), with the highest impact being in sub-Saharan Africa (Murray et al., 2012). Ascariasis accounted for 1,315,000, trichuriasis 638,000 and hookworm disease had the highest with 3,231,000 DALYs (Murray et al., 2012). There was a 42.5% reduction in DALY impact by STH from 1990 to 2010, with the highest reduction being in ascariasis (68.8%) and the lowest (17.9%) in hookworm infections

(Murray et al., 2012). The great reduction in ascariasis may be due to the high efficacy of benzimidazoles against *A. lumbricoides* and improvements in sanitation, especially in China and Brazil (Fenghua et al., 1998; Pullan and Brooker, 2012).

1.3.3 Prevalence

It is estimated that over a billion of the world's population is infected with at least one of the STH (Bethony et al., 2006) and more than 300 million suffer from severe morbidity (WHO, 2010). The majority of STH are found in the tropics and subtropics (Figure 1.4) and WHO estimates that 130 countries are endemic (WHO, 2008). China, Southeast Asia, Central Africa, and the coastal regions of West Africa have the highest prevalence of *A. lumbricoides* infection (de Silva et al., 2003). *T. trichiura* prevalence is highest in Central Africa, southern India and Southeast Asia, while hookworm infections are common throughout most of sub-Saharan Africa, south China and Southeast Asia (de Silva et al., 2003).

Although there was a great decline in STH estimated prevalence in Latin America and the Caribbean (LAC) from 1994 to 2003, due to widespread control efforts, the risk of transmission remains high due to the highly permissive climate of this region (de Silva et al., 2003; Pullan and Brooker, 2012). Prevalence estimates for LAC in 2003 were 16% for *A. lumbricoides*, 19% for *T. trichiura* and 10% for hookworm (de Silva et al., 2003). However, the prevalence varies within regions and countries based on socioeconomic factors, urbanization and sanitation (Pullan and Brooker, 2012). STH prevalence tends to be higher in rural areas due to lower socioeconomic levels, farming practices and poor sanitation (Brooker, 2010; Hotez et al., 2005). Economic development of rural areas has

been linked to a reduction in STH prevalence, as seen in the Jiangsu province of China (Fenghua et al., 1998).




Figure 1.4: Global distribution of Soil-transmitted helminths. Adapted by permission from Macmillan Publishers Ltd: Nature Reviews Microbiology (Savioli and Albonico, 2004) (Appendix A).

1.3.4 Diagnosis

Since the symptoms of soil-transmitted helminthiasis are mostly non-specific, diagnosis of STH is primarily reliant on demonstration of helminth eggs in feces. The most commonly used method, and one that is recommended by WHO for surveillance of STH is the Kato-Katz method (Montresor et al., 1998). It was initially developed for the detection of *Schistosoma mansoni* in feces (Katz et al., 1972) but has since been adopted for use in detection of STH as well (Endris et al., 2013; Funk et al., 2013; WHA, 2001). The method is both quantitative and qualitative and has proved to be relatively robust and field-friendly as it does not require the use of expensive equipment like a centrifuge. However, hookworm eggs tend to collapse after approximately 30 minutes of the slide

being prepared and this could lead to erroneous results if there is a delay in examination of the slide (Booth et al., 2003). Another disadvantage is the loss of sensitivity in the test due to the small amount (41.7mg) of feces that is examined at a time (Booth et al., 2003; Endris et al., 2013). However, this may be overcome by the preparation of two or three slides per sample of feces and also repeated sampling for two to three consecutive days to overcome the day-to-day variation in egg shedding by adult worms (Booth et al., 2003), although it would be more time consuming and costly (Utzinger et al., 2008). Tarafder et al. (2010), used a Bayesian approach to determine the sensitivity and specificity of Kato-Katz method in the absence of a gold standard, and found that one stool sample was adequate with regard to detection of *Ascaris* and *Trichuris* but not hookworm eggs, and that two samples (from two days) gave better results.

The formalin ethyl acetate (FEAC) concentration/sedimentation method (Allen and Ridley, 1970; Young et al., 1979) is also used to detect STH eggs as well as many other parasite eggs, larvae, oocysts and cysts. Because of the formalin, the eggs are preserved and the sample can be examined long after collection (Baker, 2007). This is advantageous for samples that need to be transported for examination at a distant/referral facility. Drawbacks include the large amount of debris, the requirement of a centrifuge and safe disposal of reagents like ethyl acetate (Allen and Ridley, 1970; Baker, 2007). In a study in Côte d'Ivoire, FEAC yielded more *Trichuris* positives than the Kato-Katz method (Glinz et al., 2010). When compared to Kato-Katz, FEAC detected more hookworm positives and this was believed to be because of the larger quantity of feces that can be examined (Endris et al., 2013).

Spontaneous sedimentation, also known as the Hoffman-Pons-Janer method, is an easy and inexpensive method because it only utilizes water and a conical flask (Carvalho et al., 2012). However, it requires that the sample be examined the same day of the sedimentation or be refrigerated to prevent the hatching of hookworm eggs. If the sample is unpreserved, hookworm L1 larvae may be confused with *Stongyloides stercoralis* and need to be differentiated, since the treatment for the two is different (Garcia, 2001). In a comparison, the Kato-Katz method was found to perform better in detection of helminths than spontaneous sedimentation (Carvalho et al., 2012).

Flotation procedures which can be both quantitative and qualitative have not been used widely in human parasitology but are widely used in veterinary parasitology. The McMaster method (MAFF, 1986) typically uses sodium chloride solution as a flotation medium and is useful for the recovery of helminth eggs with the exception of operculated and schistosome eggs (Truant et al., 1981). In a study that compared the sensitivity of Kato-Katz method with that of the McMaster method, the results for the two were comparable except with regard to *Ascaris*, where Kato-Katz detected significantly more positive samples and higher egg counts (Levecke et al., 2011). In testing the McMaster method against an ether-based sedimentation method for detection of *Trichuris* in primate fecal samples, the McMaster test was less sensitive but detected significantly more eggs than the ether-based concentration method (Levecke et al., 2009).

Zinc sulfate solution (Specific gravity (SG) 1.18) (Faust et al., 1938) is another flotation medium that has been used extensively in veterinary parasitology (Dryden et al., 2005), with and without centrifugation, although centrifugation resulted in detection of more positive samples and higher egg counts (Dryden et al., 2005; Zajac et al., 2002). The

centrifugal force created causes heavier fecal debris to sink to the bottom of the tube therefore prevents obscuring of parasite eggs (Dryden et al., 2005). When used on human fecal samples, zinc sulfate was more effective for the detection of hookworm eggs than FEAC, but not as effective for *Ascaris* and *Trichuris* eggs (Truant et al., 1981). Brandelli et al. (2012) used centrifugal flotation with zinc sulfate on human fecal samples in conjunction with other methods but no comparison of methods were reported. Another flotation medium that is widely used in veterinary parasitology but not in human parasitology is the sucrose (Sheather's) solution (SG 1.25-1.27). Sucrose has been used in a few human parasitology studies with success (Shakya et al., 2006) although information on how it performs in comparison with other helminth egg concentration methods is lacking. When compared with other flotation solutions, it was the best for gastrointestinal strongyle egg visualization and maintained good parasite egg quality for longer (Cringoli et al., 2004). Due to the higher SG, sucrose flotation results in higher fecal egg counts than zinc sulfate and, like zinc sulfate flotation, centrifugation enhances results (Cringoli et al., 2004; Dryden et al., 2005).

The FLOTAC method was developed more recently (Cringoli, 2006) that utilizes a FLOTAC[®] apparatus. The apparatus is used to perform flotation in a centrifuge and the retention of the upper portion of the floating suspension can then be visualized in the FLOTAC chamber under a microscope (Cringoli, 2006). FLOTAC can be both a quantitative and qualitative technique and utilizes up to a gram of feces. There is a choice as to the flotation medium that can be used in the apparatus (Cringoli et al., 2004) and this allows for flexibility. In comparison to triplicate Kato-Katz in two different STH endemic settings, the FLOTAC was found to be more sensitive in detection of STH eggs

but yielded lower fecal egg counts (Glinz et al., 2010; Knopp et al., 2009). Depending on the flotation medium used, also allowed for detection of larvae, cysts and oocysts of protozoa (Cringoli, 2006). FLOTAC performed superior to FEAC in recent surveys (Glinz et al., 2010; Utzinger et al., 2008). When tested only for hookworm egg detection, FLOTAC was much more sensitive than Kato-Katz or FEAC, implying that there may be an underestimation of prevalence of human hookworm infections and probably other STHs and *S. mansoni* by routinely used methods (Kato-Katz and FEAC) (Utzinger et al., 2008).

Serology for antibody detection using enzyme linked immunosorbent assays (ELISA) have been developed and used to detect intestinal parasites such as *O. bifurcum* and *N. americanus* (Pit et al., 2001; Polderman et al., 1993). However, they are not very sensitive or specific and not able to differentiate current from past infection (Basuni et al., 2011; Verweij et al., 2001).

Coproculture is useful for differentiating nematode infective larvae for strongylids such as *N. americanus*, *Ancylostoma* sp, *O. bifurcum*, *Trichostrongylus* spp and *S. stercoralis* (Harada and Mori, 1955; Okazaki et al., 1988; Yong et al., 2007). However, coproculture is time consuming, taking six or more days, is labor intensive, and requires skilled personnel (Basuni et al., 2011; de Gruijter et al., 2005).

The development of PCR for the detection and identification of intestinal parasites in the last two decades has proved very helpful in parasitology (Verweij et al., 2001), and, together with the sequencing of the entire mitochondrial genomes of *N. americanus* and *A. duodenale*, study of the genetic variability in hookworms has become feasible (Chilton

and Gasser, 1999; Hu et al., 2002). Hookworm eggs are difficult to differentiate by microscopy because of their similarity in size.

Gasser et al. (1993) and Romstad et al. (1997) found the ribosomal (r)DNA region encompassing the internal transcribed spacer (ITS-1), 5.8S, ITS-2 and part of the 28S rRNA gene as being ideal for PCR amplification as it was well conserved among strongylid nematodes but had adequate sequence difference between species for delineation (Figure 1.5). Genetic markers in the ITS-2 region were used develop species-specific primers to distinguish *N. americanus* from *O. bifurcum* (Romstad et al., 1997; Verweij et al., 2001) and to differentiate *N. americanus* eggs from those of *A. duodenale* in fecal samples using a semi-nested PCR technique (de Gruijter et al., 2005).




Figure 1.5. Schematic representation of part of the rDNA transcriptional unit and relative locations of primers NC5, NC1 and NC2 commonly used for hookworm species differential diagnosis. Adapted from (Romstad et al., 1997; Yong et al., 2007).

1.3.5 Treatment and Control

False negative results have been reported with PCR and have been attributed to the small amount of fecal sample used as compared to that used in coproculture or fecal concentration methods and also possible DNA degradation from long periods of storage (de Gruijter et al., 2005; Ngui et al., 2012b). There is a high degree of similarity between the ITS-2 regions of *A. duodenale* and *A. ceylanicum* (a zoonotic hookworm species) with only a few nucleotide differences, therefore, de Gruijter et al. (2005) found it necessary to recommend sequencing of the resulting PCR amplicons to differentiate the

two species. Real-time PCR assays have also been developed for simultaneous diagnosis of hookworms and other species of intestinal parasites (Basuni et al., 2011). A PCR-linked restriction fragment length polymorphism (RFLP) was developed and used by Hawdon (1996) to differentiate *N. americanus* from *A. duodenale* using endonucleases and it was found that *Taq1* worked best to differentiate single and mixed infections of hookworm.

With the growing international attention that neglected tropical diseases (NTD) were receiving in the 1990s, WHO developed a clear policy for the control of STH in endemic areas (Hotez et al., 2008; Montresor et al., 1998). The policy was based on prevalence and intensity of infection of the STH in a community (WHA, 2001). The Mass Drug Administration (MDA) concept was first applied to the control of lymphatic filariasis in China and was adopted by WHO for incorporation in control programs for STH and other NTDs (Hotez et al., 2007). MDA involves the distribution and administration of drugs to whole communities, rather than individuals infected or at risk of infection with preventable diseases that are endemic (Brown et al., 2009). In order to properly administer drugs for STH, WHO recommended that surveys be done to determine the prevalence and intensity of infection (Table 1.2) by STH in a community, and developed survey criteria guidelines (Montresor et al., 1998).

Table 1.2: Classes of intensity of infection. Adapted from Montresor et al. (1998)

	Light intensity	Moderate intensity	Heavy intensity
<i>A. lumbricoides</i>	1-4999 epg	5000-49999 epg	50000 epg
<i>T. trichiura</i>	1-999 epg	1000-9999 epg	10000 epg
Hookworms	1-1999 epg	2000-3999 epg	4000 epg

Community categories were developed as a guide for MDA and WHO recommends MDA for communities with 50% or higher prevalence of STH, irrespective of intensity of infection. However, with regard to hookworm, when the prevalence is over 20% and anemia is prevalent, then MDA is recommended (Montresor et al., 1998).

The drugs that are commonly used for MDA are the benzimidazoles, albendazole and mebendazole (WHA, 2001). Benzimidazole drugs bind selectively to β -tubulin of nematodes, cestodes and trematodes and inhibit microtubule formation. This causes a slow starvation of the parasite and inhibition of egg production (Martin, 1997). In addition to benzimidazoles, WHO also recommends praziquantel (tetrahydropyrimidine) and levamisole (imidazothiazole) for STH (WHA, 2001). These drugs are nicotinic agonists at acetylcholine receptors of nematode muscle and cause spastic paralysis (Martin, 1997). In some cases, a combination of drugs resulted in better efficacy of treatment than when a single drug was used. For the treatment of hookworm, a combination of mebendazole and levamisole showed significantly higher efficacy (Albonico et al., 2003) than when either drug was used alone. In other studies, the addition of ivermectin to albendazole or mebendazole showed higher efficacy against *T. trichiura* (Ismail and Jayakody, 1999; Knopp et al., 2010). Ivermectin causes paralysis of the pharyngeal pumping of nematodes. With the reported relatively low efficacy of benzimidazoles, and other drugs recommended by WHO against *T. trichiura* (Keiser and Utzinger, 2008), oxantel pamoate, a nicotinic agonist like pyrantel, were used against *T. trichiura* and found to have higher efficacy, particularly when combined with mebendazole (Keiser et al., 2013). A differential anthelmithic effect by levamisole has been reported among the human hookworm species, in which a greater efficacy was

observed against *A. duodenale* than *N. americanus* (Albonico et al., 2003; Albonico et al., 1998).

Concerns regarding the possible development of drug resistance against benzimidazoles has grown since the implementation of MDA in most STH endemic areas (Vercruysse et al., 2011). In investigating drug efficacy, fecal egg reduction (FER) rate is recommended, rather than cure rate, as it is a closer representation of worm burden (Albonico et al., 2004; Vercruysse et al., 2011). The egg hatch assay, which is useful for assessing hookworm resistance, is another test that can be used to determine drug efficacy, as hookworm eggs hatch within 1-2 days (Albonico et al., 2004). In light of reported FER rates of <90% in studies in Ghana (Humphries et al., 2013) and Zanzibar (Albonico et al., 2003), closer monitoring of MDA programs was called for in other regions as well. The association between the response to albendazole treatment and β -tubulin genotype frequencies was studied for STH from three regions of the world, and although *Ascaris* and hookworm did not show much modification in β -tubulin genotype frequencies, *Trichuris* showed evidence of resistance even in benzimidazole naïve populations (Diawara et al., 2013). This may explain the reduced efficacy of benzimidazoles in single drug administration protocols noted by other researchers.

Improved sanitation has had mixed reported effects on parasite infection. For the most part there have been reductions in prevalence and intensities of infections especially with ascariasis (Esrey et al., 1991; Jeffery, 1960; Ziegelbauer et al., 2012). However, the impact on hookworm infections has been less remarkable and in some cases lacking (Esrey et al., 1991; Gross et al., 1989). Although deworming remains the most effective way of controlling the spread of STH (de Silva et al., 2003; Hotez et al., 2006), it is

recommended that improvements to sanitation and hygiene education be included in control programs for STH in order to reduce infection incidence and re-infections (Ziegelbauer et al., 2012).

1.4 Human Hookworm Infection

A. duodenale and *N. americanus* are the most important hookworms in humans. *A. duodenale* is found mainly in subtropical climates up to 52°N (Araújo et al., 1988), mostly in the Middle East, northern Africa, India, Australia and Europe (de Silva et al., 2003). *N. americanus* has a more tropical distribution and is endemic in South and Southwest China, South India, Southeast Asia, sub-Saharan Africa, and Central and South America (Brooker et al., 2004) (Figure 1.6). However, mixed infections do occur usually with prominence of one or the other (Araújo et al., 1988). There have also been a few unexpected findings with regard to distribution of hookworm species. For instance, Jonker et al. (2012) using PCR, in Malawi, a sub-Saharan country, unexpectedly found that *A. duodenale* was prevalent. In Malaysia, Ngui et al. (2012a) also using molecular techniques, unexpectedly found *A. ceylanicum* where initially they thought only have *N. americanus* was present in humans. In Ghana, *A. duodenale* was found in a population that had predominantly *N. americanus* (de Gruijter et al., 2005).

Mixed infections with other stongylids are also possible. Pit et al. (1999) found hookworm with *Oesophagostomum bifurcum* in Togo and Ghana and Sato et al. (2010) in Laos, found *Trichostrongylus colubriformis* as well as *N. americanus*, *A. duodenale* and *A. caninum* in human fecal samples using PCR. Since it is not possible to differentiate the eggs of human hookworm spp. from each other by light microscopy, coproculture or molecular methods are needed to definitively identify the species hookworm present.

Zoonotic hookworms, *A. ceylanicum* (Traub et al., 2008) and *A. caninum* (Landmann and Prociv, 2003; Prociv and Croese, 1996) have also been detected in Thailand, Laos and Australia. *A. ceylanicum* can cause patent infections (Knopp et al., 2010) but *A. caninum* does not. Instead, *A. caninum* oral infection results in the development of an immature adult in the intestine with associated eosinophilia, intestinal ulceration and abdominal pain (Prociv and Croese, 1996).




Figure 1.6: Climatic suitability for hookworm transmission defined by land surface temperature and aridity. Areas were defined as stable (dark green), unstable (light green, where infection prevalence <2%), or no risk (light grey). Adopted from Pullan and Brooker (2012).

Hookworm disease is characterized by iron deficiency anemia (Brooker et al., 2004).

This anemia can be particularly severe in children and women of reproductive age who also have concurrent malnutrition (Brooker et al., 2004). However, even in adults without malnutrition, chronic infection can cause a reduction in hemoglobin (Santos et al., 2013). Iron deficiency anemia can result in low birth weight, premature birth and maternal death in pregnant women as well as low weight gain, reduced appetite and poor physical fitness in children (Crompton, 2000; Stephenson et al., 1993). The anemia is brought about by

the blood loss associated with the hematophagous L₄ and adult stages of the hookworm. Hookworms also produce anticoagulants which cause the intestinal mucosa to continue bleeding after the hookworm has stopped feeding. *A. duodenale* causes greater blood loss and it is estimated that the daily loss is 0.14 – 0.26 ml of blood per worm, while *N. americanus* is estimated at 0.02 – 0.07 ml per worm (Albonico et al., 1998; Pawlowski et al., 1991; Santos et al., 2013). Infections caused by *A. duodenale* have been associated with higher levels of iron deficiency and anemia in children (Albonico et al., 1998). Low intensity infections with multiple parasites are also likely to cause clinically significant anemia (Ezeamama et al., 2005).

1.4.1 Hookworm in Brazil

The tropical climate in Brazil is highly conducive for the development of free-living stages of hookworm in the environment (Mudenda et al., 2012). Socio-economic factors play an important role in facilitating or deterring the transmission of hookworm (Brooker et al., 1999). The prevalence of hookworm has been reported as being as low as 0.27% among higher income people (Santos et al., 2013) as opposed to rural and poor populations where it was found to be 9% among patients in the Amazon region (Araujo and Fernández, 2005). In poor rural communities, the prevalence of hookworm has been reported to be as high as 80.2% (Gonçalves et al., 1990).

Few studies have gone on to determine the species of hookworm in Brazil. In one study in the state of Minas Gerais, only *N. americanus* was found (Fleming et al., 2006). In the northeastern region of Brazil, Gonçalves et al. (1990) found 70.9% of the hookworms were *N. americanus*, 2.1% were *A. duodenale* and the rest could not be identified. Okazaki et al. (1988) reported similar findings in child out-patients in a farming

community in the same region, albeit with *A. duodenale* up to 12.5%. The Harada-Mori method of coproculture was used to identify larvae morphologically in both studies.

Prevalence of hookworm tends to be higher among males compared to females (Santos et al., 2013) and this has been attributed to activities that males may be involved in that put them at greater risk for infection. In a study among upper and middle class patients in Salvador, Brazil, the 30-50 year age group was the most commonly infected with hookworms (Santos et al., 2013) and there is general agreement among researchers that prevalence tends to increase with age and plateaus in adulthood (Fleming et al., 2006; Stoltzfus et al., 1997). However, others have found that neither age nor sex was correlated with prevalence where all family members were involved in agricultural activities (Barreto et al., 2000; Gonçalves et al., 1990; Okazaki et al., 1988).

1.5 Common Intestinal Parasites of Dogs

Intestinal parasites in dogs are widespread and are important because they cause morbidity and mortality in dogs as well as having a zoonotic potential for dog owners and those working with dogs (Blagburn et al., 1996). In dogs, intestinal parasites may cause enteritis, anemia, bloody diarrhea, weight loss, dermatitis and vomiting (Bowman, 2009) and contamination of the environment with dog feces containing eggs, oocysts, and cysts of parasites presents a public health risk. In addition to risk to humans, dog parasites can cause infection in other animals. For instance, *Toxocara canis* can cause visceral larval migrans (VLM) in cats and rabbits.

1.5.1 Biology and infection

1.5.1.1 Hookworm

The species of hookworm found in dogs are *Ancylostoma caninum*, *A. braziliense*, *A. ceylanicum* and *Uncinaria stenocephala*. *A. caninum* occurs worldwide, *A. braziliense* is mostly in the southern US and central/ South America, *A. ceylanicum* is mostly found in Asia (Conlan et al., 2012; Smout et al., 2013; Traub et al., 2008) and *U. stenocephala* is found in the cooler climate of the northern America and Europe (Taylor et al., 2007; Traub et al., 2004). Canine hookworms have a zoonotic potential and cause cutaneous larvae migrans (CLM) or ‘creeping eruption’ when they penetrate human skin (Bowman et al., 2010). *A. braziliense* is the most common cause of CLM, while *A. caninum* has been reported to cause eosinophilic enteritis in humans in Australia (Prociv and Croese, 1996) and has been implicated as a cause for diffuse subacute neuroretinitis (Harto et al., 1999). In the cases of eosinophilic enteritis encountered, only a single adult worm was found. *A. ceylanicum* is the only canine hookworm capable of causing a patent infection in humans and it has been demonstrated experimentally as well as by natural infection (Ngu et al., 2012b; Traub et al., 2008).

The lifecycle of canine hookworm is similar to the human hookworm lifecycle. Canine hookworms also infect via skin penetration and follow the alveolar route of tissue migration before settling in the small intestine and maturing to hematophagous adults (Figure 1.7). However, the migrating larvae of canine hookworms may also undergo arrested development in the muscles and intestinal walls of the host and serve as source of trans-mammary infection (ingestion of L₃) to puppies (Bowman et al., 2010; Stone and Girardeau, 1968). Arrested *A. caninum* larvae tend to reactivate spontaneously, regardless

of whether the intestine has been evacuated of adults or with immune suppression (Schad and Page, 1982). Human infection by canine hookworms is also by skin penetration, but the larvae migrate in the epidermis forming erythematous tracts called cutaneous larva migrans (Bowman et al., 2010). Only *A. ceylanicum* is able to migrate to the intestine, mature, mate and produce eggs. The prepatent period for *A. caninum* via ingestion is 2-3 weeks and via skin penetration it is 4-5 weeks (Bowman, 2009). For *A. braziliense*, the prepatent period is reported as 13 or more days by either oral or percutaneous infection (Liotta et al., 2012a).




Figure 1.7. Zoonotic Hookworm (CDC, 2012)

Hookworms produce anticoagulants while feeding (Furmidge et al., 1996; Hotez and Cerami, 1983; Kasper et al., 2001; Stassens et al., 1996) and *A. caninum* causes blood loss of approximately 0.1ml per day per worm. This level of blood loss can cause severe and life threatening anemia in puppies. The LD₅₀ of *A. caninum* in nursing puppies is

reported to be 75 worms per pound of body weight (Miller, 1971). The peak of the blood loss occurs even before eggs appear in the feces due to blood sucking by the L₄ stage, but it coincides with bloody diarrhea (Bowman, 2009). *A. caninum* is the most pathogenic of the canine hookworms because of the amount of blood loss associated with it.

1.5.1.2 *Toxocara canis* (roundworm)

T. canis is an ascarid similar to *A. lumbricoides* in humans. It is the largest nematode (10-15cm long) found in the digestive tract of dogs and other canids (Bowman, 2009). It has a worldwide distribution and the eggs are quite resistant to environmental conditions, able to survive for up to four years (Slifko et al., 2000). Eggs are unable to develop when exposed to temperatures less than -11°C (Slifko et al., 2000) and more than 34°C (Gamboa, 2005). Larvae appear in the eggs in 2-3 weeks depending on environmental conditions (Anderson, 2000) with higher humidity and temperature accelerating the development (Gamboa, 2005). Transmission is by ingestion of the infective egg, which contains infective larvae. The eggs hatch in the intestine and the larvae penetrate the intestinal mucosa and migrate through the hepato-tracheal route like *A. lumbricoides* (Beaver, 1969). They mature into adults in the small intestine, mate and lay eggs which are passed unembryonated in feces (Figure 1.8). The prepatent period is 2-4 weeks and one *T. canis* female can produce up to 85,000 eggs per day. Larvae are also capable of encysting in the intestinal walls and somatic tissues, particularly in adult immune dogs, where they become dormant for prolonged periods of time. In female dogs in late pregnancy, tissue larvae reactivate and infect puppies transplacentally or through the transmammary route, although the latter is considered a minor route of transmission (Bowman, 2009). Heavy prenatal infection of puppies can cause severe abdominal

distension and discomfort and life-threatening obstruction of intestines, bile and pancreatic ducts may occur (Bowman, 2009). Infection can also occur by ingestion of paratenic hosts (eg. small mammals) in which the larvae hatch and encyst in tissues (Okulewicz et al., 2012). Ingestion of encysted larvae by dogs is followed by maturation and oviposition by adults in the small intestine.




Figure 1.8. *Toxocara canis* life cycle (CDC, 2013)

If humans get infected accidentally by ingestion of infective eggs or encysted larvae in paratenic hosts, larvae fail to mature and instead migrate extensively in the body causing inflammation in various organs like the liver, brain, lungs, muscle and heart. This condition is called visceral larva migrans (VLM) and if the migration is in ocular tissue, ocular larva migrans (OLM) (Bowman, 2009). Seroprevalence of *T. canis* in a national survey in humans in the USA was 14% and was associated with poverty and dog ownership among other risk factors (Won et al., 2008).

T. leonina is an ascarid that occurs both in canids and felids (Okulewicz et al., 2012). It is similar in size, to *T. canis*, but they differ morphologically in that *T. leonina* males have a conical tail with no caudal alae while *T. canis* males have a digitiform appendage and caudal alae (Okulewicz et al., 2012). Although the eggs of the two species have similar dimensions, *T. canis* have the pits of the eggshell on the outer layer of the eggs, while *T. leonina* eggs are more translucent with a smooth shell, no striations or albuminous coat on the outer layer (González et al., 2007). Larvae of *T. leonina* develop to the infective stage in eggs in as few as three days at 30°C (Anderson, 2000). *T. leonina* is also more tolerant to cold temperature extremes with eggs being able to survive temperatures of -15°C for up to 40 days while those of *T. canis* do not (Okulewicz et al., 2012). *T. leonina* is sometimes referred to as the ‘Northern roundworm’ because it is most prevalent in cool climates. *T. leonina* has been reported to also cause VLM in humans although prevalence information is lacking (Bowman, 1987; Slifko et al., 2000).

The co-occurrence of *T. canis* and *T. leonina* is possible but one species tends to dominate. Age prevalence is also different in that *T. canis* is more prevalent in young dogs less than six months old while *T. leonina* tends to be more prevalent in older dogs (Okulewicz et al., 2012). This may be attributed to the somatic migration and encystation of larvae of *T. canis* which is minimal in puppies compared to older dogs and therefore puppies shed more eggs in feces. Due to the high prevalence of transplacental infection and development of patent adult infections of *T. canis* in puppies, contamination of the environment tends to be high and attributed to infection in puppies. Infection of older puppies and adult dogs mostly results in encystations of larvae.

1.5.1.3 *Trichuris vulpis* (whipworm)

T. vulpis is found worldwide in dogs and other canids e.g. foxes and cayotes.

Unembryonated eggs measuring 70-80 x 30-40 µm are released into the environment in feces (Taylor et al., 2007). The eggs are highly resistant to desiccation and UV light and embryonate in 9-21 days depending on environmental conditions. Like *Toxocara*, they can also remain viable in the environment for years (Traversa, 2011). *T. vulpis* has a direct lifecycle like *T. trichiura* in humans. Infection occurs when the embryonated eggs with L₃ larvae are ingested and they hatch in the small intestine. The larvae penetrate the mucosa and develop further for another 2-15 days, emerge and proceed to the cecum or colon where they mature and live (Kirkova and Dinev, 2005; Traversa, 2011). Adult *T. vulpis* measure between 4.5 -7.5 cm and are shaped like a whip, with a thick posterior end and a thin long anterior end, hence the name ‘whipworm’ (Taylor et al., 2007). The adult worms possess an oral stylet with which to lacerate mucosa and capillaries as they thread their anterior ends into the superficial mucosa and feed on blood, fluids and mucosa, causing hemorrhagic typhilitis or colitis (Kirkova and Dinev, 2005; Traversa, 2011). In severe infections, bloody diarrhea, anemia, lethargy and dehydration can occur and may lead to death. The prepatent period for *T. vulpis* is 74-90 days and females lay, on average, 2,000 eggs per day (Bowman, 2009). There have been a few human cases where *T. vulpis* was implicated based on larger trichuroid egg size compared to *T. trichiura* and identification of partial adult worms, but was not identified definitively (Traversa, 2011). Therefore, *T. vulpis* is currently not considered zoonotic.

1.5.1.4 *Cystoisospora* spp.

Common coccidian parasites infecting dogs are *C. canis* and the *C. ohioensis*-complex (*C. ohioensis*, *C. neorivolta*, and *C. burrowsi*). Infection is by ingestion of sporulated oocysts. The sporozoites excyst in the small intestine and invade epithelial cells, where they multiply asexually (schizogony) and sexually (gametogony) and release unsporulated oocysts (Altreuther et al., 2011). The prepatent period is 8-12 days for *C. canis* and 4-10 days for *C. ohioensis* (Altreuther et al., 2011; Houk et al., 2013). In heavy infections, they cause stunting of villi and reduction in the absorptive area leading to diarrhea (Altreuther et al., 2011; Houk et al., 2013). This is more so in young puppies but infection is believed to be exacerbated by concurrent viral infection or other immunosuppressive conditions (Taylor et al., 2007). Oocysts found in feces need to be differentiated from other coccidian oocysts such as those of *Sarcocystis* spp and *Hammondia* sp. (Bowman, 2009). Oocysts of *C. canis* are the largest in the dog and measure 32-42 x 27-33 μm , while *C. ohioensis* oocysts measure 19-27 x 18-22 μm (Bowman, 2009; Houk et al., 2013). Sporulation occurs in two or more days (Altreuther et al., 2011; Houk et al., 2013). Crowding and lack of sanitation promote the spread of coccidiosis and as such should be avoided or corrected (Taylor et al., 2007).

1.5.1.5 *Dipylidium caninum*

D. caninum is the most common cestode (tapeworm) parasite of dogs, cats and other canids and is found worldwide (Molina et al., 2003; Taylor et al., 2007). The tapeworm sheds gravid proglottids that are elongate and resemble a grain of rice, with a pore opening on each margin. Eggs (oncospheres) which measure 25-50 μm contain a hexacanth embryo and are contained in egg packets with up to 30 eggs (Taylor et al.,

2007). The eggs are released actively by the egg packet or on its disintegration. Fleas (*Ctenocephalides* spp, *Pulex irritans*) and lice (*Trichodectes canis*) serve as intermediate hosts. The eggs are ingested by the intermediate host and in the hemocoel develop into cysticercoids. Development in the louse takes about 30 days while in the flea, it can take several months. The final host is infected upon ingestion of the intermediate host containing the cysticercoids and the prepatent period is approximately three weeks (Okulewicz et al., 2012). *D. caninum* has been found to infect man occasionally, especially young children that become infected by accidental ingestion of the intermediate host (Molina et al., 2003). The cysticercoid matures in the small intestine, attaches to the mucosa by its scolex and develops a neck which buds off immature segments that develop to mature and then gravid segments in a long strobila. The adult tapeworm can measure up to 60 cm and is non-pathogenic, although the shed proglottids may cause discomfort as they crawl around the perineum (Taylor et al., 2007).

1.5.1.6 *Giardia duodenalis*

G. duodenalis is also known as *G. intestinalis* or *G. lamblia*. It is a flagellated protozoan parasite of dogs, cats and other mammals including humans (Taylor et al., 2007; Thompson, 2004). It is found worldwide and is transmitted fecal-orally. It has a simple and direct lifecycle, with trophozoites dividing by binary fission to form other trophozoites. Trophozoites encyst, are passed in feces and are fully infective (Taylor et al., 2007). The cysts measure 8-12 x 7-10 μm and are resistant to UV light, chlorine and iodine (Jarroll and Hoff, 1988). *G. duodenalis* may cause a variable semiformed mucoid, blood flecked to projectile diarrhea in dogs resulting in lethargy and weight loss. Species cross transmission is controversial as to whether dogs serve as a reservoir for human

infection (Bowman, 1987). Dogs become infected by direct fecal-oral contact from other dogs or the environment. Water contamination has led to outbreaks in humans although they can also get infected from inadvertently coming into contact with feces accompanied by poor hand hygiene (Shields et al., 2008; Thompson, 2004). Human to human transmission also occurs and it is quite common in child day-care facilities (Thompson, 2000).

1.5.2 Epidemiology of Canine Intestinal Parasites.

Intestinal parasites occur worldwide although there may be some differences in species based on the region of the world. The susceptibility of a parasite to environmental conditions may also limit its distribution. For instance among hookworm species, *U. stenocephala* has higher tolerance for low temperatures than *A. caninum* (Balasingam, 1964). Intestinal parasites occur both in urban and rural areas, with prevalence in rural areas tending to be generally higher (Bwalya et al., 2011; Dubná et al., 2007). However, among the ascarids, *T. canis* was found to contaminate soil more in urban areas than in rural areas, while prevalence was not significantly different, but prevalence with *T. leonina* was higher in rural areas (Okulewicz et al., 2012). However, the greatest influence on parasitism is how well cared for a dog is. Dogs that receive regular veterinary care and live indoors are less likely to harbor intestinal parasites as opposed to those that rarely receive veterinary care and are free roaming (Little et al., 2009). Studies in shelter dogs have shown that parasitism is extensive in this population (Blagburn et al., 1996; Papini et al., 2005). A big proportion of shelter dogs are owner surrendered, implying that the environment they are coming from is highly contaminated with parasites. Shelters may keep dogs for several months without adequate antiparasitic

therapy, thus allowing these dogs to continue shedding parasite eggs/oocysts/cysts in the premises of the animal shelter. Such dogs serve as source of infection or re-infection for newly acquired dogs in the shelter (Dubná et al., 2007). Overcrowding and poor sanitation are particularly important in enhancing the spread of coccidiosis and giardiasis (Dubná et al., 2007; Papini et al., 2005; Taylor et al., 2007; Thompson, 2004). Feeding of wet commercial food in a shelter was also found to be a contributing factor to high prevalence of giardiasis in Rome (Papini et al., 2005).

Within the USA, national surveys have been conducted on shelter dogs and owned (pet) dogs (Table 1.3) in the recent past. As has been seen in other parts of the world, shelter dogs in the USA have a high prevalence of intestinal parasites compared to pet dogs (Blagburn et al., 1996; Blagburn et al., 2012; Little et al., 2009). The most prevalent parasite in shelter dogs was hookworm, while in pet dogs, it was the protozoan parasites. This may be due to the fact that pet dogs receive anthelmintic treatment often as part of the heartworm prevention, and this helps to keep the helminth infections in check. However, in the southeastern part of the country, hookworm was the most prevalent even among pet dogs, with protozoan parasites a close second. This situation may be attributed to the highly conducive tropical environmental conditions for free-living stages of hookworm to develop in the soil and remain infective longer than they would in the cooler and drier parts of the country. In a study that was conducted among dogs brought to the veterinary teaching hospital at Louisiana State University 30 years ago (Hoskins et al., 1982), the prevalence of parasites was high, comparable to the national study on shelter dogs in 2012 for the southeastern region (Blagburn et al., 2012). It is evident that pet dogs are in a much better state with regard to parasitism now than they were 30 years

Table 1.3. Prevalence of common dog intestinal parasites from national and regional surveys in shelter and pet dogs.

	Shelter dogs 1996 (Blagburn et al., 1996)		Shelter dogs 2012 (Blagburn et al., 2012)	Pet dogs (Little et al., 2009)		Owned dogs (Hoskins et al., 1982)
	National %	Southeastern %	National %	National %	Southeastern %	Louisiana %
Hookworm	19.2	36.5	30.1	2.5	4.0	38.5
<i>T. canis</i>	14.5	17.7	12.5	2.2	1.6	8.5
<i>T. vulpis</i>	14.3	19.9	18.6	1.2	1.5	14.9
<i>Cystoisospora</i> <i>spp.</i>	4.8	5.7	<i>C. canis</i> 4.8, <i>C. ohioensis</i> 8.3	4.4	3.0	2.8
<i>Giardia spp.</i>	0.6	0.7	3.4	4.0	2.3	0.8
<i>D. caninum</i>	0.1	0.2	0.9			2.2

ago and it can be attributed to better veterinary care, better anthelmintics, especially since the discovery of ivermectin and its use as heartworm preventative (Bogan and Duncan, 1984; Kreeger et al., 1990). The national prevalence of parasites in shelter dogs has not changed much in the last 20 years which highlights the fact that there is still a population of dogs or dog owners that remains beyond the reach of the veterinarian, and that calls for enhanced public education on the need for healthcare for dogs are warranted (Blagburn et al., 2012).

1.5.3 Diagnosis of Canine Gastrointestinal Parasites

Canine intestinal parasites can be diagnosed using various flotation and sedimentation methods. The most commonly used methods in the United States surveys are centrifugal sucrose or zinc sulfate flotation (Blagburn et al., 1996; Dryden et al., 2005; Liotta et al., 2012b; Little et al., 2009). These methods are good for diagnosis of nematode eggs, protozoan oocysts and cysts but not for the heavier cestode and operculated trematode eggs (Dryden et al., 2005; Liccioli et al., 2012). Zinc sulfate flotation is particularly important for the demonstration of *Giardia* cysts. Direct smears are commonly used in the clinical setting and are useful for detecting motile protozoa (Dryden et al., 2005). Sedimentation methods are useful for the detection of cestode and trematode eggs (Dryden et al., 2005) and when saline is used as a sedimentation medium, then operculated eggs of some flukes with fully developed miracidia in eggs, such as *Heterobilharzia americanum* and *Schistosoma* spp., are prevented from hatching (Baker, 2007). The Baermann sedimentation apparatus, which allow the release of larvae out of a fecal sample and their collection in water, is useful for the isolation of lung worm

(*Angiostrongylus vasorum*) and *Strongyloides stercoralis* larvae as they exit the body via the gastrointestinal tract (Baker, 2007).

Preliminary identification of *D. caninum* may be made from fresh gravid proglottids, or adhered dried proglottids ('rice granules') are typically seen around the perineum so that diagnosis is often made on physical examination. The elongate shape, presence of two reproductive organs and shed egg packets can be seen under a light microscope by making 'squash preparations' of shed proglottids (Taylor et al., 2007). Commercial ELISA assays which are highly sensitive and specific are available for the diagnosis of *G. duodenalis* (Papini et al., 2005).

The diagnosis of CLM is by visualization of pruritic, erythematous tracts, 1-5cm long, most commonly found on parts of the body that have been in contact with moist soil, such as feet, hands, buttocks and the back as well as biopsy and histology with demonstration of migrating larvae in the epidermis (Bowman, 2009; Caumes et al., 2002; Jelinek et al., 1994). Diagnosis of VLM in humans is by aided by history of exposure to puppies, serology (antibodies to *T. canis*) and eosinophilia (Won et al., 2008).

Canine hookworm species eggs are difficult to differentiate microscopically except for *U. stenocephala* which has a notably bigger size (Traub et al., 2004). Differences in size and shape have been documented although they require a trained microscopist to detect them (Lucio-Forster et al., 2012). Molecular methods have been used to diagnose intestinal parasites in dogs, as well as to differentiate hookworm species (Gasser et al., 1996; Liotta et al., 2012b; Ngui et al., 2012b; Traub et al., 2004). PCR-RFLP by ITS fragment digestion using endonucleases was developed and used to differentiate *Ancylostoma* spp

of hookworm in dogs (e Silva et al., 2006; Liotta et al., 2012b; Traub et al., 2004). The performance of four endonucleases was compared for the differentiation of *A.*

braziliense, *A. caninum* and *A. ceylanicum* and *HinfI* produced the best profiles (e Silva et al., 2006).

1.5.4 Treatment and control

T. canis infection in puppies most often results in heavy oviposition and they are therefore considered a major source of environmental contamination and should be prevented from exposure to infection (Bowman, 2009). Hookworm infection is also most severe in puppies, commonly presenting with peracute disease due to severe anemia. CDC recommends that puppies be treated every two weeks for the first two to three months of life to clear intestinal infections before egg shedding occurs, thereby reducing public health dangers of VLM and OLM (Bowman, 2009; CAPC, 2013). In dogs where high levels of somatic larvae are likely, pregnant bitches can be cleared of tissue larvae by deworming daily with an effective dewormer like fenbendazole from 40 days of pregnancy until two weeks after whelping or by 3-4 treatments with elevated doses of ivermectin or other macrocyclic lactone drugs during pregnancy. (Bowman, 2009). *A. caninum* has been reported as being resistant to pyrantel in Australia (Kopp et al., 2007) and therefore attention must be paid to correct dosing of dogs and performing post treatment fecal exams in order to prevent and watch for the possible development of resistance in other parts of the world.

When dogs are acquired from shelters, they should be examined and treated for intestinal parasites and the treatment should continue for at least three months to get rid of reactivated or migratory larvae as they mature (Bowman, 2009; Traversa, 2011). It

should be noted adult dogs which are more commonly infected with *T. trichiura* than puppies should be treated as well (Traversa, 2011).

CLM in humans is treated with topical thiabendazole (Jelinek et al., 1994), ivermectin and/or albendazole (Caumes et al., 2002).

Tapeworm infections can be treated with praziquantel or epsiprantel and in the case of *D. caninum*, this therapy should be accompanied by use of insecticides to get rid of flea vectors and their immature stages both on the animal and in the environment (Taylor et al., 2007).

Coccidiosis can be treated with sulfonamides or toltrazural (Altreuther et al., 2011; Harder and Haberkorn, 1989). The infection can be prevented or controlled by implementing good sanitation practices, isolation of shedding dogs and avoiding crowding (Taylor et al., 2007). *G. duodenalis* can be treated with benzimidazoles like fenbendazole and nitroimidazoles like metronidazole (Taylor et al., 2007). Prompt removal of feces from the environment can aid in the control of intestinal parasites.

1.6 References

- Aaron, P.S., 1950. Developmental Studies Concerning the eggs of *Ascaris lumbricoides* var. suum. American Midland Naturalist 43, 450-470.
- Akgun, Y., 1996. Intestinal obstruction caused by *Ascaris lumbricoides*. Dis Colon Rectum 39, 1159-1163.
- Albonico, M., Bickle, Q., Ramsan, M., Montresor, A., Savioli, L., Taylor, M., 2003. Efficacy of mebendazole and levamisole alone or in combination against intestinal nematode infections after repeated targeted mebendazole treatment in Zanzibar. Bulletin of the World Health Organization 81, 343-352.
- Albonico, M., Engels, D., Savioli, L., 2004. Monitoring drug efficacy and early detection of drug resistance in human soil-transmitted nematodes: a pressing public health agenda for helminth control. International Journal for Parasitology 34, 1205-1210.

- Albonico, M., Stoltzfus, R., Savioli, L., Tielsch, J., Chwaya, H., Ercole, E., Cancrini, G., 1998. Epidemiological evidence for a differential effect of hookworm species, *Ancylostoma duodenale* or *Necator americanus*, on iron status of children. *International Journal of Epidemiology* 27, 530-537.
- Allen, A.V., Ridley, D.S., 1970. Further observations on the formol-ether concentration technique for faecal parasites. *J Clin Pathol* 23, 545-546.
- Altreuther, G., Gasda, N., Schroeder, I., Joachim, A., Settje, T., Schimmel, A., Hutchens, D., Krieger, K., 2011. Efficacy of emodepside plus toltrazuril suspension (Procox® Oral Suspension for Dogs) against prepatent and patent infection with *Isospora canis* and *Isospora ohioensis*-complex in dogs. *Parasitology Research* 109, 9-20.
- Anderson, R.C., 2000. Nematode parasites of vertebrates: Their development and transmission 2nd Edition. CABI Publishing, Wallingford, UK.
- Araújo, A., Ferreira, L.F., Confalonieri, U., Chame, M., 1988. Hookworms and the peopling of America. *Cadernos de Saúde Pública* 4, 226-233.
- Araujo, C.F., Fernández, C.L., 2005. Prevalência de parasitoses intestinais na cidade de Eirunepé, Amazonas. *Rev Soc Bras Med Trop* 38.
- Bailey, J.K., Warner, P., 2010. Respiratory arrest From *Ascaris lumbricoides*. *Pediatrics* 126, e712-e715.
- Baker, D.G. 2007. Flynn's Parasites of Laboratory Animals, In: Blackwell Publishing, Ames, Iowa, 1-10.
- Balasingam, E., 1964. Comparative studies on the effects of temperature on free-living stages of *Placoconus lotoris*, *Dochmoides stenocephala* and *Ancylostoma caninum*. *Canadian Journal of Zoology* 42, 907-918.
- Barreto, M., Rodrigues, L., Silva, R., Assis, A., Reis, M., Santos, C., Blanton, R., 2000. Lower hookworm incidence, prevalence, and intensity of infection in children with a Bacillus Calmette-Guerin vaccination scar. *Journal of Infectious Diseases* 182, 1800-1803.
- Basuni, M., Muhi, J., Othman, N., Verweij, J.J., Ahmad, M., Miswan, N., Rahumatullah, A., Aziz, F.A., Zainudin, N.S., Noordin, R., 2011. A Pentaplex Real-Time Polymerase Chain Reaction Assay for Detection of Four Species of Soil-Transmitted Helminths. *The American Journal of Tropical Medicine and Hygiene* 84, 338-343.
- Beaver, P.C., 1969. The nature of visceral larva migrans. *The Journal of parasitology*, 3-12.

- Beaver, P.C., 1988. Light, long-lasting *Necator* infection in a volunteer. *Am J Trop Med Hyg* 39, 369-372.
- Bergquist, R., Rinaldi, L., 2010. Health research based on geospatial tools: a timely approach in a changing environment. *J Helminthol* 84, 1-11.
- Bethony, J., Brooker, S., Albonico, M., Geiger, S.M., Loukas, A., Diemert, D., Hotez, P.J., 2006. Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *The Lancet* 367, 1521-1532.
- Blagburn, B.L., Lindsay, D.S., L., V.J., S., R.N., C., W.J., 1996. Prevalence of canine parasites based on fecal flotation. *The compendium on continuing education of the practising vet* 18, 483-508.
- Blagburn, B.L., Little, S.E., Butler, J.M., Land, T.M., Mount, J.D., Wright, J.C., Griffin, B., Hostetler, J. 2012. National prevalence of parasites in shelter dogs based on centrifugal fecal flotation. In: American Association of Veterinary Parasitologists 57th annual meeting, San Diego, CA, August 4-7, 76.
- Bogan, J.A., Duncan, J.L., 1984. Anthelmintics for dogs, cats and horses. *British Veterinary Journal* 140, 361-367.
- Booth, M., Vounatsou, P., N'Goran, E., Tanner, M., Utzinger, J., 2003. The influence of sampling effort and the performance of the Kato-Katz technique in diagnosing *Schistosoma mansoni* and hookworm co-infections in rural Cote d'Ivoire. *Parasitology* 127, 525 - 531.
- Bowman, D.D., 1987. Diagnostic morphology of four larval ascaridoid nematodes that may cause visceral larva migrans: *Toxascaris leonina*, *Baylisascaris procyonis*, *Lagochilascaris sprengi*, and *Hexameta leidy*. *The Journal of parasitology*, 1198-1215.
- Bowman, D.D., 2009. *Georgis' Parasitology for Veterinarians*, 9th Edition. W. B. Saunders Elsevier, St. Louis, MO.
- Bowman, D.D., Montgomery, S.P., Zajac, A.M., Eberhard, M.L., Kazacos, K.R., 2010. Hookworms of dogs and cats as agents of cutaneous larva migrans. *Trends in parasitology* 26, 162-167.
- Brandelli, C.L.C., Carli, G.A.d., Macedo, A.J., Tasca, T., 2012. Intestinal parasitism and socio-environmental factors among Mbyá-Guarani indians, Porto Alegre, Rio Grande do Sul, Brazil. *Revista do Instituto de Medicina Tropical de São Paulo* 54, 119-122.
- Brooker, S., 2010. Estimating the global distribution and disease burden of intestinal nematode infections: Adding up the numbers – A review. *International Journal for Parasitology* 40, 1137-1144.

- Brooker, S., Bethony, J., Hotez, P.J. 2004. Human Hookworm Infection in the 21st Century, In: Advances in Parasitology. Academic Press, 197-288.
- Brooker, S., Clements, A., Bundy, D., 2006a. Global epidemiology, ecology and control of soil-transmitted helminth infections. *Adv Parasitol* 62, 221 - 261.
- Brooker, S., Clements, A.C.A., Bundy, D.A.P. 2006b. Global Epidemiology, Ecology and Control of Soil-Transmitted Helminth Infections, In: Simon I. Hay, A.G., David, J.R. (Eds.) *Advances in Parasitology*. Academic Press, 221-261.
- Brooker, S., Peshu, N., Warn, P.A., Mosobo, M., Guyatt, H.L., Marsh, K., Snow, R.W., 1999. The epidemiology of hookworm infection and its contribution to anaemia among pre-school children on the Kenyan coast. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 93, 240-246.
- Brown, A.S., Raff, S., Hotez, P., 2009. Soil-Transmitted Helminths: Worms Are the Most Common. *Essentials of Public Health Biology: A Guide for the Study of Pathophysiology*, 287.
- Brown, H.W., 1927. Studies on the rate of development and viability of the eggs of *Ascaris lumbricoides* and *Trichuris trichiura* under field conditions. *The Journal of Parasitology* 14, 1-15.
- Bwalya, E.C., Nalubamba, K.S., Hankanga, C., Namangala, B., 2011. Prevalence of canine gastrointestinal helminths in urban Lusaka and rural Katete Districts of Zambia. *Preventive Veterinary Medicine* 100, 252-255.
- C.D.C. 2013a. Ascariasis.
- C.D.C. 2013b. Hookworm.
- C.D.C. 2013c. Trichiasis.
- Campbell, J.E., Wynne, R.A., 2011. *Introduction to Remote Sensing*, 5th Edition. The Guilford Press, New York.
- CAPC 2013. Intestinal parasites.
- Carvalho, G.L.X.d., Moreira, L.E., Pena, J.L., Marinho, C.C., Bahia, M.T., Machado-Coelho, G.L.L., 2012. A comparative study of the TF-Test®, Kato-Katz, Hoffman-Pons-Janer, Willis and Baermann-Moraes coprologic methods for the detection of human parasitosis. *Memórias do Instituto Oswaldo Cruz* 107, 80-84.
- Caumes, E., Ly, F., Bricaire, F., 2002. Cutaneous larva migrans with folliculitis: report of seven cases and review of the literature. *British Journal of Dermatology* 146, 314-316.
- CDC 2012. Zoonotic Hookworm.

CDC 2013. *Toxocara* lifecycle.

Chandler, A.C., 1929. Hookworm Disease. Macmillan Company, New York.

Chilton, N.B., Gasser, R.B., 1999. Sequence differences in the internal transcribed spacer ribosomal DNA among four species of hookworm (Ancylostomatoidea: *Ancylostoma*). International Journal for Parasitology 29, 1971-1977.

Conlan, J.V., Khamlome, B., Vongxay, K., Elliot, A., Pallant, L., Sripa, B., Blacksell, S.D., Fenwick, S., Thompson, R.C.A., 2012. Soil-transmitted helminthiasis in Laos: a community-wide cross-sectional study of humans and dogs in a mass drug administration environment. The American Journal of Tropical Medicine and Hygiene 86, 624-634.

Cooper, E.S., Whyte-Alleng, C.A.M., Finzi-Smith, J.S., MacDonald, T.T., 1992. Intestinal nematode infections in children: the pathophysiological price paid. Parasitology 104, S91-S103.

Cringoli, G., 2006. FLOTAC®, a novel apparatus for a multivalent faecal egg count technique. Parasitologia 48, 381-384.

Cringoli, G., Rinaldi, L., Veneziano, V., Capelli, G., Scala, A., 2004. The influence of flotation solution, sample dilution and the choice of McMaster slide area (volume) on the reliability of the McMaster technique in estimating the faecal egg counts of gastrointestinal strongyles and *Dicrocoelium dendriticum* in sheep. Veterinary parasitology 123, 121-131.

Crompton, D.W.T., 2000. The public health importance of hookworm disease. Parasitology 121, 39-50.

Crompton, D.W.T., Whitehead, R.R., 1993. Hookworm infections and human iron metabolism. Parasitology 107, S137-S145.

Daash, A., Srivastava, A., Nagpal, B., Saxena, R., Gupta, S.K., 2009. Geographical information system (GIS) in decision support to control malaria—a case study of Koraput district in Orissa, India. J Vector Borne Dis 46, 72-74.

de Gruijter, J.M., van Lieshout, L., Gasser, R.B., Verweij, J.J., Brienens, E.A.T., Ziem, J.B., Yelifari, L., Polderman, A.M., 2005. Polymerase chain reaction-based differential diagnosis of *Ancylostoma duodenale* and *Necator americanus* infections in humans in northern Ghana. Tropical Medicine & International Health 10, 574-580.

de Silva, N.R., Brooker, S., Hotez, P.J., Montresor, A., Engels, D., Savioli, L., 2003. Soil-transmitted helminth infections: updating the global picture. Trends in Parasitology 19, 547-551.

- Diawara, A., Halpenny, C.M., Churcher, T.S., Mwandawiro, C., Kihara, J., Kaplan, R.M., Streit, T.G., Idaghdour, Y., Scott, M.E., Basáñez, M.-G., Prichard, R.K., 2013. Association between response to albendazole treatment and β -tubulin genotype frequencies in soil-transmitted helminths. *PLoS Negl Trop Dis* 7, e2247.
- Dold, C., Holland, C.V., 2011. *Ascaris* and ascariasis. *Microbes and Infection* 13, 632-637.
- Dongus, S., Mwakalinga, V., Kannady, K., Tanner, M., Killeen, G. 2011. Participatory mapping as a component of operational malaria vector control in Tanzania, In: *Geospatial Analysis of Environmental Health*. Springer, 321-336.
- Dryden, M.W., Payne, P.A., Ridley, R., Smith, V., 2005. Comparison of common fecal flotation techniques for the recovery of parasite eggs and oocysts. *Vet Ther* 6, 15-28.
- Dubná, S., Langrová, I., Nápravník, J., Jankovská, I., Vadlejš, J., Pekár, S., Fechtner, J., 2007. The prevalence of intestinal parasites in dogs from Prague, rural areas, and shelters of the Czech Republic. *Veterinary Parasitology* 145, 120-128.
- Dutra, L.H., Molento, M.B., Naumann, C.R.C., Biondo, A.W., Fortes, F.S., Savio, D., Malone, J.B., 2010. Mapping risk of bovine fasciolosis in the south of Brazil using geographic information systems. *Vet Parasitol* 169, 76-81.
- e Silva, L.M.C., Miranda, R.R.C., Santos, H.A., Rabelo, É.M.L., 2006. Differential diagnosis of dog hookworms based on PCR-RFLP from the ITS region of their rDNA. *Veterinary Parasitology* 140, 373-377.
- Endris, M., Tekeste, Z., Lemma, W., Kassu, A., 2013. Comparison of the Kato-Katz, Wet Mount, and Formol-Ether Concentration Diagnostic Techniques for Intestinal Helminth Infections in Ethiopia. *ISRN Parasitology* 2013, 5.
- Eppig, C., Fincher, C., Thornhill, R., 2010. Parasite prevalence and the worldwide distribution of cognitive ability. *Proc Biol Sci* 277, 3801 - 3808.
- Esrey, S.A., Potash, J.B., Roberts, L., Shiff, C., 1991. Effects of improved water supply and sanitation on ascariasis, diarrhoea, dracunculiasis, hookworm infection, schistosomiasis, and trachoma. *Bulletin of the World Health Organization* 69, 609-621.
- Ezeamama, A.E., Friedman, J.F., Olveda, R.M., Acosta, L.P., Kurtis, J.D., Mor, V., McGarvey, S.T., 2005b. Functional significance of low-intensity polyparasite helminth infections in anemia. *Journal of Infectious Diseases* 192, 2160-2170.
- Faust, E.C., D'Antoni, J.S., Odom, V., Miller, M.J., Peres, C., Sawitz, W., Thomen, L.F., Tobie, J., Walker, J.H., 1938. A critical study of clinical laboratory technics for the diagnosis of protozoan cysts and helminth eggs in feces: I. Preliminary

- Communication. The American Journal of Tropical Medicine and Hygiene s1-18, 169-183.
- Fenghua, S., Zhongxing, W., Yixing, Q., Hangqun, C., Haichou, X., Hainan, R., Shuhua, X., Bin, Z., Hawdon, J.M., Zheng, F., Hotez, P.J., 1998. Epidemiology of human intestinal nematode infections in Wujiang and Pizhou counties, Jiangsu Province, China. *Southeast Asian J Trop Med Public Health* 29, 605-610.
- Fleming, F.M., Brooker, S., Geiger, S.M., Caldas, I.R., Correa-Oliveira, R., Hotez, P.J., Bethony, J.M., 2006. Synergistic associations between hookworm and other helminth species in a rural community in Brazil. *Trop Med Int Health* 11, 56-64.
- Fuentes, M.V., Sainz-Elipse, S., Nieto, P., Malone, J.B., Mas-Coma, S., 2005. Geographical information systems risk assessment models for zoonotic fascioliasis in the South American Andes region. *Parassitologia* 47, 151-156.
- Funk, A.L., Boisson, S., Clasen, T., Ensink, J.H.J., 2013. Comparison of Kato-Katz, ethyl-acetate sedimentation, and Midi Parasep® in the diagnosis of hookworm, *Ascaris* and *Trichuris* infections in the context of an evaluation of rural sanitation in India. *Acta Tropica* 126, 265-268.
- Furmidge, B.A., Horn, L.A., Pritchard, D.I., 1996. The anti-haemostatic strategies of the human hookworm *Necator americanus*. *Parasitology* 112, 81-87.
- Gamboa, M.I., 2005. Effects of temperature and humidity on the development of eggs of *Toxocara canis* under laboratory conditions. *Journal of Helminthology* 79, 327-331.
- Garcia, L.S., 2001. *Diagnostic Medical Parasitology*, 4th Edition. ASM Press, Washington, DC.
- Gasser, R.B., Chilton, N.B., Hoste, H., Beveridge, I., 1993. Rapid sequencing of rDNA from single worms and eggs of parasitic helminths. *Nucleic Acids Research* 21, 2525-2526.
- Gasser, R.B., Stewart, L.E., Speare, R., 1996. Genetic markers in ribosomal DNA for hookworm identification. *Acta Tropica* 62, 15-21.
- Geenen, P.L., Bresciani, J., Boes, J., Pedersen, A., Eriksen, L., Fagerholm, H.P., Nansen, P., 1999. The morphogenesis of *Ascaris suum* to the infective third-stage larvae within the egg. *The Journal of Parasitology* 85, 616-622.
- Glinz, D., Silué, K.D., Knopp, S., Lohourignon, L.K., Yao, K.P., Steinmann, P., Rinaldi, L., Cringoli, G., N'Goran, E.K., Utzinger, J., 2010. Comparing Diagnostic Accuracy of Kato-Katz, Koga Agar Plate, Ether-Concentration, and FLOTAC for *Schistosoma mansoni* and Soil-Transmitted Helminths. *PLoS Negl Trop Dis* 4, e754.

- Gonçalves, J.F., Tanabe, M., Medeiros, F.d.P.d.M., Gonçalves, F.J., Aca, I.d.S., Motta, S.R.N.d., Tateno, S., Takeuchi, T., 1990. Parasitological and serological studies on amoebiasis and other intestinal parasitic infections in the rural sector around Recife, Northeast Brazil. *Rev Inst Med Trop São Paulo* 32, 428-435.
- González, P., Carbonell, E., Urios, V., Rozhnov, V.V., 2007. Coprology of *Panthera tigris altaica* and *Felis bengalensis euphilurus* From the Russian Far East. *Journal of Parasitology* 93, 948-950.
- Gross, R., Schell, B., Molina, M.C.B., Leão, M.A.C., Strack, U., 1989. The impact of improvement of water supply and sanitation facilities on diarrhea and intestinal parasites: a Brazilian experience with children in two low-income urban communities. *Revista de Saúde Pública* 23, 214-220.
- Gyapong, J.O., Kyelem, D., Kleinschmidt, I., Agbo, K., Ahouandogbo, F., Gaba, J., Owusu-Banahene, G., Sanou, S., Sodahlon, Y.K., Biswas, G., Kale, O.O., Molyneux, D.H., ROUNGOU, J.B., Thomson, M.C., Remme, J., 2002. The use of spatial analysis in mapping the distribution of bancroftian filariasis in four West African countries. *Annals of Tropical Medicine and Parasitology* 96, 695-705.
- Haque, R., Ahmed, T., Wahed, M.A., Mondal, D., Rahman, A.S., Albert, M.J., 2010. Low-dose beta-carotene supplementation and deworming improve serum vitamin A and beta-carotene concentrations in preschool children of Bangladesh. *Journal of health, population, and nutrition* 28, 230-237.
- Harada, U., Mori, O., 1955. A new method for culturing hookworm. *Yonago Acta Med* 1, 177-179.
- Harder, A., Haberkorn, A., 1989. Possible mode of action of toltrazuril: studies on two *Eimeria* species and mammalian and *Ascaris suum* enzymes. *Parasitology Research* 76, 8-12.
- Harto, M., Rodriguez-Salvador, V., Avino, J., Duch-Samper, A., Menezo, J., 1999. Diffuse unilateral subacute neuroretinitis in Europe. *Eur J Ophthalmol* 9, 58-62.
- Hawdon, J.M., 1996. Differentiation between the human hookworms *Ancylostoma duodenale* and *Necator americanus* Using PCR-RFLP. *The Journal of Parasitology* 82, 642-647.
- Horton, J., 2003. Human gastrointestinal helminth infections: are they now neglected diseases? *Trends in Parasitology* 19, 527-531.
- Hoskins, J.D., Malone, J.B., Smith, P.H., Uhl, S.A., 1982. Prevalence of parasitism diagnosed by fecal examination in Louisiana dogs. *American Journal of Veterinary Research* 43, 1106-1109.
- Hotez, P., 2008. Hookworm and Poverty. *Ann NY Acad Sci* 1136, 38-44.

- Hotez, P., Brindley, P., Bethony, J., King, C., Pearce, E., Jacobson, J., 2008. Helminth infections: the great neglected tropical diseases. *J Clin Invest* 118, 1311 - 1321.
- Hotez, P., Molyneux, D., Fenwick, A., Kumaresan, J., Erlich Sachs, S., Sachs, J., Savioli, L., 2007. Control of neglected tropical diseases. *N Engl J Med* 357, 1018 - 1027.
- Hotez, P.J., Bethony, J., Bottazzi, M.E., Brooker, S., Buss, P., 2005. Hookworm: "The Great Infection of Mankind". *PLoS Med* 2, e67.
- Hotez, P.J., Bundy, D.A., Beegle, K., Brooker, S., Drake, L., de Silva, N., Montresor, A., Engels, D., Jukes, M., Chitsulo, L., 2006. Helminth infections: soil-transmitted helminth infections and schistosomiasis.
- Hotez, P.J., Cerami, A., 1983. Secretion of a proteolytic anticoagulant by *Ancylostoma* hookworms. *The Journal of Experimental Medicine* 157, 1594-1603.
- Houk, Alice E., O'Connor, T., Pena, Hilda F.J., Gennari, Solange M., Zajac, Anne M., Lindsay, David S., 2013. Experimentally Induced Clinical *Cystoisospora canis* Coccidiosis in Dogs with Prior Natural Patent *C. ohioensis*-like or *C. canis* Infections. *Journal of Parasitology*.
- Hu, M., Chilton, N.B., Gasser, R.B., 2002. The mitochondrial genomes of the human hookworms, *Ancylostoma duodenale* and *Necator americanus* (Nematoda: Secernentea). *International Journal for Parasitology* 32, 145-158.
- Humphries, D., Simms, B.T., Davey, D., Otchere, J., Quagrain, J., Terryah, S., Newton, S., Berg, E., Harrison, L.M., Boakye, D., Wilson, M., Cappello, M., 2013. Hookworm Infection among School Age Children in Kintampo North Municipality, Ghana: Nutritional Risk Factors and Response to Albendazole Treatment. *The American Journal of Tropical Medicine and Hygiene* 89, 540-548.
- Ismail, M.M., Jayakody, R.L., 1999. Efficacy of albendazole and its combinations with ivermectin or diethylcarbamazine (DEC) in the treatment of *Trichuris trichiura* infections in Sri Lanka. *Annals of Tropical Medicine and Parasitology* 93, 501-504.
- Jardim-Botelho, A., Raff, S., DeÁvila Rodrigues, R., Hoffman, H.J., Diemert, D.J., Corrêa-Oliveira, R., Bethony, J.M., Gazzinelli, M.F., 2008. Hookworm, *Ascaris lumbricoides* infection and polyparasitism associated with poor cognitive performance in Brazilian schoolchildren. *Tropical Medicine & International Health* 13, 994-1004.
- Jarroll, E.L., Hoff, J.C., 1988. Effect of disinfectants on *Giardia* cysts. *Critical Reviews in Environmental Control* 18, 1-28.
- Jeffery, G.M., 1960. A three-year epidemiologic study of intestinal parasites in a selected group of mental patients. *American Journal of Hygiene* 71, 1-8.

- Jelinek, T., Maiwald, H., Nothdurft, H., Löscher, T., 1994. Cutaneous larva migrans in travelers: synopsis of histories, symptoms, and treatment of 98 patients. *Clinical infectious diseases* 19, 1062-1066.
- Jones, L., 2012. Geographic information systems. *Essentials of Environmental Epidemiology for Health Protection: A Handbook for Field Professionals*, 121.
- Jonker, F.A.M., Calis, J.C.J., Phiri, K., Brienens, E.A.T., Khoffi, H., Brabin, B.J., Verweij, J.J., van Hensbroek, M.B., van Lieshout, L., 2012. Real-time PCR demonstrates *Ancylostoma duodenale* is a key factor in the etiology of severe anemia and iron deficiency in Malawian pre-school children. *PLoS Negl Trop Dis* 6, e1555.
- Kasper, G., Brown, A., Eberl, M., Vallar, L., Kieffer, N., Berry, C., Girdwood, K., Eggleton, P., Quinnell, R., Pritchard, D.I., 2001. A calreticulin-like molecule from the human hookworm *Necator americanus* interacts with C1q and the cytoplasmic signalling domains of some integrins. *Parasite Immunology* 23, 141-152.
- Katz, N., Chaves, A., Pellegrino, J., 1972. A simple device for quantitative stool thick-smear technique in schistosomiasis mansoni. *Rev Inst Med Trop São Paulo* 14, 397 - 400.
- Keiser, J., Tritten, L., Silbereisen, A., Speich, B., Adelfio, R., Vargas, M., 2013. Activity of oxantel pamoate monotherapy and combination chemotherapy against *Trichuris muris* and hookworms: revival of an old drug. *PLoS neglected tropical diseases* 7, e2119.
- Keiser, J., Utzinger, J., 2008. Efficacy of current drugs against soil-transmitted helminth infections: Systematic review and meta-analysis. *JAMA* 299, 1937-1948.
- King, C.H., 2010. Parasites and poverty: The case of schistosomiasis. *Acta Tropica* 113, 95-104.
- Kirkova, Z., Dinev, I., 2005. Morphological changes in the intestine of dogs, experimentally infected with *Trichuris vulpis*. *Bulgarian Journal of Veterinary Medicine* 8, 239-243.
- Knopp, S., Mohammed, K.A., Speich, B., Hattendorf, J., Khamis, I.S., Khamis, A.N., Stothard, J.R., Rollinson, D., Marti, H., Utzinger, J., 2010. Albendazole and mebendazole administered alone or in combination with ivermectin against trichuris trichiura: a randomized controlled Trial. *Clinical Infectious Diseases* 51, 1420-1428.
- Knopp, S., Rinaldi, L., Khamis, I.S., Stothard, J.R., Rollinson, D., Maurelli, M.P., Steinmann, P., Marti, H., Cringoli, G., Utzinger, J., 2009. A single FLOTAC is more sensitive than triplicate Kato-Katz for the diagnosis of low-intensity soil-transmitted helminth infections. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 103, 347-354.

- Kopp, S.R., Kotze, A.C., McCarthy, J.S., Coleman, G.T., 2007. High-level pyrantel resistance in the hookworm *Ancylostoma caninum*. *Veterinary parasitology* 143, 299-304.
- Kreeger, T.J., Seal, U.S., Callahan, M., Beckel, M., 1990. Treatment and Prevention with Ivermectin of Dirofilariosis and Ancylostomiasis in Captive Gray Wolves (*Canis lupus*). *Journal of Zoo and Wildlife Medicine* 21, 310-317.
- Landmann, J.K., Prociv, P., 2003. Experimental human infection with the dog hookworm, *Ancylostoma caninum*. *Medical journal of Australia* 178, 69-71.
- Levecke, B., Behnke, J.M., Ajjampur, S.S., Albonico, M., Ame, S.M., Charlier, J., Geiger, S.M., Hoa, N.T., Kamwa Ngassam, R.I., Kotze, A.C., McCarthy, J.S., Montresor, A., Periago, M.V., Roy, S., Tchuem Tchuente, L.A., Thach, D.T., Vercruysse, J., 2011. A comparison of the sensitivity and fecal egg counts of the McMaster egg counting and Kato-Katz thick smear methods for soil-transmitted helminths. *PLoS neglected tropical diseases* 5, e1201.
- Levecke, B., De Wilde, N., Vandenhouste, E., Vercruysse, J., 2009. Field Validity and Feasibility of Four Techniques for the Detection of *Trichuris* in Simians: A Model for Monitoring Drug Efficacy in Public Health? *PLoS Negl Trop Dis* 3, e366.
- Liccioli, S., Catalano, S., Kutz, S.J., Lejeune, M., Verocai, G.G., Duignan, P.J., Fuentealba, C., Ruckstuhl, K.E., Massolo, A., 2012. Sensitivity of double centrifugation sugar fecal flotation for detecting intestinal helminths in coyotes (*Canis latrans*). *Journal of Wildlife Diseases* 48, 717-723.
- Lindsay, S.W., Thomas, C.J., 1999. Mapping and estimating the population at risk from lymphatic filariasis in Africa. *Trans R Soc Trop Med Hyg* 94, 37-45.
- Liotta, J.L., Lee, A.C., Aksel, S., Alkhalife, I., Cruz-Reyes, A., Youn, H., Bienhoff, S.E., Bowman, D.D., 2012a. Obtaining an isolate of *Ancylostoma braziliense* from dogs without the need for necropsy. *Journal of Parasitology* 98, 1034-1036.
- Liotta, J.L., Youn, H., Aksel, S., Bienhoff, S.E., Bowman, D.D., 2012b. Prevalence of *Ancylostoma braziliense* in Dogs From Alachua and Marion Counties, Florida, United States. *Journal of Parasitology* 98, 1039-1040.
- Little, S.E., Johnson, E.M., Lewis, D., Jaklitsch, R.P., Payton, M.E., Blagburn, B.L., Bowman, D.D., Moroff, S., Tams, T., Rich, L., Aucoin, D., 2009. Prevalence of intestinal parasites in pet dogs in the United States. *Veterinary Parasitology* 166, 144-152.
- Looss, A., 1911. The anatomy and life history of *Ancylostoma duodenale*. *Records of the Egyptian Government School of Medicine* 4, 167-616.
- Lucio-Forster, A., Liotta, J.L., Yaros, J.P., Briggs, K.R., Mohammed, H.O., Bowman, D.D., 2012. Morphological differentiation of eggs of *Ancylostoma caninum*,

- Ancylostoma tubaeforme*, and *Ancylostoma braziliense* from dogs and cats in the United States. *Journal of Parasitology* 98, 1041-1044.
- MAFF, 1986. *Manual of Veterinary Parasitological Laboratory Techniques: Fisheries and Food Reference Book*, Vol 418. HMSO, London.
- Mahalanabis, D., Jalan, K.N., Maitra, T.K., Agarwal, S.K., 1976. Vitamin A absorption in ascariasis. *Am J Clin Nutr* 29, 1372-1375.
- Malone, J.B., 2005. Biology-based mapping of vector-borne diseases by geographic information systems and remote sensing. *Parasitologia* 47, 27-50.
- Malone, J.B., Williams, T.E., A., M.R., P., G.J., F., L.A., 1987. Fascioliasis in cattle in Louisiana: development of a system to predict disease risk by climate, using the Thornthwaite water budget. *American Journal of Veterinary Research* 48, 1167-1170.
- Malone, J.B., Yilma, J.M., McCarroll, J.C., Erko, B., Mukaratirwa, S., Zhou, X., 2001. Satellite climatology and the environmental risk of *Schistosoma mansoni* in Ethiopia and east Africa. *Acta Tropica* 79, 59-72.
- Martin, R., 1997. Modes of action of anthelmintic drugs. *The Veterinary Journal* 154, 11-34.
- McMaster, G.S., Wilhelm, W.W., 1997. Growing degree-days: one equation, two interpretations. *Agricultural and Forest Meteorology* 87, 291-300.
- Miller, T.A., 1971. Vaccination against the canine hookworm diseases. *Adv Parasitol* 9, 153-183.
- Mischler, P., Kearney, M., McCarroll, J.C., Scholte, R.G., Vounatsou, P., Malone, J.B., 2012. Environmental and socio-economic risk modelling for Chagas disease in Bolivia. *Geospatial Health* 6, S59-S66.
- Molina, C.P., Ogburn, J., Adegboyega, P., 2003. Infection by *Dipylidium caninum* in an Infant. *Archives of Pathology & Laboratory Medicine* 127, e157-e159.
- Montresor, A., Crompton, D.W.T., Hall, H., Bundy, D.A.P., Savioli, L. 1998. Guidelines for the evaluation of soil-transmitted helminthiasis and schistosomiasis at community level. A guide for control programme managers. (Geneva, World Health Organisation.).
- Mudenda, N.B., Malone, J.B., Kearney, M.T., Mischler, P., Nieto, P., McCarroll, J.C., Vounatsou, P., 2012. Modelling the ecological niche of hookworm in Brazil based on climate. *Geospatial Health* 6, 111-123.
- Murray, C.J.L., Vos, T., Lozano, R., Naghavi, M., Flaxman, A.D., Michaud, C., Ezzati, M., Shibuya, K., Salomon, J.A., Abdalla, S., al.et, 2012. Disability-adjusted life

- years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *The Lancet* 380, 2197-2223.
- Nawalinski, T.A., Schad, G.A., 1974. Arrested development in *Ancylostoma duodenale*: Course of a self-induced infection in man. *The American Journal of Tropical Medicine and Hygiene* 23, 895-898.
- Ngui, R., Ching, L.S., Kai, T.T., Roslan, M.A., Lim, Y.A.L., 2012a. Molecular Identification of Human Hookworm Infections in Economically Disadvantaged Communities in Peninsular Malaysia. *The American Journal of Tropical Medicine and Hygiene* 86, 837-842.
- Ngui, R., Lim, Y.A., Traub, R., Mahmud, R., Mistam, M.S., 2012b. Epidemiological and genetic data supporting the transmission of *Ancylostoma ceylanicum* among human and domestic animals. *PLoS neglected tropical diseases* 6, e1522.
- Nichols, R.L., 1956. The etiology of visceral Larva migrans. II. Comparative larval morphology of *Ascaris lumbricoides*, *Necator americanus*, *Strongyloides stercoralis* and *Ancylostoma caninum*. *Journal of Parasitology* 42, 363-399.
- Nieto, P., Malone, J.B., Bavia, M.E., 2006. Ecological niche modeling for visceral leishmaniasis in the state of Bahia, Brazil, using genetic algorithm for rule-set prediction and growing degree day-water budget analysis. *Geospatial Health* 1, 115-126.
- Okazaki, M., Okazaki, M., Miranda, P., Neto, J., Diegues, V., Alves, J., Cauas, M., Tanabe, M., Kobayashi, S., Kaneko, N., 1988. Parasitological and serological studies on amoebiasis and other intestinal parasitic infections in Recife and its suburban area, Northeast Brazil. *Revista do Instituto de Medicina Tropical de São Paulo* 30, 313-321.
- Okulewicz, A., Perek-Matysiak, A., Buńkowska, K., Hildebrand, J., 2012. *Toxocara canis*, *Toxocara cati* and *Toxascaris leonina* in wild and domestic carnivores. *Helminthologia* 49, 3-10.
- Olsen, L.S., Kelley, G.W., Sen, H.G., 1958. Longevity and egg-production of *Ascaris suum*. *Transactions of the American Microscopical Society* 77, 380-383.
- Papini, R., Gorini, G., Spaziani, A., Cardini, G., 2005. Survey on giardiasis in shelter dog populations. *Veterinary Parasitology* 128, 333-339.
- Pawlowski, Z.S., Schad, G., Stott, G., 1991. Hookworm infection and anaemia: approaches to prevention and control.
- Pit, D., Polderman, A., Baeta, S., Schulz-Key, H., Soboslay, P., 2001. Parasite-specific antibody and cellular immune responses in humans infected with *Necator americanus* and *Oesophagostomum bifurcum*. *Parasitology Research* 87, 722-729.

- Pit, D.S., Rijcken, F.E., Raspoort, E.C., Baeta, S.M., Polderman, A.M., 1999. Geographic distribution and epidemiology of *Oesophagostomum bifurcum* and hookworm infections in humans in Togo. *The American Journal of Tropical Medicine and Hygiene* 61, 951-955.
- Polderman, A., Krepel, H., Verweij, J., Baeta, S., Rotmans, J., 1993. Serological diagnosis of *Oesophagostomum* infections. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 87, 433-435.
- Prociv, P., Croese, J., 1996. Human enteric infection with *Ancylostoma caninum*: hookworms reappraised in the light of a “new” zoonosis. *Acta Tropica* 62, 23-44.
- Pullan, R.L., Brooker, S.J., 2012. The global limits and population at risk of soil-transmitted helminth infections in 2010. *Parasites & Vectors* 5, 81.
- Romstad, A., Gasser, R.B., Monti, J.R., Polderman, A.M., Nansen, P., Pit, D.S.S., Chilton, N.B., 1997. Differentiation of *Oesophagostomum bifurcum* from *Necator americanus* by PCR using genetic markers in spacer ribosomal DNA. *Molecular and Cellular Probes* 11, 169-176.
- Salahi-Moghaddam, A., Mohebbi, M., Moshfaei, A., Majid, H., Zarei, Z., 2010. Ecological study and risk mapping of visceral leishmaniasis in an endemic area of Iran based on a geographical information systems approach. *Geospatial Health* 5, 71-77.
- Santos, F.L.N., Souza, A.M.G.C.d., Soares, N.M., 2013. Hookworm and threadworm infections and their association with hemoglobin and eosinophil concentrations in residents of Salvador-Bahia, Brazil. *Revista do Instituto de Medicina Tropical de São Paulo* 55, 233-238.
- Sato, M., Sanguankiat, S., Yoonuan, T., Pongvongsa, T., Keomoungkhoun, M., Phimmayoi, I., Boupa, B., Moji, K., Waikagul, J., 2010. Copro-molecular identification of infections with hookworm eggs in rural Lao PDR. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 104, 617-622.
- Savioli, L., Albonico, M., 2004. Focus: Soil-transmitted helminthiasis. *Nat Rev Micro* 2, 618-619.
- Schad, G.A., Page, M.R., 1982. *Ancylostoma caninum*: Adult worm removal, corticosteroid treatment, and resumed development of arrested larvae in dogs. *Experimental Parasitology* 54, 303-309.
- Schad, G.A., Warren, K.S. 1990. *Hookworm Disease, Current Status and New Directions* (Taylor & Francis Ltd.).
- Shakya, B., Rai, S.K., Singh, A., Shrestha, A., 2006. Intestinal parasitosis among the elderly people in Kathmandu Valley.

- Shields, J.M., Gleim, E.R., Beach, M.J., 2008. Prevalence of *Cryptosporidium* spp. and *Giardia intestinalis* in swimming pools, Atlanta, Georgia. *Emerging infectious diseases* 14, 948.
- Shuttleworth, W.J. 1983. Evaporation Models in the Global Water Budget, In: Street-Perrott, A., Beran, M., Ratcliffe, R. (Eds.) *Variations in the Global Water Budget*. Springer Netherlands, 147-171.
- Slifko, T.R., Smith, H.V., Rose, J.B., 2000. Emerging parasite zoonoses associated with water and food. *International Journal for Parasitology* 30, 1379-1393.
- Smout, F.A., Thompson, R.C.A., Skerratt, L.F., 2013. First report of *Ancylostoma ceylanicum* in wild canids. *International Journal for Parasitology: Parasites and Wildlife* 2, 173-177.
- Stassens, P., Bergum, P.W., Gansemans, Y., Jespers, L., Laroche, Y., Huang, S., Maki, S., Messens, J., Lauwereys, M., Cappello, M., 1996. Anticoagulant repertoire of the hookworm *Ancylostoma caninum*. *Proceedings of the National Academy of Sciences* 93, 2149-2154.
- Stephenson, L., Latham, M., Adams, E., Kinoti, S., Pertet, A., 1993. Physical fitness, growth and appetite of Kenyan school boys with hookworm, *Trichuris trichiura* and *Ascaris lumbricoides* infections are improved four months after a single dose of albendazole. *J Nutr* 123, 1036 - 1046.
- Stoltzfus, R.J., Dreyfuss, M.L., Chwaya, H.M., Albonico, M., 1997. Hookworm control as a strategy to prevent iron deficiency. *Nutrition Reviews* 55, 223-332.
- Stone, W.M., Girardeau, M., 1968. Transmammary passage of *Ancylostoma caninum* larvae in dogs. *The Journal of parasitology*, 426-429.
- Suwannatrai, A., Suwannatrai, S., Haruay, S., Piratae, S., Thammasiri, C., Khamphooa, P., Kulsantiwong, J., Prasopdee, S., Tarbsripair, P., Suwanwerakamtorn, R., 2011. Effect of soil surface salt on the density and distribution of the snail *Bithynia siamensis goniomphalos* in northeast Thailand. *Geospatial Health* 5, 183-190.
- Svensson, R., 1925. Observations on the development and longevity of hookworm larvae in different temperature conditions. *Chinese Medical Journal* 39, 667-673.
- Svensson, R.M., Kessel, J.F., 1926. Morphological Differences between *Necator* and *Ancylostoma* Larvae. *The Journal of Parasitology* 13, 146-153.
- Tarafder, M.R., Carabin, H., Joseph, L., Balolong Jr, E., Olveda, R., McGarvey, S.T., 2010. Estimating the sensitivity and specificity of Kato-Katz stool examination technique for detection of hookworms, *Ascaris lumbricoides* and *Trichuris trichiura* infections in humans in the absence of a 'gold standard'. *International Journal for Parasitology* 40, 399-404.

- Taylor, M.A., Coop, R.L., Wall, R.L., 2007. *Veterinary Parasitology*, 3rd Edition. Blackwell Publishing, Oxford, UK.
- Thompson, R., 2004. Epidemiological and molecular evidence supports the zoonotic transmission of *Giardia* among humans and dogs living in the same community.
- Thompson, R.C.A., 2000. Giardiasis as a re-emerging infectious disease and its zoonotic potential. *International Journal for Parasitology* 30, 1259-1267.
- Thomson, M.C., Connor, S.J., Milligan, P., Flasse, S.P., 1997. Mapping malaria risk in Africa: what can satellite data contribute? *Parasitol Today* 13, 313-318.
- Traub, R.J., Inpankaew, T., Sutthikornchai, C., Sukthana, Y., Thompson, R.C.A., 2008. PCR-based coprodiagnostic tools reveal dogs as reservoirs of zoonotic ancylostomiasis caused by *Ancylostoma ceylanicum* in temple communities in Bangkok. *Veterinary Parasitology* 155, 67-73.
- Traub, R.J., Robertson, I.D., Irwin, P., Mencke, N., Thompson, R., 2004. Application of a species-specific PCR-RFLP to identify *Ancylostoma* eggs directly from canine faeces. *Veterinary Parasitology* 123, 245-255.
- Traversa, D., 2011. Are we paying too much attention to cardio-pulmonary nematodes and neglecting old-fashioned worms like *Trichuris vulpis*. *Parasit Vectors* 4, 32.
- Truant, A.L., Elliott, S.H., Kelly, M.T., Smith, J.H., 1981. Comparison of formalin-ethyl ether sedimentation, formalin-ethyl acetate sedimentation, and zinc sulfate flotation techniques for detection of intestinal parasites. *Journal of Clinical Microbiology* 13, 882-884.
- Udonsi, J.K., Atata, G., 1987. *Necator americanus*: Temperature, pH, light, and larval development, longevity, and desiccation tolerance. *Exp Parasitol* 63, 136-142.
- Utzinger, J., Rinaldi, L., Lohourignon, L.K., Rohner, F., Zimmermann, M.B., Tschannen, A.B., N'Goran, E.K., Cringoli, G., 2008. FLOTAC: a new sensitive technique for the diagnosis of hookworm infections in humans. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 102, 84-90.
- Valencia-López, N., Malone, J.B., Carmona, C.G., Velásquez, L.E., 2012. Climate-based risk models for *Fasciola hepatica* in Colombia. *Geospat Health* 6, S67-S85.
- Vercruysse, J., Albonico, M., Behnke, J.M., Kotze, A.C., Prichard, R.K., McCarthy, J.S., Montresor, A., Levecke, B., 2011. Is anthelmintic resistance a concern for the control of human soil-transmitted helminths? *International Journal for Parasitology: Drugs and Drug Resistance* 1, 14-27.
- Verweij, J.J., Pit, D.S.S., Van Lieshout, L., Baeta, S.M., Dery, G.D., Gasser, R.B., Polderman, A.M., 2001. Determining the prevalence of *Oesophagostomum*

- bifurcum* and *Necator americanus* infections using specific PCR amplification of DNA from faecal samples. Tropical Medicine & International Health 6, 726-731.
- Villamizar, E., Mendez, M., Bonilla, E., Varon, H., de Onatra, S., 1996. *Ascaris lumbricoides* infestation as a cause of intestinal obstruction in children: experience with 87 cases. J Pediatr Surg 31, 201-204; discussion 204-205.
- WHA 2001. Resolution WHA54.19: Schistosomiasis and soil-transmitted helminth infections.
- WHO, 2008. Soil-transmitted helminthiasis. Progress report on number of children treated with anthelmintic drugs: an update towards the 2010 global target. Relevé épidémiologique hebdomadaire / Section d'hygiène du Secrétariat de la Société des Nations = Weekly epidemiological record / Health Section of the Secretariat of the League of Nations 82, 237-252.
- WHO 2010. Working to overcome the global impact of neglected tropical diseases (Geneva).
- Won, K.Y., Kruszon-Moran, D., Schantz, P.M., Jones, J.L., 2008. National Seroprevalence and Risk Factors for Zoonotic *Toxocara* spp. Infection. The American Journal of Tropical Medicine and Hygiene 79, 552-557.
- Wu, Z.X., Peng, J.M., 1965. Studies on the morphological differentiation of the infective larvae of *Ancylostoma duodenale* and *Necator americanus*. Acta Parasitol Sin 2, 280-290.
- Yang, G.-J., Gao, Q., Zhou, S.S., Malone, J.B., McCarroll, J.C., M., T., Vounatsou, P., Bergquist, R., Utzinger, J., Zhou, X.-N., 2010. Mapping and predicting malaria transmission in the People's Republic of China, using integrated biology-driven and statistical models. Geospatial Health 5, 11-22.
- Yap, P., Du, Z.-W., Chen, R., Zhang, L.-P., Wu, F.-W., Wang, J., Wang, X.-Z., Zhou, H., Zhou, X.-N., Utzinger, J., Steinmann, P., 2012. Soil-transmitted helminth infections and physical fitness in school-aged Bulang children in southwest China: results from a cross-sectional survey. Parasites & Vectors 5, 50.
- Yong, T.-S., Lee, J.-H., Sim, S., Lee, J., Min, D.-Y., Chai, J.-Y., Eom, K.S., Sohn, W.-M., Lee, S.-H., Rim, H.-J., 2007b. Differential diagnosis of *Trichostrongylus* and hookworm eggs via PCR using ITS-1 sequence. The Korean journal of parasitology 45, 69-74.
- Young, K.H., Bullock, S.L., Melvin, D.M., Spruill, C.L., 1979. Ethyl acetate as a substitute for diethyl ether in the formalin-ether sedimentation technique. Journal of Clinical Microbiology 10, 852-853.

- Zajac, A.M., Johnson, J., King, S.E., 2002. Evaluation of the Importance of Centrifugation as a Component of Zinc Sulfate Fecal Flotation Examinations. *Journal of the American Animal Hospital Association* 38, 221-224.
- Zhou, X.N., Malone, J.B., Kristensen, T.K., Bergquist, N.R., 2001. Application of geographic information systems and remote sensing to schistosomiasis control in China. *Acta Trop* 79, 97-106.
- Ziegelbauer, K., Speich, B., Mäusezahl, D., Bos, R., Keiser, J., Utzinger, J., 2012. Effect of Sanitation on Soil-Transmitted Helminth Infection: Systematic Review and Meta-Analysis. *PLoS Med* 9, e1001162.

2 MODELING THE ECOLOGICAL NICHE OF HOOKWORM IN BRAZIL BASED ON CLIMATE

2.1 Introduction

Hookworm infection is a worldwide public health problem, particularly in developing countries in Asia, Africa and Latin America, where poverty levels are high. An estimated 740 million people are infected (WHO, 2006). Of great concern in hookworm infection is the resultant iron deficiency anemia due to intestinal blood loss, especially in children and women of reproductive age and also the reduced cognitive ability of infected children (Hotez, 2008; Stoltzfus et al., 1996). Brazil introduced a Schistosomiasis Control Program in 1976, which is an active decentralized, federal program in known endemic areas (Amaral et al., 2006). Since the Kato-Katz method utilized in this program for the detection of *Schistosoma* eggs also shows the presence of many other helminth eggs, there is a bonus in using the results with respect to other helminth infections. Studies have shown that co-infection between *Schistosoma* sp. and soil transmitted helminths (STH) is common, including hookworm, *Ascaris lumbricoides* and *Trichuris trichiura* (Booth et al., 2003; Fleming et al., 2006).

Human hookworm infections are caused by *Necator americanus* and *Ancylostoma duodenale*. In Brazil, hookworm infections are predominantly caused by *N. americanus* (Fleming et al., 2006; Geiger et al., 2004). The eggs of this species do not develop at temperatures below 15°C. The larvae prefer shady, moist areas with temperatures at or above 30°C but are killed above 45°C (Udonsi and Atata, 1987). Larvae fare best in sandy loam soils where they have been shown to migrate vertically to depths up to three feet (Palmer and Reeder, 2001) although most remain in the top 5 cm layer as long as the

soil is moist (Udonsi et al., 1980). Larvae have been shown to survive for up to six weeks although the yield becomes progressively lower over time (Beaver, 1953; Svensson, 1925; Udonsi and Atata, 1987). Other studies have shown that larvae survive for longer periods at lower temperatures (between 20 – 25°C) than at temperatures above 30°C (Svensson, 1925; Udonsi, 1988; Udonsi and Atata, 1987). When potential infectivity was assessed, it was found that larvae lost the ability to penetrate fresh rabbit skin in 11-15 days (Udonsi, 1988), apparently due to exhaustion of the worm energy stores. This evidence suggests that infectivity of hookworms in the environment is relatively short, on the order of a few weeks on average, and that the life cycle survival strategy of *N. americanus* is highly dependent on the 3-4 year longevity (Brooker et al., 2006b) of adult worms in infected human hosts.

Remote sensing and Geographic Information Systems (GIS) have been used to map the distribution of parasites and diseases and to develop prediction models of disease distribution in many regions of the world. An understanding of the environmental properties that contribute to the ecological niche of a particular parasite or vector is needed to be able to predict its distribution (Malone, 2005). Risk maps based on known biological requirements and climatic variables have been employed for a number of diseases including malaria (Yang et al., 2010), schistosomiasis (Lindsay and Thomas, 1999; Malone et al., 2001), filariasis (Lindsay and Thomas, 1999), leishmaniasis (Nieto et al., 2006; Salahi-Moghaddam et al., 2010) and others (Bergquist and Rinaldi, 2010). Such models have proven to be useful for formulating disease control programs in various parts of the world.

The concept of Growing Degree Days and Water Budget (GDD-WB) provides a measure of thermal-hydrological regime in relation to species specific knowledge of the base temperature (below which no development occurs), optimal temperature range for development, lethal upper limiting temperatures and limiting moisture regime in the context of the thermal units (GDD-WB) that must accumulate for a parasite-vector system to complete one or more generations. The GDD-WB concept has been used to generate predictive models of fascioliasis in the United States, Ethiopia, Ecuador and southern Brazil (Dutra et al., 2010; Fuentes et al., 2005; Malone et al., 1987; Malone et al., 2001), schistosomiasis in China and East Africa (Malone, 2005; Zhou et al., 2001) and visceral leishmaniasis in Bahia, Brazil (Nieto et al., 2006).

For hookworm, GIS and remote sensing methodologies have been used to study spatial distribution of the parasite and produce risk prediction models for the African continent (Brooker et al., 2006a), Kwazulu Natal in South Africa (Mabaso et al., 2003), and Ivory Coast (Raso et al., 2006). Risk maps based on biological data and climate regime have not yet been developed for hookworm infection in Brazil on a national scale.

In 2001 the World Health Assembly (WHA, 2001) World Health Organization's (WHO) member states agreed to implement or intensify control of schistosomiasis and STH infections. In 2009, the Directing Council of the Pan American Health Organization (PAHO) approved Resolution CD49.R19 urging member states to eliminate or drastically reduce, by 2015, the burden of 12 neglected infectious diseases including STH infections (PAHO, 2009). A study published in 2011 pointed out that important gaps of information on STH prevalence of infection at the first subnational level and below exist through-out the region including data of prevalence and intensity of infection by hookworm

(Schneider et al., 2011). Modeling the distribution and prevalence of STH infections could contribute to fill the existing gaps of epidemiological information.

The study reported here used hookworm prevalence data concurrently reported as part of surveillance and control programs for *S. mansoni* endemic areas in Brazil from 2000-2009. National scale risk maps on the distribution of hookworm infection were developed by applying known biological information on *N. americanus* to the climate envelope of conditions suitable for the survival, propagation and transmission of this hookworm species.

2.2 Materials and Methods

Brazil is located in South America at 10° South and 55° West (CIA, 2011). It has a mainly tropical climate with five major ecological zones (FAO, 2001). The hookworm prevalence data used was acquired from the Brazilian national notifiable diseases information system (Sistema Nacional de Agravos de Notificação – SINAN) and contained the proportion positive for hookworm at the municipality level from 2000 to 2009. The methods of surveillance were not defined in terms of numbers and ages of people examined, although examination of about 30% of a population is recommended by WHO and the Schistosomiasis Control Program in Brazil (Machado, 1982), by regular surveillance of school children. Based on the low numbers of people examined in some municipalities compared to the population, the municipality data points with less than 100 people examined were excluded to avoid bias, resulting in a total of 975 municipality data points.

The mapping and modeling software used was ArcGIS 9.3 (Environmental Systems Research Institute Inc., Redlands, CA, USA). Boundary map shape files for political units in Brazil were obtained from a South American Minimum Medical GIS database resource (Malone et al., 2007b) using data downloaded from the DIVA-GIS website (<http://diva.riu.cip.cgiar.org/index.php>). The climate data was a long-term-normal (LTN) climate grid (18x18 km cell size) of South America (Corbett, 2005). Each cell contained monthly long-term normal average data on rainfall, maximum daily temperature, minimum daily temperature, potential evapotranspiration, and the ratio of precipitation to potential evapotranspiration (PPE), also known as water budget (WB).

The Brazil administrative boundary map, prevalence data points, and climate grid data were imported into ArcGIS and re-projected to a geographic coordinate system (GCS), WGS 84 coordinate system format. The prevalence data point map and the climate grid attribute tables were joined using the polygon-to-point join function to produce a composite layer for GDD-WB analysis and calculation of the potential number of transmission cycles possible per year at hookworm positive municipalities. The values derived were used to extrapolate the model by calculating model variables for the entire national map, including municipalities where data was absent, using the layer properties/symbology function of ArcGIS.

2.2.1 GDD

Growing degree days (GDD) for hookworm were calculated by the field calculator function of ArcMap using a base temperature of 15°C for each month as follows (Malone et al., 2007a):

$$(Mean\ monthly\ temperature - 15) * Number\ of\ days\ in\ a\ particular\ month$$

Months where mean temperature was lower than 15°C were considered as having zero GDD.

2.2.2 Moisture Adjusted GDD Model (MA-GDD)

The accumulated annual GDD was calculated after excluding the GDD accumulated in dry months, identified as a month when the water budget was less than 0.4. (Soil saturation or ‘field capacity’ is reached at a value of 1.0 and monthly values above 1.0 have surplus/runoff water.) This cut-off was used because below 0.4 the prevalence of hookworm in the 2000-2009 hookworm survey database from Brazil was found to be absent or low (<5%).

2.2.3 GDD-Water Budget Gradient Model (GDD X WB)

The product of accumulated GDD and WB for each data point was used to develop a ‘gradient index’ that relates temperature to a moisture gradient (water budget). The gradient index model was calculated for each month and the annual value was calculated by summing the monthly GDDxWB values.

2.2.4 Calculation of Mean Transmission Time and Potential Transmission Cycles per Year

The mean transmission time was defined as the time spent in the free-living state of the hookworm, from deposition of eggs into the environment to infection of the host by viable infective third stage larvae. It was calculated based on literature reports (Beaver, 1953; Svensson, 1925; Udonsi, 1988) by adding development time from egg to infective larva (14 days) and time as active infective larva (14 days).

$$\text{Transmission cycle GDD} = (\text{Mean transmission time}) * \text{Daily GDD}$$

Potential Transmission cycles per year values were calculated based on the number of potential transmission cycles per annum by dividing annual MA-GDD by Transmission cycle GDD (224). Models were then developed using WB, GDD and transmission cycle values to predict potential risk zones for hookworm transmission in Brazil.

2.2.5 Model Validation

A literature search was conducted to review reports on independent surveys for prevalence of hookworm and other STHs in Brazil. Results were used to validate the MA-GDD and GDDxWB models.

2.2.6 Statistical Analysis

Statistical Analysis Software (SAS®, Cary, NC, USA), was used to determine correlations between prevalence of hookworm and the variables MA-GDD and GDDxWB using Pearson Correlation. The data were then log₁₀ transformed for normalizing. Analysis of variance (ANOVA) followed by Tukey's test was used to compare MA-GDD means among low (1-4.9%), moderate (5-19.9%) and high (<20%) categories of hookworm prevalence. These categories were adapted from WHO guidelines for the control of STHs (WHO, 2006). The same was done for the GDDxWB categories. Data points with prevalence less than 1% were assumed negligible due to possible migration and were excluded from ANOVA, resulting in 564 data points of the 975 municipalities with data. Polynomial regression was used to determine the relationship between mean transmission cycles and the GDDxWB index.

2.3 Results

The mean prevalence of hookworm among the municipalities in the data set was 4.99% (range 0-70.22; Standard Deviation 8.93). The prevalence distribution at 975 municipality sites is shown in Table 2.1, categorized as very low, low, moderate and high prevalence.

Table 2.1. Distribution of hookworm prevalence among the 975 municipalities surveyed with corresponding MA-GDD and GDDxWB values.

Categories	Hookworm Prevalence (%)	Number of municipalities	MA-GDD Mean (range; SEM)	GDDxWB Mean (range; SEM)
Very Low	<1	411	1672.2 (0-3586; 30.0)	2029.9 (0-6618; 41.5)
Low	1-4.9	303	1752.7 (540-3280; 29.2)	2274.4 (666-7117; 60.5)
Moderate	5-19.9	187	1891.0 (1088-3099; 28.0)	2563.0 (1016-7041; 71.0)
High	>20	74	2057.7 (913-3014; 58.5)	3493.8 (1134-7875; 207.0)
	Total	975		

The Pearson correlation coefficients between hookworm prevalence and MA-GDD and between hookworm prevalence and GDDxWB index were 0.24 ($p<0.001$) and 0.27 ($p<0.001$) respectively. The MA-GDD and GDDxWB \log_{10} means in the three categories of prevalence (low, moderate, high) were all significantly different ($p<0.001$) (i.e. the MA-GDD \log_{10} mean among the, low prevalence group was significantly different from that in the moderate and high prevalence groups and that between the moderate prevalence group was different from the high prevalence group). However, the MA-GDD

range in the high prevalence group overlapped with the moderate prevalence group range (Table 2.1).

The relationship between the mean transmission cycles and GDDxWB index was approximated by the formula $y=2.3+0.0031x-2E-07x^2$ ($R^2=0.522$) where y is mean number of transmission cycles and x is GDDxWB.

The MA-GDD model (Figure 2.1) and the GDDxWB model (Figure 2.2) both predicted a low hookworm prevalence of less than 5% in the northeastern inland (Caatinga) region and in the southern region of Brazil. The MA-GDD model (Figure 2.3) showed that where the transmission cycles were less than 5 per year, or more than 14 per year, the hookworm prevalence was low (<5%). The GDDxWB index model showed a low end cut off at 1300 (Figure 4) below which the prevalence of hookworm fell below 5%.

Thirty independent reports published between 1990 and 2010 by various authors were reviewed and used for the purpose of validating the climate risk prediction models developed in the current study (Aguiar et al., 2007; Alves et al., 2003; Bóia et al., 2009; Braga et al., 1998; Brooker et al., 2007; Buschini et al., 2007; Cabral et al., 2000; Carvalho-Costa et al., 2007; Cury et al., 1994; Ferrari et al., 1992; Ferreira et al., 1994; Geiger et al., 2004; Gomes et al., 2002; Gonçalves et al., 1990; Heukelbach et al., 2004; Lopes et al., 2006; Machado et al., 2008a; Machado et al., 2008b; Melo et al., 2010; Miranda et al., 1998; Miranda et al., 1999; Moraes and Cairncross, 2004; Oliveira et al., 2003; Rocha et al., 2000; Santos et al., 1995; Scolari et al., 2000; Silva and da Silva, 2010; Silva et al., 2007; Souza et al., 2007; Tsuyuoka et al., 1999). These articles covered 16 states out of Brazil's 26 states and hookworm prevalence ranged from 0 to 80.2% with

a mean of 20.3%. The number of people examined ranged from 54 to 13,279 and sampled populations ranged from school children to whole communities. There were eight low, 11 moderate and 11 high hookworm prevalence sites. While the 30 data points available from literature were limited there was a general agreement of survey data to the predicted GDDxWB index risk map showing high hookworm prevalence risk in the amazon basin (Figure 2.2). The Pearson correlation coefficient between the prevalence and GDDxWB was 0.26 ($p = 0.158$) and between prevalence and MA-GDD was 0.15 ($p = 0.4116$).




Figure 2.1. Risk Model for Hookworm using Moisture Adjusted GDD. Showing moderate to high prevalence is most likely to occur between five and 14 transmission cycles per year (yellow to dark brown). Areas with less than five transmission cycles per year (yellow) are prohibitive to hookworm transmission. Blue dots represent municipality data points.




Figure 2.2. Hookworm prevalence risk model using GDDxWB index. Areas of low (yellow) to high (dark brown) prevalence risk are shown. Blue dots represent municipality data points. Red graduated circles denote hookworm prevalence from independent survey reports.




Figure 2.3. Scatter plot of hookworm prevalence from municipality data points against transmission cycles per year. The plot shows a lower end cut off at five transmission cycles below which the prevalence is less than 5% and an upper end cut off at about 14 transmission cycles above which the prevalence is less than 5%. Black line is a trend line ($y=0.9702x$).




Figure 2.4. Scatter plot of Hookworm Prevalence against GDD-WB index. Prevalence levels increase dramatically above 1300 with no obvious upper limit. Black line is a trend line ($y=0.0031x$).

2.4 Discussion

In the current study on development of ecological niche models for hookworms in Brazil, there was a need for a standard measure of the developmental suitability of a site for propagation and transmission of the parasite based on climate and known biological parameters that can distinguish low, moderate and high risk conditions for hookworm disease.

Temperature and moisture environmental parameters are fundamental determinants of the distribution range and abundance of free-living species (Andrewartha and Birch, 1954) and suitable ranges are vital for hookworm eggs to hatch and for the larvae to survive and be transmitted to susceptible hosts (Udonsi and Atata, 1987). We developed and compare here two climate-based models to evaluate geographic risk of hookworm disease, the MA-GDD and GDDxWB indices, using a LTN climate grid (18 km^2) of Brazil. Both indices take into account environmental temperatures conducive to the development of

the free living stages of the parasite as well as prevailing moisture conditions. The MA-GDD index considers moisture above a given threshold of 0.4 and the GDDxWB index considers the entire range of a gradient of moisture.

2.4.1 MA-GDD Index and Transmission Cycles

The MA-GDD index reflects suitable days for development above the base temperature of 15°C, from which can be calculated the number of ‘potential transmission cycles’ that can be completed per year. Hookworm infected individuals have been reported to shed eggs continuously for long periods of time, leading to development of infective larvae (L₃) and potential year-round transmission. The developmental time of the hookworm from L₃ to egg producing adults in the host is fairly constant, with a mean of eight weeks (Beaver, 1988; Hoagland and Schad, 1978). The variation in generation time (from egg to egg-producing adult) arises from the free-living stages of the parasite, here referred to as the mean transmission cycle period, which is highly susceptible to variation in environmental conditions. Because of the high dependence on moisture of hookworms for movement, and to prevent desiccation, annual GDD was adjusted (MA-GDD) to exclude months where mean water budget was less than 0.4 as explained above. This result suggests that larvae development and L₃ transmission is impaired as soils get drier (Nwosu and Anya, 1980; Udonsi and Atata, 1987). Dry months were thus not included in the index calculations based on the assumption that the GDD from those months do not contribute to the annual accumulated GDD.

We propose use of the transmission cycle parameter as a biology-based criterion for regional risk modeling based on the GDD-WB concept. Literature reports indicate that the development time from egg to L₃ can be up to 14 days at temperatures less than 40°C

(Brooker et al., 2006b; Schad and Warren, 1990), and the period of time that the majority of infective larva remain active in the environment ranges from one to four weeks (Beaver, 1953; Svensson, 1925; Udonsi, 1988). Therefore, taking two weeks for development to the L₃ stage and two weeks longevity (on average) in which L₃ must find a definitive host, the transmission time was estimated to be four weeks (28 days). This criterion was used as an indicator of biological suitability of different climate thermal-hydrological regimes for the purposes of this study. However, depending on the basis for calculation, which references are used, as well as availability of human hosts the transmission time can potentially range from 9 days (Palmer and Reeder, 2001) if infection occurs immediately after molting to L₃, to approximately 51 days (Palmer and Reeder, 2001; Svensson, 1925; Udonsi, 1988) if the larvae survive in the environment. Our estimation of the transmission cycle takes into account factors that may delay the embryonation and hatching of eggs (eg. low temperature) and also the period in which the L₃ are reported to be active in the environment prior to infection (Table 2.2).

The average annual mean temperature prevalent in Brazil is 23°C, a thermal regime expected to promote longevity of the larvae in the environment. *N. americanus*, unlike *A. duodenale*, does not exhibit hypobiosis (Loukas and Prociv, 2001) which could interrupt continuous egg shedding. Thus it is possible to use mean transmission cycles per year for describing *N. americanus*'s distribution and to generate risk map surfaces assuming adult populations are long lived (i.e. 4 years).

In the MA-GDD model, most areas with prevalence above 5% lay between index values of 1500 to 3000, allowing 5-14 mean transmission cycles per year (Figure 2.3). Areas with index values below this range lay in the Caatinga ecological zone, which is

Table 2.2. Important biological and climatic variables for the transmission of *N. americanus* used as baseline values for calculations based on the GDD-WB concept.

Variables	Value	References
Base Temperature	15 °C	(Udonsi and Atata, 1987)
Mean Annual Temperature	23 °C	From Brazil climate data
Daily GDD	8	
Mean transmission cycle time (free-living phase)	28 days (range 9-51)	(Beaver, 1953; Croll and Matthews, 1973; Palmer and Reeder, 2001; Udonsi, 1988; Udonsi and Atata, 1987)
Transmission cycle GDDs	224	
Mean Egg life expectancy	13-14 days	(Schad and Warren, 1990)
Adult life span	3-4 years	(Schad and Warren, 1990)

characterized by deserts and xeric shrub and in the southern region of the country where mean temperatures are less than 21°C. Regions with the highest MA-GDD index lay in the Amazon region, the northern coast area and the Bahia coastal forests, which are characterized by moist broadleaf forests. Early work (Chandler, 1929) more recent literature reports suggest the Amazon region has high prevalence in indigenous populations (Bóia et al., 2009; Botero, 1981; Miranda et al., 1998). In the Bahia coastal forests of high annual rainfall the low prevalence suggests that hookworms do not survive long where there is frequent flooding and run-off and that larval stages do not survive well in water without suitable substrate (Schad and Warren, 1990).

2.4.2 GDDxWB Index

The GDDxWB index model had a higher correlation coefficient for hookworm prevalence as compared to MA-GDD (0.27 versus 0.24). This may be because MA-GDD did not have a completely linear relationship to hookworm prevalence, having an upper

end cut off (Figure 3). Risk maps based on both the MA-GDD and GDDxWB indices results suggest low temperature (mean temperatures less than 21°C) inhibit hookworm transmission in southern Brazil where climate is much cooler, with minimum temperatures commonly falling below 15°C, the base temperature for egg hatching of hookworm (Schad and Warren, 1990).

In contrast to the MA-GDD, the GDDxWB index provides the advantage of measuring a moisture *gradient* of suitability but not the number of transmission cycles per year. However, it is possible to relate transmission cycles and the GDDxWB index by use of a quadratic equation (described above) to calculate the number of transmission cycles expected at a given GDDxWB index value. For both the MA-GDD and the GDDxWB index there was a significant difference in log₁₀ means of GDDxWB between areas of low, moderate and high prevalence of hookworm (Figure 4). At above a certain climatic threshold, however, it is typical to observe considerable variation in actual versus potential transmission levels at individual sites even in high risk climates because of location-specific factors related to poverty, occupational-agricultural practices or local environment, such as location along riverine wetland plains. Climate risk is therefore an indicator of *potential* transmission related to mean transmission of individual sites in a broader regional climatic envelope (Malone, 2005).

2.4.3 Data Validation

A relatively limited number of 30 survey reports were available in the literature to validate the climate risk models developed in the current study based on data collected as part of the national schistosomiasis control program which focused on areas with known or suspected schistosomiasis endemicity. Results from the correlation coefficients

suggest the majority of the validation points fit better in the GDDxWB model (Figure 2) than the MA-GDD model. The high hookworm prevalence predicted for the Amazon basin by both the GDD-WB and GDDxWB models is consistent with results reported by Chandler (1929) and earlier reports that hookworm occurs in most of the country, including the Amazon, and has lower prevalence in the drier areas found in the northeastern part of the country (DNERu, 1962). In order to further validate the climate-based models produced in the current study, a comprehensive national survey for STH is needed, since hookworm often co-infects with other STH (Fleming et al., 2006). A comprehensive survey should include areas of limited surveillance for schistosomiasis such as in the Amazon basin and the Caatinga region.

2.4.4 Other Environmental Factors

The objective of the current study was to develop climate based models that enable mapping of the climate envelop of hookworms in Brazil and did not take into account other factors relating to the prevalence of hookworm that can be used to further develop and refine climate risk maps in future studies. Other environmental variables identified in literature reports as important in hookworm propagation and transmission include soil type, slope, altitude and land surface temperature measurements derived from environmental satellites (Brooker et al., 2006b; Mabaso et al., 2003; Udonsi et al., 1980). Social-economic status has long been recognized as an important variable relating to parasitic infections including hookworm (Hotez, 2008) and a similar trend has been shown in Brazil (Fleming et al., 2006). Vegetation cover or shade have been shown to be important for the longevity of larvae (Udonsi and Atata, 1987), suggesting an impact of deforestation. Further studies are underway to refine maps based on climate suitability

alone and to validate the influence of these additional factors on the ecological niche of *N. americanus*.

In summary, the prevalence of hookworm in Brazil appears to be strongly related to both warmer and moister climatic conditions where the GDDxWB is above 1300 and where the MA-GDD dictates a minimum number of five transmission cycles per year. Low prevalence is predicted in areas that have mean monthly temperatures below 21°C. The current study supports the validity of the GDD-WB concept for use in mapping risk areas for hookworm. There is need for a systematic national survey for hookworm and other STH in Brazil to elucidate STH disease risk outside the schistosomiasis endemic areas where very limited surveillance exists. The models developed can be used to identify high hookworm prevalence risk areas for purposes of planning further surveillance and control efforts in Brazil.

2.5 References

- Aguiar, J.I.A., Gonçalves, A.Q., Sodré, F.C., Pereira, S.d.R., Bóia, M.N., Lemos, E.R.S.d., Daher, R.R., 2007. Intestinal protozoa and helminths among Terena Indians in the state of Mato Grosso do Sul: high prevalence of *Blastocystis hominis*. *Rev Soc Bras Med Trop* 40, 631-634.
- Alves, J.R., Macedo, H.W., Ramos Jr., A.N., Ferreira, L.F., Gonçalves, M.L.C., Araújo, A., 2003. Parasitoses intestinais em região semi-árida do Nordeste do Brasil: resultados preliminares distintos das prevalências esperadas. *Cad Saude Publica* 19, 667-670.
- Amaral, R.S.d., Tauil, P.L., Lima, D.D., Engels, D., 2006. An analysis of the impact of the Schistosomiasis Control Programme in Brazil. *Mem Inst Oswaldo Cruz* 101, 79-85.
- Andrewartha, H.G., Birch, L.C., 1954. The distribution and abundance of animals. University of Chicago Press, Chicago.
- Beaver, P.C., 1953. Persistence of hookworm larvae in soil. *Am J Trop Med Hyg* 2, 102-108.

- Beaver, P.C., 1988. Light, long-lasting *Necator* infection in a volunteer. *Am J Trop Med Hyg* 39, 369-372.
- Bergquist, R., Rinaldi, L., 2010. Health research based on geospatial tools: a timely approach in a changing environment. *J Helminthol* 84, 1-11.
- Bóia, M.N., Carvalho-Costa, F.A., Sodré, F.C., Porras-Pedroza, B.E., Faria, E.C., Magalhães, G.A.P., Silva, I.M.d., 2009. Tuberculose e parasitismo intestinal em população indígena na Amazônia brasileira. *Rev Saúde Pública* 43, 176-178.
- Booth, M., Vounatsou, P., N'goran, E.K., Tanner, A., Utzinger, J., 2003. The influence of sampling effort and the performance of the Kato-Katz technique in diagnosing *Schistosoma mansoni* and hookworm co-infections in rural Côte d'Ivoire. *Parasitology* 127, 525-531.
- Botero, D., 1981. Persistence of the intestinal parasites endemic in Latin America. *Bull Pan Am Health Organ* 15, 241-248.
- Braga, L.L., Mendonca, Y., Paiva, C.A., Sales, A., Cavalcante, A.L.M., Mann, B.J., 1998. Seropositivity for and intestinal colonization with *Entamoeba histolytica* and *Entamoeba dispar* in individuals in northeastern Brazil. *J Clin Microbiol* 36, 3044-3045.
- Brooker, S., Clements, A., Hotez, P., Hay, S., Tatem, A., Bundy, D., Snow, R., 2006a. The co-distribution of *Plasmodium falciparum* and hookworm among African schoolchildren. *Malar J* 5, 99.
- Brooker, S., Clements, A.C.A., Bundy, D.A.P. 2006b. Global Epidemiology, Ecology and Control of Soil-Transmitted Helminth Infections, In: Simon I. Hay, A.G., David, J.R. (Eds.) *Advances in Parasitology*. Academic Press, 221-261.
- Brooker, S., Jardim-Botelho, A., Quinnell, R.J., Geiger, S.M., Caldas, I.R., Fleming, F., Hotez, P.J., Correa-Oliveira, R., Rodrigues, L.C., Bethony, J.M., 2007. Age-related changes in hookworm infection, anaemia and iron deficiency in an area of high *Necator americanus* hookworm transmission in south-eastern Brazil. *Trans R Soc Trop Med Hyg* 101, 146-154.
- Buschini, M.L.T., Pittner, E., Czervinski, T., Moraes, I.d.F., Moreira, M.M., Sanches, H.F., Monteiro, M.C., 2007. Spatial distribution of enteroparasites among school children from Guarapuava, state of Paraná, Brazil. *Rev Bras Epidemiol* 10, 568-578.
- Cabral, D.D., Mundim, M.J.S., Oliveira, P.R.d., Santos, S.d.F.O., Gomes, S.M., 2000. Intestinal parasitoses in school children from Gouvelândia, Goiás State. *Bioscience J* 16, 75-85.
- Carvalho-Costa, F.A., Gonçalves, A.Q., Lassance, S.L., Silva Neto, L.M.d., Salmazo, C.A.A., Bóia, M.N., 2007. *Giardia lamblia* and other intestinal parasitic

- infections and their relationships with nutritional status in children in Brazilian Amazon. *Rev I Med Trop Sao Paulo* 49, 147-153.
- Chandler, A.C., 1929. Hookworm Disease. Macmillan Company, New York.
- CIA 2011. The World Factbook. (<https://www.cia.gov/library>).
- Corbett, J.D. 2005. South America Minimum Medical GIS Database, Malone, J.B., Wiles, S.E., Nieto, P., McCarroll, J.C., Bavia, M., eds.
- Croll, N.A., Matthews, B.E., 1973. Activity, ageing and penetration of hookworm larvae. *Parasitology* 66, 279-289.
- Cury, G.C., Salles, P.G., Reis, M.C., Rego, V.M., Arndt, A.W., de Souza Filho, C.B., Moreira, H.B., de Abreu, M.S., Leite, P.M., 1994. Prevalence of *Schistosoma mansoni* and parasitic intestinal diseases among students of the rural area of the municipality of Jaboticatubas, MG, 1992-1993. *Rev Soc Bras Med Trop* 27, 217-220.
- DNERu, 1962. Departamento Nacional de Endemias Rurais. Combate a endemias rurais no Brasil. Ministerio da Saude.
- Dutra, L.H., Molento, M.B., Naumann, C.R.C., Biondo, A.W., Fortes, F.S., Savio, D., Malone, J.B., 2010. Mapping risk of bovine fasciolosis in the south of Brazil using geographic information systems. *Vet Parasitol* 169, 76-81.
- FAO 2001. Global Ecological Zoning for the Global Forest Resources Assessment 2000 (Rome), 91-105.
- Ferrari, J.O., Ferreira, M.U., Aranha Camargo, L.M., Ferreira, C.S., 1992. Intestinal parasites among Karitiana indians from Rondônia State, Brazil. *Rev I Med Trop Sao Paulo* 34, 223-225.
- Ferreira, C.S., Ferreira, M.U., Nogueira, M.R., 1994. The prevalence of infection by intestinal parasites in an urban slum in Sao Paulo, Brazil. *J Trop Med Hyg* 97, 121-127.
- Fleming, F.M., Brooker, S., Geiger, S.M., Caldas, I.R., Correa-Oliveira, R., Hotez, P.J., Bethony, J.M., 2006. Synergistic associations between hookworm and other helminth species in a rural community in Brazil. *Trop Med Int Health* 11, 56-64.
- Fuentes, M.V., Sainz-Elipse, S., Nieto, P., Malone, J.B., Mas-Coma, S., 2005. Geographical information systems risk assessment models for zoonotic fascioliasis in the South American Andes region. *Parassitologia* 47, 151-156.
- Geiger, S.M., Massara, C.L., Bethony, J., Soboslay, P.T., CorrêA-Oliveira, R., 2004. Cellular responses and cytokine production in post-treatment hookworm patients from an endemic area in Brazil. *Clin Exp Immunol* 136, 334-340.

- Gomes, T.C., Almeida, M.F., Miura, L.A., Granja, J., Santos, D.V.G., Oliveira, R.M.F., Lopes, A., Sequeira, B.P., Rolemberg, A.A., Moraes, A.L., Santos, C.S., 2002. Helminthoses intestinais em população de rua da cidade do Rio de Janeiro. *Rev Soc Bras Med Trop* 35, 531-532.
- Gonçalves, J.F., Tanabe, M., Medeiros, F.d.P.d.M., Gonçalves, F.J., Aca, I.d.S., Motta, S.R.N.d., Tateno, S., Takeuchi, T., 1990. Parasitological and serological studies on amoebiasis and other intestinal parasitic infections in the rural sector around Recife, Northeast Brazil. *Rev Inst Med Trop São Paulo* 32, 428-435.
- Heukelbach, J., Winter, B., Wilcke, T., Muehlen, M., Albrecht, S., Oliveira, F.A.S.d., Kerr-Pontes, L.R.S., Liesenfeld, O., Feldmeier, H., 2004. Selective mass treatment with ivermectin to control intestinal helminthiasis and parasitic skin diseases in a severely affected population. *Bull World Health Org* 82, 563-571.
- Hoagland, K.E., Schad, G.A., 1978. *Necator americanus* and *Ancylostoma duodenale*: Life history parameters and epidemiological implications of two sympatric hookworms of humans. *Exp Parasitol* 44, 36-49.
- Hotez, P., 2008. Hookworm and Poverty. *Ann NY Acad Sci* 1136, 38-44.
- Lindsay, S.W., Thomas, C.J., 1999. Mapping and estimating the population at risk from lymphatic filariasis in Africa. *Trans R Soc Trop Med Hyg* 94, 37-45.
- Lopes, F.M.R., Gonçalves, D.D., dos Reis, C.R., Breganó, R.M., Filho, F.A., Murad, V.A., Dutra de Menezes, M.C.N., Freire, R.L., de Freitas, J.C., Santana, M.A.Z., 2006. Occurrence of enteroparasitosis in schoolchildren of the municipal district of Jataizinho, state of Paraná, Brazil. *Acta Sci-Health Sci* 28, 107-111.
- Loukas, A., Prociv, P., 2001. Immune responses in hookworm infections. *Clin. Microbiol. Rev.* 14, 689-703.
- Mabaso, M.L.H., Appleton, C.C., Hughes, J.C., Gouws, E., 2003. The effect of soil type and climate on hookworm (*Necator americanus*) distribution in KwaZulu-Natal, South Africa. *Trop Med Int Health* 8, 722-727.
- Machado, E.R., de Souza, T.S., da Costa, J.M., Costa-Cruz, J.M., 2008a. Enteroparasites and commensals among individuals living in rural and urban areas in Abadia dos Dourados, Minas Gerais state, Brazil. *Parasitologia Latinoamericana* 63, 34-39.
- Machado, E.R., Santos, D.S., Costa-Cruz, J.M., 2008b. Enteroparasites and commensals among children in four peripheral districts of Uberlândia, state of Minas Gerais. *Rev Soc Bras Med Trop* 41, 581-585.
- Machado, P.A., 1982. The Brazilian program for schistosomiasis control, 1975-1979. *Am J Trop Med Hyg* 31, 76-86.

- Malone, J.B., 2005. Biology-based mapping of vector-borne diseases by geographic information systems and remote sensing. . *Parasitologia* 47, 27-50.
- Malone, J.B., McCarroll, J.C., Wiles, S.E. 2007a. Geospatial Health and Environment: A lessons manual and CDROM Data Resource, University, L.S., ed. (Baton Rouge).
- Malone, J.B., Seklau, E.W., McCarroll, J.C., Bavia, M. 2007b. South American Region Minimum Medical GIS Database.
- Malone, J.B., Williams, T.E., Muller, R. A., Geaghan, J. P., Loyacano, A. F., 1987. Fascioliasis in cattle in Louisiana: development of a system to predict disease risk by climate, using the Thornthwaite water budget. *American Journal of Veterinary Research* 48, 1167-1170.
- Malone, J.B., Yilma, J.M., McCarroll, J.C., Erko, B., Mukaratirwa, S., Zhou, X., 2001. Satellite climatology and the environmental risk of *Schistosoma mansoni* in Ethiopia and east Africa. *Acta Tropica* 79, 59-72.
- Melo, G.C., Reyes-Lecca, R.C., Vitor-Silva, S., Monteiro, W.M., Martins, M., Benzecry, S.G., Alecrim, M.d.G.C., Lacerda, M.V.G., 2010. Concurrent helminthic infection protects schoolchildren with *Plasmodium vivax* from anemia. *PLoS ONE* 5, e11206.
- Miranda, R.A., Xavier, F.B., Menezes, R.C., 1998. Intestinal parasitism in a Parakanã indigenous community in southwestern Pará State, Brazil. *Cad Saude Publica* 14, 507-511.
- Miranda, R.A., Xavier, F.B., Nascimento, J.R., de Menezes, R.C., 1999. The prevalence of intestinal parasitism in native villages of the Tembê tribe, the Brazilian eastern Amazon. *Rev Soc Bras Med Trop* 32, 389-393.
- Moraes, L.R.S., Cairncross, S., 2004. Environmental interventions and the pattern of geohelminth infections in Salvador, Brazil. *Parasitology* 129, 223-232.
- Nieto, P., Malone, J.B., Bavia, M.E., 2006. Ecological niche modeling for visceral leishmaniasis in the state of Bahia, Brazil, using genetic algorithm for rule-set prediction and growing degree day-water budget analysis. . *Geospatial Health* 1, 115-126.
- Nwosu, A.B.C., Anya, A.O., 1980. Seasonality in human hookworm infection in an endemic area of Nigeria, and its relationship to rainfall. *Tropenmedizin und Parasitologie* 31, 201-208.
- Oliveira, M.C., Silva, C.V.d., Costa-Cruz, J.M., 2003. Intestinal parasites and commensals among individuals from a landless camping in the rural area of Uberlândia, Minas Gerais, Brazil. *Rev Inst Med Trop São Paulo* 45, 173-176.

- PAHO, 2009. Elimination of Neglected Diseases and Other Poverty-Related Infections. 49th Directing Council Resolution CD49.R19.
- Palmer, P.E.S., Reeder, M.M. 2001. Imaging of tropical diseases: with epidemiological, pathological and clinical correlation.(Berlin, Springer-Verlag, 52-65.
- Raso, G., Vounatsou, P., Gosoni, L., Tanner, M., N'Goran, E.K., Utzinger, J., 2006. Risk factors and spatial patterns of hookworm infection among schoolchildren in a rural area of western Côte d'Ivoire. *Int J Parasitol* 36, 201-210.
- Rocha, R.S., Silva, J.G., Peixoto, S.V., Caldeira, R.L., Firmo, J.O., Carvalho O dos, S., Katz, N., 2000. Assessment of schistosomiasis and other intestinal parasitoses in school children of the Bambuí municipality, Minas Gerais, Brazil. *Rev Soc Bras Med Trop* 33, 431-436.
- Salahi-Moghaddam, A., Mohebbi, M., Moshfai, A., Majid, H., Zarei, Z., 2010. Ecological study and risk mapping of visceral leishmaniasis in an endemic area of Iran based on a geographical information systems approach. *Geospatial Health* 5, 71-77.
- Santos, R.V., Coimbra Júnior, C.E., Flowers, N.M., Silva, J.P., 1995. Intestinal parasitism in the Xavante Indians, central Brazil. *Rev Inst Med Trop Sao Paulo* 37, 145-148.
- Schad, G.A., Warren, K.S. 1990. Hookworm Disease, Current Status and New Directions Taylor & Francis Ltd.
- Schneider, M.C., Aguilera, X.P., Barbosa da Silva Junior, J., Ault, S.K., Najera, P., Martinez, J., Requejo, R., Nicholls, R.S., Yadon, Z., Silva, J.C., Leanes, L.F., Periago, M.R., 2011. elimination of neglected diseases in Latin America and the Caribbean: a mapping of selected diseases. *PLoS Negl Trop Dis* 5, e964.
- Scolari, C., Torti, C., Beltrame, A., Matteelli, A., Castelli, F., Gulletta, M., Ribas, M., Morana, S., Urbani, C., 2000. Prevalence and distribution of soil-transmitted helminth (STH) infections in urban and indigenous schoolchildren in Ortigueira, State of Paraná, Brasil: implications for control. *Trop Med Int Health* 5, 302-307.
- Silva, L.P., da Silva, R.M.G., 2010. Occurrence of enteroparasites in centers education child of the municipality of Patos of Minas, MG, Brazil. *Bioscience J* 26, 147-151.
- Silva, R.d.C.R., Barreto, M.L., Assis, A.M.O., de Santana, M.L.P., Parraga, I.M., Reis, M.G., Blanton, R.E., 2007. The relative influence of polyparasitism, environment, and host factors on schistosome Infection. *Am J Trop Med Hyg* 77, 672-675.
- Souza, E.A.d., Silva-Nunes, M.d., Malafronte, R.d.S., Muniz, P.T., Cardoso, M.A., Ferreira, M.U., 2007. Prevalence and spatial distribution of intestinal parasitic infections in a rural Amazonian settlement, Acre State, Brazil. *Cad Saúde Pública* 23, 427-434.

- Stoltzfus, R.J., Albonico, M., Chwaya, H.M., Savioli, L., Tielsch, J., Schulze, K., Yip, R., 1996. Hemoquant determination of hookworm-related blood loss and its role in iron deficiency in african children. *Am J Trop Med Hyg* 55, 399-404.
- Svensson, R., 1925. Observations on the development and longevity of hookworm larvae in defferent temperature conditions. *Chin Med J* 39, 667-673.
- Tsuyuoka, R., Bailey, J.W., Guimarães, A.M.d.A.N., Gurgel, R.Q., Cuevas, L.E., 1999. Anemia and intestinal parasitic infections in primary school students in Aracaju, Sergipe, Brazil. *Cad Saúde Pública* 15, 413-421.
- Udonsi, J.K., 1988. Experimental and field studies on the viability of eggs, and the infectivity of third-stage larvae of the human hookworm *Necator americanus* recovered from the creek waters of the Niger Delta, Nigeria, and their epidemiological significance. *Parasitology* 96, 111-117.
- Udonsi, J.K., Atata, G., 1987. *Necator americanus*: Temperature, pH, light, and larval development, longevity, and desiccation tolerance. *Exp Parasitol* 63, 136-142.
- Udonsi, J.K., Nwosu, A.B.C., Anya, A.O., 1980. *Necator americanus*: population structure, distribution, and fluctuations in population densities of infective larvae in contaminated farmlands. *Parasitol Res* 63, 251-259.
- WHA 2001. Resolution WHA54.19: Schistosomiasis and soil-transmitted helminth infections.
- WHO, 2006. Preventive chemotherapy in human helminthiasis: coordinated use of anthelmintic drugs in control interventions: a manual for health professionals and programme managers. World Health Organization, Geneva.
- Yang, G.-J., Gao, Q., Zhou, S.S., Malone, J.B., McCarroll, J.C., M., T., Vounatsou, P., Bergquist, R., Utzinger, J., Zhou, X.-N., 2010. Mapping and predicting malaria transmission in the People's Republic of China, using integrated biology-driven and statistical models. *Geospatial Health* 5, 11-22.
- Zhou, X.N., Malone, J.B., Kristensen, T.K., Bergquist, N.R., 2001. Application of geographic information systems and remote sensing to schistosomiasis control in China. *Acta Trop* 79, 97-106.

3 HOOKWORM AND OTHER SOIL TRANSMITTED HELMINTHS AMONG PATIENTS IN MUTUÍPE, BRAZIL

3.1 Introduction

Soil transmitted helminths (STH) which include hookworm, *Ascaris lumbricoides* and *Trichuris trichiura* are found worldwide. It is estimated that more than 1.5 billion people are infected with at least one species of STH globally (Bethony et al., 2006; WHO, 2013). As a group, they represent the most prevalent parasitic infection of man and are most common in tropical and subtropical countries in East Asia, sub-Saharan Africa, and the Americas where they are associated with high levels of poverty (Brooker et al., 2006b). Co-infection with all three parasite species is also common (Bethony et al., 2006) and children infected with STH have been found to have physical and intellectual retardation, reduced cognitive ability and malnutrition (Bethony et al., 2006; Eppig et al., 2010; Jardim-Botelho et al., 2008; Stephenson et al., 1993; Yap et al., 2012). Other symptoms of infection include diarrhea, anemia, and abdominal pain (Beaver, 1975; Ezeamama et al., 2005).

Mass drug administration, which refers to the periodic distribution and administration of drugs to at-risk populations in endemic areas (WHA, 2001), is the main approach used in the control of STH (Hotez et al., 2007). The primary drugs used in the treatment and control of STH are benzimidazoles, particularly albendazole and mebendazole (Hotez et al., 2008). Control strategies are based on prevalence and intensity of infection in a community, with intensity of infection being estimated from fecal egg counts (Montresor et al., 1998).

Hookworm infection is the most important infection among the intestinal nematodes and ranks fourth among parasitic infections worldwide after malaria, leishmaniasis and lymphatic filariasis (Hotez et al., 2005) as determined by disability-affected life years (DALY) . It is estimated that approximately 740 million people worldwide are infected with hookworm (de Silva et al., 2003). Hookworm disease is often characterized by iron deficiency anemia which is of great concern in women of reproductive age and children (Brooker et al., 1999; Dreyfuss et al., 2000; Hotez et al., 2005).

The two human hookworms that are most prevalent are *Necator americanus*, occurring mostly in the tropics and *Ancylostoma duodenale*, found in more temperate climates. The few studies done so far in Brazil (Minas Gerais and São Paulo states) that have identified the hookworm as to species reported found predominantly *N. americanus* (Fleming et al., 2006; Geiger et al., 2004; Kobayashi et al., 1995). However, a few *A. duodenale* infections were reported in Pernambuco state, in northeastern Brazil (Gonçalves et al., 1990; Okazaki et al., 1988). Because the two species have different epidemiological and ecological characteristics, it is important to differentiate them to be able to properly plan control measures (Brooker et al., 2006a), particularly with regard to public awareness and sanitation.

Most parasitological surveys for intestinal helminths utilize the Kato-Katz thick smear for quick diagnosis of parasite infection as recommended by WHO (Montresor et al., 1998). This method, though cost effective and technologically easy to perform lacks sensitivity and specificity for hookworm species detection (Basuni et al., 2011) when compared to time consuming classic morphological analysis or molecular methods like Polymerase Chain Reaction (PCR).

Levecke et al. (2011) compared the Kato-Katz method to the McMaster flotation method that is commonly used in veterinary parasitology and they found that Kato-Katz was more sensitive and resulted in higher fecal egg counts for *A. lumbricoides* than the McMaster method. However, there was no significant difference when used for *T. trichiura* and hookworm egg detection.

The use of PCR to differentiate hookworm species has proved useful because hookworm eggs are difficult to differentiate by microscopy as they are of similar size. Coproculture is useful for differentiating strongylid nematode infective larvae and may also be used to differentiate *N. americanus* from the *Ancylostoma* sp. However, it is time consuming and requires skilled personnel (de Gruijter et al., 2005). Using molecular methods, the internal transcribed spacer (ITS) region of ribosomal (r)DNA was identified as being ideal for amplification as it was a well conserved region in strongylid nematodes (Romstad et al., 1997) and had adequate sequence difference between species for delineation. Genetic markers in ITS-2 rDNA were used to distinguish *Necator americanus* from *Oesophagostomum bifurcum* (Romstad et al., 1997) and later de Gruijter et al. (2005) used it successfully to differentiate *N. americanus* and *A. duodenale* in fecal samples.

The tropical climate in Brazil is highly favorable to the development and transmission of hookworm and other STH and many independent studies have confirmed this (Mudenda et al., 2012). The hookworm prevalence varies based on environmental factors and socio-economic status and prevalence can range from less than 1% among city dwellers with good sanitation and nutrition (Santos et al., 2013) to over 80% (Gonçalves et al., 1990) in poorer rural communities.

The objectives of this study were to use a flotation method to determine the prevalence of soil transmitted nematodes in Mutuípe, a rural municipality in Bahia state, in Northeast Brazil and to determine the hookworm species present.

3.2 Materials and Methods

3.2.1 Study Area

Mutuípe municipality (latitude -13.206189, longitude -39.502773), located in Bahia state, Brazil, has a population of approximately 22000 people. Patients were referred to the Central Municipality Laboratory from six health posts scattered around the municipality namely: L'dio Santana, Teobaldo Pinheiro, Antonio Ribeiro leal, Centro (Moisés Gongalves I, Moisés Gongalves II) and Cajazeira. Mutuípe is a rural municipality with farming as the main occupation and contains a small city, Mutuípe, where the central health laboratory was located. The age, gender and location i.e. rural/urban were noted for each person. The study was done in tandem with other routine laboratory investigations of the Central Laboratory, Ministry of Health (Laboratório do centro de saúde, Ministério da Saúde). Authorization was obtained from the Health Secretary of the municipality and verbal consent of participants as part of the project "Recurrent Treatment: Effect on rural and urban *S. mansoni* population structure". (Exemption approval in appendix for IRB #EM-13-39) (Appendix C).

3.2.2 Parasitological Survey

The survey was carried out over a period of six days in July, 2012. All fecal samples that were submitted to the laboratory during that time were examined for STH eggs. Patients were routinely required to submit three fecal samples on consecutive days regardless of presenting complaint/symptoms. Fecal flotation was performed on 2g of feces using a the

double centrifugation technique (Baker, 2007; Sloss et al., 1994) with a sucrose solution of 1.27 specific gravity. Two grams of feces were placed in a 15 ml conical tube and filled with water up to the 10ml mark. The feces were homogenized with the water using a small kebab wooden stick, then the tube was filled with water and a cap screwed on. The tubes were centrifuged for 5 min at 2000 rpm using a fixed head, 80-2B Centribio centrifuge (Curitiba, Brazil) and the supernatant was discarded. The tube was filled with sucrose solution up to the 10ml mark and homogenized as before. The tube were then centrifuged for 5 min at 2000 rpm and then placed upright in a rack. The tube was then filled with sucrose solution till a positive meniscus formed and a cover slip was place on top of the tube. The tube was left to stand for a minimum of 20 min before the cover slip was removed, placed on a glass slide and examined by light microscope (Nikon YS100, Nikon, Melville, NY). The fecal egg count was recorded and divided by two to obtain the eggs per gram (epg) for each parasite species seen.

3.2.3 Larvae Culture and Examination

One gram of feces from each hookworm positive sample was cultured for six to nine days using the Harada-Mori method (Harada and Mori, 1955)(Figure 3.1). Briefly, a gram of feces was smeared on a 1.5 x 11cm filter paper strip (medium porosity, slow flow rate, P5 grade, Fisher Scientific, Pittsburgh, PA); 4ml of filtered water was placed in a 15 ml tube and the filter paper was inserted into the tube till the water was almost at the level of the fecal smear but not touching it so that the filter paper did not reach the bottom of the tube. The cap was then placed loosely on the tube. Cultures were done in triplicate at room temperature (23-24°C) for 1-6 days before they could be moved into a 29°C incubator at the Oswaldo Cruz (FIOCRUZ) laboratory in Salvador, Bahia. The culture

tubes were removed from the incubator and the filter paper carefully removed and discarded into a biohazard bag. The contents of the three tubes from each sample were combined into one tube and centrifuged at 2000rpm for 5 minutes to concentrate the larvae. The supernatant was aspirated with a plastic pipette leaving 1ml of water. The larvae pellet was resuspended and 0.1ml of the suspension was placed on a glass slide with a cover slip and examined under a light microscope at 10x power. The larvae present were measured and the species identified morphologically using published keys and descriptions (Nichols, 1956; Svensson and Kessel, 1926; Wu and Peng, 1965). The remainder of the suspension was placed in a well of a 24 well plate and the larvae were counted using an inverted microscope at 4x magnification, then placed in 1.5ml eppendorf tubes and stored at -20°C until DNA extraction was performed.



Figure 3.1: Harada-Mori culture. Infective hookworm larvae accumulate at the bottom of the tube.

3.2.4 Preparation and Storage of Fecal Samples

A 1-2 grams sample of individual hookworm positive fecal submissions were mixed thoroughly with 1% saline, strained through a sieve (tea strainer) and the filtrate was centrifuged at 1500rpm for 5 minutes. The supernatant was discarded and the pellet in the 15ml tube was stored at -20°C until DNA extraction could be done. A random sample of 29 hookworm negative samples were similarly processed individually.

3.2.5 DNA Extraction from Infective Larvae and Fecal Samples

Eppendorf tubes (1.5ml) containing infective larvae in water were centrifuged at 2000 rpm for 5 minutes and the water removed leaving the last 0.1ml in the tube. The tubes were gently vortexed to resuspend the pellet. DNA was extracted as per manufacturer's instructions using the QiaAmp® DNA Mini kit (Qiagen, Vinlo, Netherlands). A freeze-thaw cycle using dry ice and 56°C was incorporated in the incubation part of the protocol to enhance cell lysis with Proteinase K. DNA was extracted from the frozen fecal samples using the QiaAmp® DNA Stool Mini kit (Qiagen, Vinlo, Netherlands) as per manufacturer's instructions. The DNA was stored at 4°C until further examination.

3.2.6 PCR for *N. americanus* and *A. duodenale* Differentiation

PCR reactions were performed in 25µl volumes using 10.5µl of fecal or hookworm larvae DNA sample, 12.5µl of 2X AmpliTaq Gold® PCR Master Mix (AmpliTaq Gold DNA Polymerase, 250U; GeneAmp PCR Gold Bugger, 30 mM Tris/HCl, pH 8.05, 100mM KCl; dNTP, 400 µM each; MgCl₂, 5mM, proprietary stabilizers) (Life Technologies, Coralville, IA) and 1µl (0.4µM) of each primer. A semi-nested PCR protocol was followed (de Gruijter et al., 2005) using the forward primer NC1 (5' - ACGTCTGGTTCAGGGTTGTT-3') and reverse primer NC2 (5'-

TTAGTTTCTTTTCCTCCGCT-3') (Gasser et al., 1993) for the first round of amplification in a Bio-Rad PTC 0200 DNA Engine Thermal Cycler (Hercules, CA). In brief, amplification consisted of 94°C for 5 min, 25 cycles at 94°C for 30s; 55°C for 30s; 72°C for 30s, followed by a final extension step of 5 min at 72°C. The amplicon (2µl) from the first round of amplification was used in the second amplification procedure using forward primer AD1 (5'-CGACTTTAGAACGTTTCGGC-3') and NC2 as the reverse primer. The second round of amplification consisted of 94°C for 5 min, 35 cycles at 94°C for 1 min; 55°C for 1 min, 72°C for 1 min, and finally 72°C for 5 min. Primers were obtained from Integrated DNA Technologies (Coralville, IA). 5µl of each second-round amplicon was mixed with 1µl of 6X TrackIt™ Cyan/Yellow Loading Buffer (Invitrogen, Carlsbad CA) and examined on 1% agarose-TBE (IM Tris, 0.9M Boric acid, 0.01M EDTA, pH 8.4) (Invitrogen, Carlsbad, CA) gel with SYBR® Safe gel stain (TrackIt™, Invitrogen, Carlsbad, CA) with an Ultraviolet (UV) transilluminator (Fluorchem®Q, Alpha Innotech, San Leandro, CA).

PCR was performed first on the larvae DNA samples, and their amplicons were sequenced by the GenLab, School of Veterinary Medicine, Louisiana State University, using a 3130 Genetic Analyzer (Applied Biosystems®/Hitachi, Life Technologies, Coralville, IA). Sequence analysis was done using MEGA 5.2 software (Tamura et al., 2011) and similarity comparison was made against the GenBank nucleotide database using BLAST (NCBI). Those confirmed to be *N. americanus* were subsequently used as positive control samples for fecal DNA PCR. Molecular biology grade water was used as a negative control. All amplicons that appeared dissimilar to the *N. americanus* expected

base pair size on the agarose gel were sequenced and searched against the GenBank database.

Statistical analysis was done using Statistical Analysis Software 9.3 (SAS, Cary, NC). Chi-square test and Fisher's Exact test were used to compare categorical data where appropriate. A p value < 0.05 was considered significant.

3.3 Results

A total of 159 patients referred from the five health posts in Mutuípe municipality submitted fecal samples (Table 3.1). Of these, only 38 patients submitted three fecal samples during the survey period, therefore 277 fecal samples were collected and examined (Table 3.2). Patient age ranged from 3 months to 96 years old.

Table 3.1: Prevalence of parasites in out-patients from health posts of Mutuípe municipality.

Health Post	Number of patients	Parasite positive (%)	Hookworm (%)	<i>A. lumbricoides</i> (%)	<i>T. trichiura</i> (%)
Centro	51	11 (21.6)	8 (15.7)	4 (7.8)	4 (7.8)
Cajazeira	33	6 (18.1)	4 (12.1)	1 (3.1)	2 (6.2)
A. R. Leal	18	3 (16.7)	2 (11.1)	0	2 (11.1)
T. Pinheiro	27	11 (40.7)	10 (37.0)	1 (3.8)	1 (3.8)
L. Santana	30	7 (23.3)	7 (23.3)	0	0
Total	159	38 (23.9)	31 (19.5)	6 (3.8)	9 (5.7)

There were 61 males and 98 females and 20 (32.8%) males and 18 (18.4%) females had parasites. There was no significant difference by gender among parasite infected people.

Table 3.2: STH egg detection by number of fecal samples submitted by patients.

Samples submitted	Number of patients	Hookworm	<i>A. lumbricoides</i>	<i>T. trichiura</i>
1	79	19	3	3
2	40	6	1	2
3	38	6	2	4

Most of the infections were single parasite infections including one case with *E. vermicularis*. *A. lumbricoides* tended to occur with other parasites (Table 3.3).

Table 3.3: Occurrence of single and multi-parasite infections of STH in out-patients.

Infections	Parasite	Number of patients
Single	Hookworm	25
	<i>A. lumbricoides</i>	1
	<i>T. trichiura</i>	5
Double	Hookworm, <i>A. lumbricoides</i>	2
	Hookworm, <i>T. trichiura</i>	1
	<i>T. trichiura</i> , <i>A. lumbricoides</i>	0
Triple	Hookworm, <i>A. lumbricoides</i> , <i>T. trichiura</i>	3

The fecal egg counts of the STH in the patients were all of low intensity (Table 3.4) except for one child that had had an *A. lumbricoides* egg count of 8762/g.

Table 3.4: Fecal egg counts of STH in patients

	Hookworm	<i>A. lumbricoides</i>	<i>T. trichiura</i>
Mean EPG	113.8	2307.2	26.3
Range	1-633	1-8762	3-80

With regard to location, 63 (39.6%) of the patients came from urban Mutuípe and 96 (60.4%) came from a rural area (Figure 3.2). Although the number of parasitized patients was higher among rural dwellers (28/96; 29.2%) than those from the urban area (10/63; 15.9%), there was no significant difference between the two groups ($p=0.054$).

Hookworm had the highest prevalence (19.5%) of the nematode eggs seen. *Enterobius vermicularis* was found in one child by fecal examination.




Figure 3.2: Proportions of parasitized and non-parasitized patients in urban and rural Mutuípe

3.3.1 Hookworm Infection

The ages of hookworm positive individuals ranged from 1.5 to 96 years old. The majority of these (70.97%) were adults over 20 years old. There was no significant difference in prevalence between patients less than 20 years old and those over 20 years old and there was very poor correlation between fecal egg count and age ($R=0.065$). Prevalence of hookworm increased with age, as did the mean epg and prevalence was highest in the >50 years age group but the mean epg was highest in the 36-50 years age group (Figure 3.3).

However, significantly more males (17/31; 54.8%) than females (14/31; 45.2%) were infected ($p = 0.035$). The mean hookworm fecal egg count was 110.3 (SD 152.4, Range 0.3-633) (Table 3.5).

There were 63 individuals from urban Mutuípe and 96 from the rural area of the municipality (Figure 3.4). Hookworm prevalence was higher among the rural dwellers (60.4%) compared to the urban dwellers (39.6%) although the difference was not significant ($p=0.08$).




Figure 3.3: Prevalence (A) and Mean hookworm EPG (B) among age groups.




Figure 3.4: Distribution of hookworm positive patients among health posts. *urban; ^rural

Infective larvae were recovered from 18 of 31 cultured fecal samples. The mean length was 667.4 μ m (SD 39.5, Range 571.8 – 738.7 μ m) and they matched the morphological characteristics of *N. americanus* (Table 3.5 and Figure 3.5).

Seventeen out of 18 of the hookworm larvae DNA samples were successfully identified by PCR and confirmed by sequencing. Sixteen samples had *N. americanus* and one had *A. ceylanicum*. PCR on the 31 hookworm positive fecal samples yielded 21 positive for *N. americanus* and one mixed infection with *N. americanus* and *A. ceylanicum*. The total number of samples identified using PCR as *N. americanus* was 23/31 with one sample also having *A. ceylanicum*. All of the fecal samples that were negative with microscopy were negative for hookworm using PCR. The *A. ceylanicum* sample was 99% identical (one nucleotide difference) with a query cover of 100% to GenBank accession number JQ673421.1, a sample originating from canine feces in Malaysia.

Table 3.5: Morphological characteristics of hookworm infective larvae in Mutuípe

Characteristics	<i>Ancylostoma sp.</i>	<i>N. americanus</i>	Mutuípe hookworm (<i>N. americanus</i>)
Length of body	660 μm	590 μm	
Length of sheath	720 μm	660 μm	667.4 μm
Anterior end	blunt	rounded	rounded
Intestine	Anterior end narrower than esophageal bulb No gap between esophagus and intestine	Anterior end as wide as esophageal bulb Gap between esophagus and intestine	Anterior end as wide as esophageal bulb. Gap between esophagus and intestine
Tail	Serrations not clear	Serrations clear	Serrations clear



Figure 3.5. *N. americanus* infective larvae. Arrows point to the gap between esophagus and intestine.

3.4 Discussion

Mutuípe is located in a region where climatic conditions are highly conducive to the development of STH in the environment. Mutuípe falls within the permissive zone for hookworm development with regard to climatic temperature and moisture (Mudenda et al., 2012). The prevalence of parasites, particularly hookworm is comparable to what other researchers have found in similar environments in rural Brazil (Machado et al., 2008; Silva and da Silva, 2010) and in some cases, higher (Cury et al., 1994; Silva et al., 2007). One reason for the higher prevalence may be that this study was based on out-patients while most of the other studies were based on school children or whole communities. However, in comparison to what other studies have found in patients in various regions of Brazil (Santos et al., 2013), the prevalence of hookworm in this study is much higher. The convenience sampling in this study may be a source of bias and further studies are necessary to have a better understanding of the status of the community with regard to parasite prevalence. The method for STH egg detection was also different from that employed by previous researchers in that a flotation technique was used that concentrated the nematode eggs in the quantity of feces examined (Baker, 2007; Dryden et al., 2005). The quantity of fecal sample used (2g) was also much more than what is used in the routine Kato-Katz method (41.7mg), even when three Kato-Katz slides are prepared. The double centrifugation flotation method therefore was more efficient at detecting low intensity infections, which may be more prevalent in areas where deworming is regularly practiced (Stoltzfus et al., 1997). The double centrifugation method used is also more conducive for detecting hookworm eggs that are quickly cleared in a Kato-Katz smear, as the eggs remain intact for hours after the slide is

prepared. The flotation method is also less labor intensive with regard to the number of slides that need to be examined and eggs are easier to detect due to sedimentation of most fecal debris.

N. americanus was the predominant STH among patients in Mutuípe. Since most were low intensity infections (epg < 2000) (Montresor et al., 1998), anemia would be expected to be uncommon if present at all (Stoltzfus et al., 1996; Stoltzfus et al., 1997). Other researchers (Santos et al., 2013) found that low intensity infections may result in lower than normal hemoglobin content even in well-nourished individuals. If low intensity infections went undetected, patients would remain untreated, resulting in continued contamination of soils with hookworm eggs in populated areas. According to this study, the highest prevalence of hookworm and intensity of infection (epg) were found in adults over 35 years old and this finding is consistent with what other researchers have found (Santos et al., 2013; Stoltzfus et al., 1997). Therefore the use of anthelmintics in all age groups in this region is justified. The current policy in the municipality indicates that all people who test positive for parasites with a Kato-Katz method, be treated for free (Jackson Costa, personal communication). However, the method used in the municipality was spontaneous sedimentation, which is not ideal for recovery of hookworm, and performed less efficiently than other methods (Inês et al., 2011). This underlines the need for an effective and sensitive method for detecting low intensity parasite infections. According to WHO recommendations, MDA should be used in areas where prevalence of STH and Schistosomiasis exceeds 50%. However, with regard to hookworm infection, if prevalence is over 20% and anemia is prevalent, then the whole community should be treated (Montresor et al., 1998). Future studies can therefore focus determining the

prevalence of hookworm and other STH in the general population, including adults and including assessment of the nutritional status and hemoglobin concentrations in order to determine the best approach to control.

Significantly more males than females were infected with hookworm, a finding comparable to other studies (Behnke et al., 2000; Santos et al., 2013) and has been attributed to males being involved in occupations or behaviors that put them at greater risk of infection, such as agriculture (Behnke et al., 2000; Stoltzfus et al., 1997).

Hookworm was found to be more prevalent in patients from rural areas than those living in urban Mutuípe, a finding although not statistically significant, is in agreement with what other researchers have found (Machado et al., 2008; Silva et al., 2007).

The use of a sucrose flotation method normally applied to veterinary parasitology proved to be adequate for the concentration of human intestinal nematode eggs. This method proved more sensitive than a semi-nested PCR procedure for hookworm egg identification. This may be because the amount of feces used in the flotation was much more than the amount from which DNA was extracted (2g and 200mg respectively). Also, the PCR on larvae was more successful (17/18 positive) than on fecal samples (21/31 positive) probably because the larvae samples were 'cleaner' with regard to PCR inhibitors normally found in feces. PCR on fecal samples proved to be faster and less tedious for identification of the species of hookworm than using the Harada-Mori culture method. The low recovery rate of infective larvae from the culture method may also be attributed to the low fecal egg count and the lesser number of culture days that some of

the samples were kept, since it is recommended culture be continued for at least seven days (Navitsky et al., 1998).

The low prevalence of *A. lumbricoides* and *T. trichiura* could be explained by their lifecycle and mode of transmission. They are transmitted orally rather than by skin penetration by *N. americanus*. Therefore good hand hygiene may be deterring the transmission even if they may be present in high numbers in the environment.

The identification of *N. americanus* as the hookworm species in Mutuípe adds to the body of knowledge on the distribution of hookworm species in Brazil. This is the first known report of *A. ceylanicum* in humans in Brazil, a zoonotic hookworm species that is most commonly found in Asia (Carroll and Grove, 1986; Traub et al., 2008).

3.4.1 Conclusions

- The prevalence of STH and hookworm among outpatients was found to be high.
- Adults and males had higher prevalence of hookworm infection compared to other groups.
- *N. americanus* was the predominant hookworm species and one case of *A. ceylanicum* was found.
- Sucrose double centrifugation method proved efficient for the concentration of STH and performed better than the Harada-Mori culture and PCR.

3.5 References

Baker, D.G. 2007. Flynn's Parasites of Laboratory Animals, In: Blackwell Publishing, Ames, Iowa, 1-10.

Basuni, M., Muhi, J., Othman, N., Verweij, J.J., Ahmad, M., Miswan, N., Rahumatullah, A., Aziz, F.A., Zainudin, N.S., Noordin, R., 2011. A pentaplex real-time

- polymerase chain reaction assay for detection of four species of soil-transmitted helminths. *The American Journal of Tropical Medicine and Hygiene* 84, 338-343.
- Beaver, P.C., 1975. Biology of soil-transmitted helminths: the massive infection. *Health Laboratory Science* 12, 116-125.
- Behnke, J.M., De Clercq, D., Sacko, M., Gilbert, F.S., Ouattara, D.B., Vercruysse, J., 2000. The epidemiology of human hookworm infections in the southern region of Mali. *Tropical Medicine & International Health* 5, 343-354.
- Bethony, J., Brooker, S., Albonico, M., Geiger, S.M., Loukas, A., Diemert, D., Hotez, P.J., 2006. Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *The Lancet* 367, 1521-1532.
- Brooker, S., Clements, A., Bundy, D., 2006a. Global epidemiology, ecology and control of soil-transmitted helminth infections. *Adv Parasitol* 62, 221 - 261.
- Brooker, S., Clements, A.C.A., Bundy, D.A.P. 2006b. Global Epidemiology, Ecology and Control of Soil-Transmitted Helminth Infections, In: Simon I. Hay, A.G., David, J.R. (Eds.) *Advances in Parasitology*. Academic Press, 221-261.
- Brooker, S., Peshu, N., Warn, P.A., Mosobo, M., Guyatt, H.L., Marsh, K., Snow, R.W., 1999. The epidemiology of hookworm infection and its contribution to anaemia among pre-school children on the Kenyan coast. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 93, 240-246.
- Carroll, S.M., Grove, D.I., 1986. Experimental infection of humans with *Ancylostoma ceylanicum*: clinical, parasitological, haematological and immunological findings. *Tropical and geographical medicine* 38, 38-45.
- Cury, G.C., Salles, P.G., Reis, M.C., Rego, V.M., Arndt, A.W., de Souza Filho, C.B., Moreira, H.B., de Abreu, M.S., Leite, P.M., 1994. Prevalence of *Schistosoma mansoni* and parasitic intestinal diseases among students of the rural area of the municipality of Jaboticatubas, MG, 1992-1993. *Rev Soc Bras Med Trop* 27, 217-220.
- de Gruijter, J.M., van Lieshout, L., Gasser, R.B., Verweij, J.J., Brienens, E.A.T., Ziem, J.B., Yelifari, L., Polderman, A.M., 2005. Polymerase chain reaction-based differential diagnosis of *Ancylostoma duodenale* and *Necator americanus* infections in humans in northern Ghana. *Tropical Medicine & International Health* 10, 574-580.
- de Silva, N.R., Brooker, S., Hotez, P.J., Montresor, A., Engels, D., Savioli, L., 2003. Soil-transmitted helminth infections: updating the global picture. *Trends in Parasitology* 19, 547-551.
- Dreyfuss, M.L., Stoltzfus, R.J., Shrestha, J.B., Pradhan, E.K., LeClerq, S.C., Khattry, S.K., Shrestha, S.R., Katz, J., Albonico, M., West, K.P., 2000. Hookworms,

- malaria and Vitamin A deficiency contribute to anemia and iron deficiency among pregnant women in the plains of Nepal. *The Journal of Nutrition* 130, 2527-2536.
- Dryden, M.W., Payne, P.A., Ridley, R., Smith, V., 2005. Comparison of common fecal flotation techniques for the recovery of parasite eggs and oocysts. *Veterinary Therapeutics* 6, 15-28.
- Eppig, C., Fincher, C.L., Thornhill, R., 2010. Parasite prevalence and the worldwide distribution of cognitive ability. *Proceedings of the Royal Society B: Biological Sciences* 277, 3801-3808.
- Ezeamama, A.E., Friedman, J.F., Olveda, R.M., Acosta, L.P., Kurtis, J.D., Mor, V., McGarvey, S.T., 2005. functional significance of low-intensity polyparasite helminth infections in anemia. *Journal of Infectious Diseases* 192, 2160-2170.
- Fleming, F.M., Brooker, S., Geiger, S.M., Caldas, I.R., Corrêa-Oliveira, R., Hotez, P.J., Bethony, J.M., 2006. Synergistic associations between hookworm and other helminth species in a rural community in Brazil. *Trop Med Int Health* 11, 56-64.
- Gasser, R.B., Chilton, N.B., Hoste, H., Beveridge, I., 1993. Rapid sequencing of rDNA from single worms and eggs of parasitic helminths. *Nucleic Acids Research* 21, 2525-2526.
- Geiger, S.M., Massara, C.L., Bethony, J., Soboslay, P.T., Corrêa-Oliveira, R., 2004. Cellular responses and cytokine production in post-treatment hookworm patients from an endemic area in Brazil. *Clin Exp Immunol* 136, 334-340.
- Gonçalves, J.F., Tanabe, M., Medeiros, F.d.P.d.M., Gonçalves, F.J., Aca, I.d.S., Motta, S.R.N.d., Tateno, S., Takeuchi, T., 1990. Parasitological and serological studies on amoebiasis and other intestinal parasitic infections in the rural sector around Recife, Northeast Brazil. *Rev Inst Med Trop São Paulo* 32, 428-435.
- Harada, U., Mori, O., 1955. A new method for culturing hookworm. *Yonago Acta Med* 1, 177-179.
- Hotez, P., Brindley, P., Bethony, J., King, C., Pearce, E., Jacobson, J., 2008. Helminth infections: the great neglected tropical diseases. *J Clin Invest* 118, 1311 - 1321.
- Hotez, P., Molyneux, D., Fenwick, A., Kumaresan, J., Erlich Sachs, S., Sachs, J., Savioli, L., 2007. Control of neglected tropical diseases. *N Engl J Med* 357, 1018 - 1027.
- Hotez, P.J., Bethony, J., Bottazzi, M.E., Brooker, S., Buss, P., 2005. Hookworm: "The Great Infection of Mankind". *PLoS Med* 2, e67.
- Inês, E.d.J., Souza, J.N., Santos, R.C., Souza, E.S., Santos, F.L., Silva, M.L.S., Silva, M.P., Teixeira, M.C.A., Soares, N.M., 2011. Efficacy of parasitological methods

for the diagnosis of *Strongyloides stercoralis* and hookworm in faecal specimens. *Acta Tropica* 120, 206-210.

- Jardim-Botelho, A., Raff, S., DeÁvila Rodrigues, R., Hoffman, H.J., Diemert, D.J., Corrêa-Oliveira, R., Bethony, J.M., Gazzinelli, M.F., 2008. Hookworm, *Ascaris lumbricoides* infection and polyparasitism associated with poor cognitive performance in Brazilian schoolchildren. *Tropical Medicine & International Health* 13, 994-1004.
- Kobayashi, J., Hasegawa, H., Forli, A.A., Nishimura, N.F., Yamanaka, A., Shimabukuro, T., Sato, Y., 1995. Prevalence of intestinal parasitic infection in five farms in Holambra, São Paulo, Brazil. *Rev Inst Med Trop São Paulo* 37, 13-18.
- Levecke, B., Behnke, J.M., Ajampur, S.S., Albonico, M., Ame, S.M., Charlier, J., Geiger, S.M., Hoa, N.T., Kamwa Ngassam, R.I., Kotze, A.C., McCarthy, J.S., Montresor, A., Periago, M.V., Roy, S., Tchuem Tchuente, L.A., Thach, D.T., Vercruysse, J., 2011. A comparison of the sensitivity and fecal egg counts of the McMaster egg counting and Kato-Katz thick smear methods for soil-transmitted helminths. *PLoS neglected tropical diseases* 5, e1201.
- Machado, E.R., de Souza, T.S., da Costa, J.M., Costa-Cruz, J.M., 2008. Enteroparasites and commensals among individuals living in rural and urban areas in Abadia dos Dourados, Minas Gerais state, Brazil. *Parasitologia Latinoamericana* 63, 34-39.
- Montresor, A., Crompton, D.W.T., Hall, H., Bundy, D.A.P., Savioli, L. 1998. Guidelines for the evaluation of soil-transmitted helminthiasis and schistosomiasis at community level. A guide for control programme managers. (Geneva, World Health Organisation.).
- Mudenda, N.B., Malone, J.B., Kearney, M.T., Mischler, P., Nieto, P., McCarroll, J.C., Vounatsou, P., 2012. Modelling the ecological niche of hookworm in Brazil based on climate. *Geospatial Health* 6, 111-123.
- Navitsky, R.C., Dreyfuss, M.L., Shrestha, J., Khatry, S.K., Stoltzfus, R.J., Albonico, M., 1998. *Ancylostoma duodenale* is responsible for hookworm infections among pregnant women in the rural plains of Nepal. *The Journal of Parasitology* 84, 647-651.
- Nichols, R.L., 1956. The etiology of visceral Larva migrans. II. Comparative larval morphology of *Ascaris lumbricoides*, *Necator americanus*, *Strongyloides stercoralis* and *Ancylostoma caninum*. *Journal of Parasitology* 42, 363-399.
- Okazaki, M., Okazaki, M., Miranda, P., Neto, J., Diegues, V., Alves, J., Cauas, M., Tanabe, M., Kobayashi, S., Kaneko, N., 1988. Parasitological and serological studies on amoebiasis and other intestinal parasitic infections in Recife and its suburban area, Northeast Brazil. *Revista do Instituto de Medicina Tropical de São Paulo* 30, 313-321.

- Romstad, A., Gasser, R.B., Monti, J.R., Polderman, A.M., Nansen, P., Pit, D.S.S., Chilton, N.B., 1997. Differentiation of *Oesophagostomum bifurcum* from *Necator americanus* by PCR using genetic markers in spacer ribosomal DNA. *Molecular and Cellular Probes* 11, 169-176.
- Santos, F.L.N., Souza, A.M.G.C.d., Soares, N.M., 2013. Hookworm and threadworm infections and their association with hemoglobin and eosinophil concentrations in residents of Salvador-Bahia, Brazil. *Revista do Instituto de Medicina Tropical de São Paulo* 55, 233-238.
- Silva, L.P., da Silva, R.M.G., 2010. Occurrence of enteroparasites in centers education child of the municipality of Patos of Minas, MG, Brazil. *Bioscience J* 26, 147-151.
- Silva, R.d.C.R., Barreto, M.L., Assis, A.M.O., de Santana, M.L.P., Parraga, I.M., Reis, M.G., Blanton, R.E., 2007. The relative influence of polyparasitism, environment, and host factors on schistosome Infection. *Am J Trop Med Hyg* 77, 672-675.
- Sloss, M.W., Kemp, R.L., Zajac, A.M., 1994. *Veterinary Clinical Parasitology*, 6th Edition. Iowa State University Press, Ames, IA.
- Stephenson, L.S., Latham, M.C., Adams, E.J., Kinoti, S.N., Pertet, A., 1993. Physical Fitness, Growth and Appetite of Kenyan School Boys with Hookworm, *Trichuris trichiura* and *Ascaris lumbricoides* Infections are Improved Four Months after a Single Dose of Albendazole. *The Journal of Nutrition* 123, 1036-1046.
- Stoltzfus, R.J., Albonico, M., Chwaya, H.M., Savioli, L., Tielsch, J., Schulze, K., Yip, R., 1996. Hemoquant determination of hookworm-related blood loss and its role in iron deficiency in african children. *Am J Trop Med Hyg* 55, 399-404.
- Stoltzfus, R.J., Dreyfuss, M.L., Chwaya, H.M., Albonico, M., 1997. Hookworm control as a strategy to prevent iron deficiency. *Nutrition Reviews* 55, 223-332.
- Svensson, R.M., Kessel, J.F., 1926. Morphological Differences between *Necator* and *Ancylostoma* Larvae. *The Journal of Parasitology* 13, 146-153.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011. MEGA5: Molecular Evolutionary Genetics Analysis Using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Molecular Biology and Evolution* 28, 2731-2739.
- Traub, R.J., Inpankaew, T., Sutthikornchai, C., Sukthana, Y., Thompson, R.C.A., 2008. PCR-based coprodiagnostic tools reveal dogs as reservoirs of zoonotic ancylostomiasis caused by *Ancylostoma ceylanicum* in temple communities in Bangkok. *Veterinary Parasitology* 155, 67-73.
- WHA 2001. Resolution WHA54.19: Schistosomiasis and soil-transmitted helminth infections.

WHO 2013. Soil-transmitted helminth infections. In Fact sheet No. 366.

Wu, Z.X., Peng, J.M., 1965. Studies on the morphological differentiation of the infective larvae of *Ancylostoma duodenale* and *Necator americanus*. Acta Parasitol Sin 2, 280-290.

Yap, P., Du, Z.-W., Chen, R., Zhang, L.-P., Wu, F.-W., Wang, J., Wang, X.-Z., Zhou, H., Zhou, X.-N., Utzinger, J., Steinmann, P., 2012. Soil-transmitted helminth infections and physical fitness in school-aged Bulang children in southwest China: results from a cross-sectional survey. Parasites & Vectors 5, 50.

4 GASTROINTESTINAL PARASITE SURVEY OF SHELTER DOGS IN SOUTHERN LOUISIANA

4.1 Introduction

Investigation of intestinal parasites in dogs is important in order to evaluate the risk to dog populations, estimate the level of environmental contamination and to assess the public health risk from zoonotic parasites. National surveys have shown that shelter dogs tend to be more parasitized than pet dogs (Blagburn et al., 2012; Little et al., 2009) due to inadequate veterinary care. Hookworm was found to be the most prevalent parasite in shelter dogs using flotation methods, particularly in the southeastern states of the USA. (Blagburn et al., 1996; Hoskins et al., 1982; Little et al., 2009). The last Louisiana survey was a retrospective study of Louisiana State University (LSU) Teaching Hospital records (Hoskins et al., 1982), therefore the level of parasitism in shelter dogs in Louisiana requires further investigation. For example, limitations in the egg/oocyst/larvae concentration methods used previously may have led to an underestimation of the prevalence of parasites such as tapeworms and *Giardia* sp. and, in order to enhance visibility and recovery of parasite eggs/oocysts, a double centrifugation flotation technique is now recommended (Liccioli et al., 2012; Sloss et al., 1994).

The most common species of canine hookworm in the US is *A. caninum*. *A. braziliense* which is the main cause of cutaneous larva migrans (CLM) in humans, has been found in Texas and Florida states in both dogs and cats (Liotta et al., 2012). However, the prevalence of *A. braziliense* in Louisiana dogs is unknown. Differentiating the eggs of hookworm species by microscopy is challenging as they are very similar in size, except for *Uncinaria stenocephala* which is larger and is more prevalent in the northern US

states (Blagburn et al., 1996; Lucio-Forster et al., 2012). Molecular methods developed for differentiating *Ancylostoma* spp. have proved useful and are less time consuming than microscopy (e Silva et al., 2006; Liotta et al., 2012).

The objectives of this study were to determine the prevalence of hookworm and other intestinal parasites in shelter dogs and to determine the relationship of sex, age and deworming status with parasite infection, as well as determine the species of hookworm in south Louisiana. The anthelmintic protocols used in animal shelters were also evaluated.

4.2 Methods

The study area was southern Louisiana and involved 10 animal shelters in nine (9) parishes, namely: Ascension, Acadia (Acadia, Crowley shelters), East Feliciana (Dixon Correctional Institute shelter), East Baton Rouge, Livingston (Walker Animal Control), Iberville, Tangipahoa, Pointe Coupee and St. Martin (Figure 1). Most of the dogs were kept in the shelters a minimum of five days with an average length of stay of two to three weeks, except in East Feliciana and Point Coupee shelters which had a ‘no-kill’ policy where animals may have been kept for years.

Fecal samples were collected before the daily cleaning from cages of individually housed dogs that had been acquired within the previous two weeks. Samples were placed in plastic containers and labeled with age, sex, breed and deworming status of the respective dog. The collection was opportunistic during the LSU Shelter Medicine course, at the time of routine visits of students and shelter medicine clinical faculty to animal shelters. Samples were submitted to the Louisiana Animal Disease Diagnostic Laboratory

(LADDL) and examined the same day. The age was determined from shelter records, where available, or estimated from examination of the dentition. The last known date of deworming was recorded if known.




Figure 4.1. Map of Louisiana with the location of parishes in the study area in green. Acadia (1), Pointe Coupee (2), St. Martin (3), Iberville (4), East Baton Rouge (5), Ascension (6), Livingston (7), Tangipahoa (8), East Feliciana (9).

4.2.1 Parasitological Examination:

A direct saline fecal smear, zinc sulfate flotation (SG = 1.18), sucrose flotation (SG = 1.25-1.27) and saline sedimentation (0.85%) (Baker, 2007) were used to process each sample. The modified Wisconsin double centrifugation method, which included an initial wash with water, was used for the flotation methods (Rehbein et al., 2011; Sloss et al., 1994). The saline sedimentation method involved three washes, allowing the tube stand for 20 minutes between washes. The supernatant was decanted, leaving the last 15mls of sediment before filling the tube with more saline. During the last decantation, a transfer

pipette was used and the sediment was then drawn into the pipette which was left upright for five minutes to allow sedimentation in the pipette. The sediment was then placed on a glass slide and examined under a light microscope. Slides were examined at 10x and 40x for morphological identification of eggs/oocysts/larvae. A gram of feces was used for each of the flotation and the sedimentation methods. Eggs per gram counts were noted for helminth eggs found by the flotation methods.

4.2.2 PCR and Restriction Fragment Length Polymorphism (RFLP) for Hookworm Speciation

Hookworm positive fecal samples were cultured using the Harada-Mori technique (Harada and Mori, 1955) and the resulting infective larvae were examined microscopically to identify *Ancylostoma* spp. and then stored at -20°C awaiting DNA extraction. For samples that did not have sufficient fecal matter for culturing after the parasitological fecal examinations were conducted, the sediment remaining from the sedimentation method was placed in 1.5ml eppendorf tubes and frozen. DNA was extracted from larvae using a QiaAmp® DNA Mini kit (Qiagen, Valencia, CA) using the protocol for DNA extraction from tissue samples as per manufacturer's instructions. A freeze-thaw cycle was incorporated during the incubation period with Proteinase K using liquid nitrogen and 56°C. DNA was extracted from fecal samples using a QiaAmp® DNA Stool Mini kit (Qiagen, Valencia, CA) and stored at -20°C until examined.

Two PCR protocols were performed using two sets of primers. In the first protocol, the forward primer NC5 (5'-GTAGGTGAACCTGCGGAAGGATCATT-3') and reverse primer NC2 (5'-TTAGTTTCTTTTCCTCCGCT-3') (Gasser et al., 1996) were used.

These primers span the ITS-1, 5.8S and ITS-2 regions of the ribosomal DNA of

Ancylostoma spp. resulting in a PCR product approximately 860 base pairs in size (e Silva et al., 2006). PCR reactions were performed in 25µl volumes using 10.5µl of fecal or hookworm larvae DNA sample, 12.5µl of 2X AmpliTaq Gold® PCR Master Mix (AmpliTaq Gold DNA Polymerase, 250U; GeneAmp PCR Gold Buffer, 30 mM Tris/HCl, pH 8.05, 100mM KCl; dNTP, 400 µM each; MgCl₂, 5mM, proprietary stabilizers) (Life Technologies, Coralville, IA) and 1µl (0.4µM) of each primer. The amplification protocol consisted of 95°C for 5 min, 39 cycles of 95°C for 30s; 50°C for 30s; 72°C for 1.5min and a final extension step of 8 min at 72°C. RFLP was then performed with 0.6µl endonuclease *Hinf*I (New England BioLabs® Inc, Beverly, MA) and 5µl of PCR amplicon DNA in a total volume of 10µl at 37°C for an hour. The reaction was stopped by incubation at 80°C for 20 mins. 5µl of each PCR-RFLP product was mixed with 1µl of 6X TrackIt™ Cyan/Yellow Loading Buffer (Invitrogen, Carlsbad CA) and examined on 3% agarose-TBE (IM Tris, 0.9M Boric acid, 0.01M EDTA, pH 8.4) gel (Invitrogen, Carlsbad, CA) with SYBR® Safe gel stain (TrackIt™, Invitrogen, Carlsbad, CA). The gel was read with an Ultraviolet (UV) transilluminator (Fluorchem®Q, Alpha Innotech, San Leandro, CA).

The second PCR protocol involved the use of forward primer NC1 (5' - ACGTCTGGTTCAGGGTTGTT-3') and reverse primer NC2 (Gasser et al., 1993). The reactions were performed in a 25µl volume as above with 1µl of each primer and 10.5µl of DNA. The amplification protocol was as follows: 94°C for 5 min, 35 cycles at 94°C for 1 min; 55°C for 1 min, 72°C for 1 min, and finally 72°C for 5 min (de Gruijter et al., 2005). The PCR product from this reaction was digested as above with *Hinf*I endonuclease.

The PCR products from the two kinds of PCR reactions were sequenced and the samples that were found to contain *A. caninum* were used as controls in the RFLP procedure. *A. braziliense* DNA from eggs was donated by Dr. Dwight Bowman, College of Veterinary Medicine, Cornell University, Ithaca, NY.

Statistical analysis software (SAS 9.3, Cary, NC) was used for statistical analysis. Chi-square or Fisher's exact tests for categorical data comparisons and student's t-test for difference in means were used. A 95% confidence interval was adopted and a *p* value <0.05 was considered significant.

4.3 Results

Of fecal samples from a total of 209 dogs examined, 163 (78%) were found to be parasitized (Table 1). Multiparasitism was present in 79 (37.8%) dogs. The most prevalent parasites found were, in descending order, *Ancylostoma* sp. (53.6%), followed by *Trichuris vulpis*, coccidia of the *Cystoisospora ohioensis* complex, *Giardia duodenalis*, *Cystispora canis*, *Toxocara canis*, *Dipylidium caninum*, *Alaria* spp., *Uncinaria stenocephala* and *Capillaria* spp. *U. stenocephalas* was found in one dog and diagnosed as a mixed infection with *Ancylostoma* spp.

4.3.1 Distribution of Parasites by Parish

Among helminth parasites, Hookworm and *T. vulpis* were recovered from dogs from all the parishes, and the other parasites were found in variable prevalence (Table 4.1). Point Coupee parish had the highest percentage of dogs with parasites (94.1%) and East Feliciana had the least (31.0%). The trend was similar for hookworm, but with regard to *T. vulpis*, Tangipahoa had the highest percentage (44.8%) and St. Martin the least (9.1%).

T. canis and *D. caninum* were found in dogs from five of the nine parishes, *Alaria* in three parishes, *Capillaria* in one parish. Among protozoan parasites, *Cystoisospora* sp. were found in dogs from all the parishes, but individual species differed in distribution, with *C. ohioensis* being most common. *G. duodenalis* was recovered from dogs in seven parishes.

4.3.2 Effect of Age on Prevalence

The age of dogs ranged from 1.5 months to 12 years. There were only 12 dogs over three years old. There were 39 dogs less than six months old and 87.2% of them had parasites compared with 75.9% of those aged six months or more (Table 4.2). There was no statistical difference ($p=0.48$) in prevalence between the two age groups, except the younger (< 6 month) age group had significantly more dogs with *T. canis* ($p = 0.0009$) and *Cystoisospora* sp ($p=0.001$) than the older age group.

Mixed parasite infections were found in 18 (52.9%) dogs less than six months and in 61 (47.3%) of dogs over six months (Table 4.3). Hookworm was the most frequently found single parasite infection (46/163) followed by *T. vulpis* and *Cystoisospora* spp. The most common combination of parasites was with hookworm and *T. vulpis* followed by hookworm and *Cystoisospora* spp. This trend was the same in the two age groups, however, with regard to proportions, the hookworm and *T. vulpis* combination was found more in older dogs (14.7%) as compared to younger dogs (11.8%). The hookworm and *Cystoisospora* spp. combination was more frequently found in younger dogs (8.8%) as compared to older dogs 5.4%. *Capillaria* spp. and *Alaria* spp. were found in mixed infections only in dogs over six months old.

Table 4.1. Prevalence of parasites found in dogs from shelters in nine parishes of south Louisiana

Parish	Acadia	Ascension	Iberville	Point Coupee	Tangipahoa	St. Martin	Livingston	East Baton Rouge	East Feliciana	Total
Total Samples	22	37	26	17	29	11	16	22	29	209
Parasite positive (%)	20 (90.9)	29 (78.4)	24 (92.3)	16 (94.1)	25 (86.2)	8 (72.7)	12 (75.0)	20 (90.9)	9 (31.0)	163 (78.0)
Hookworm (%)	13 (59.1)	23 (62.2)	17 (65.4)	16 (94.1)	21 (72.4)	2 (18.2)	5 (31.3)	13 (59.1)	2 (6.9)	112 (53.6)
<i>Trichuris</i> (%)	9 (40.9)	9 (24.3)	5 (19.2)	2 (11.8)	13 (44.8)	1 (9.1)	7 (43.8)	9 (40.9)	5 (17.2)	60 (28.7)
<i>Toxocara</i> (%)	5 (22.7)	2 (5.4)	2 (7.7)	0	2 (6.9)	2 (18.2)	0	0	0	13 (6.2)
<i>Capillaria</i> (%)	0	0	0	0	0	0	0	1 (4.5)	0	1 (0.5)
<i>D. caninum</i> (%)	0	2 (5.4)	1 (3.8)	0	2 (6.9)	2 (18.2)	0	2 (9.1)	0	9 (4.3)
<i>Alaria</i> (%)	1 (4.5)	0	0	1 (5.9)	0	0	1 (6.3)	0	0	3 (1.4)

Table 4.1. Continued

Parish	Acadia	Ascension	Iberville	Point Coupee	Tangipahoa	St. Martin	Livingston	East Baton Rouge	East Feliciana	Total
<i>Cystisospora.</i> <i>canis</i> (%)	3 (13.6)	1 (2.7)	5 (19.2)	0	0	2 (18.2)	2 (12.5)	3 (13.6)	0	16 (7.7)
<i>Cystoisospora</i> <i>ohioensis</i> <i>complex</i> (%)	3 (13.6)	3 (8.1)	7 (26.9)	2 (11.8)	8 (27.6)	0	7 (43.8)	3 (13.6)	3 (10.3)	36 (17.2)
<i>Giardia</i> sp. (%)	3 (13.6)	0	0	1 (5.9)	1 (3.4)	2 (18.2)	2 (12.5)	15 (68.2)	1 (3.4)	25 (12.0)

Table 4.2. Parasite prevalence in dogs less than 6 months old and more than 6 months old.

Age in months	Sample size	Parasite positive (%)	Hookworm (%)	<i>Trichuris</i> (%)	<i>Toxocara</i> (%)	<i>Cystoisospora</i> (%)	<i>Giardia</i> (%)
< 6	39	34 (87.2)	22 (56.4)	10 (25.6)	6 (15.4)	16 (41.0)	6 (15.4)
> 6	170	129 (75.9)	90 (52.9)	49 (28.8)	7 (4.1)	30 (17.7)	23 (13.5)
<i>p</i> value		0.125	0.695	0.69	0.009	0.001	0.762

4.3.3 Effect of Sex on Prevalence

Fecal samples were obtained from 101 female and 106 male dogs as well as two sets of litters that were housed together. The litters were not included in the analysis based on sex. More female dogs (79.2%) were parasitized compared to male dogs (76.4%), although this difference was not significant ($p=0.629$) (Table 4.4). With regard to individual parasite species, there was no statistical difference between the sexes.

4.3.4 Deworming

Shelter records indicated that 125 of the dogs had been dewormed before the day of fecal collection. The anthelmintics used were pyrantel, fenbendazole or ivermectin, singly or in combination with each other or with heartworm medication. The majority of dogs admitted to the shelters were only treated once, usually upon intake. The time period from deworming to fecal collection ranged from a day to six months with a mean of 24 days. Only 49/125 records provided the date deworming was given.

Table 4.3. Occurrence of single species or combinations of parasite species found in dogs over 6 months and less than 6 months of age.

Parasite	Number	<6months	>6months
Hookworm	46	5	41
<i>Trichuris</i>	14	2	12
<i>Toxocara</i>	3	2	1
<i>Dipylidium</i>	1	1	0
<i>Cystoisospora</i>	13	6	7
<i>Giardia</i>	7	0	7
hookworm + <i>Trichuris</i>	23	4	19
hookworm + <i>Toxocara</i>	1	0	1
hookworm + <i>Dipylidium</i>	1	0	1
hookworm + <i>Alaria</i>	1	0	1
<i>Toxocara</i> + <i>Dipylidium</i>	1	0	1
hookworm + <i>Cystoisospora</i>	10	3	7
<i>Trichuris</i> + <i>Cystoisospora</i>	5	0	5
hookworm + <i>Giardia</i>	8	2	6
<i>Trichuris</i> + <i>Giardia</i>	1	0	1
<i>Cystoisospora</i> + <i>Giardia</i>	2	0	2
hookworm + <i>Trichuris</i> + <i>Toxocara</i>	2	1	1
hookworm + <i>Trichuris</i> + <i>Dipylidium</i>	2	1	1
<i>Trichuris</i> + <i>Dipylidium</i> + <i>Capillaria</i>	1	0	1
hookworm + <i>Trichuris</i> + <i>Giardia</i>	4	0	4
hookworm + <i>Trichuris</i> + <i>Cystoisospora</i>	3	0	3
hookworm + <i>Cystoisospora</i> + <i>Toxocara</i>	2	2	0
<i>Trichuris</i> + <i>Cystoisospora</i> + <i>Toxocara</i>	1	0	1
hookworm + <i>Giardia</i> + <i>Toxocara</i>	1	0	1
hookworm + <i>Cystoisospora</i> + <i>Dipylidium</i>	2	1	1
hookworm + <i>Cystoisospora</i> + <i>Giardia</i>	2	1	1
<i>Toxocara</i> + <i>Cystoisospora</i> + <i>Giardia</i>	2	1	1
hookworm + <i>Trichuris</i> + <i>Giardia</i> + <i>Cystoisospora</i>	1	1	0
hookworm + <i>Trichuris</i> + <i>Cystoisospora</i> + <i>Alaria</i>	2	0	2
hookworm + <i>Trichuris</i> + <i>Giardia</i> + <i>Cystoisospora</i> + <i>Dipylidium</i>	1	1	0
Total number of dogs	163	34	129

Table 4.4. Parasite prevalence in male and female dogs.

Sex	Sample size	Parasite positive (%)	Hookworm (%)	<i>Trichuris</i> (%)	<i>Toxocara</i> (%)	<i>Cystoisospora</i> (%)	<i>Giardia</i> (%)
Male	106	81 (76.4)	54 (50.9)	28 (26.4)	6 (5.7)	21 (19.8)	12 (11.3)
Female	101	80 (79.2)	55 (54.5)	30 (29.7)	7 (6.9)	24 (23.7)	15 (14.8)
<i>p</i> value		0.629	0.613	0.598	0.706	0.491	0.451

Of the dogs that were dewormed, helminth eggs were found in 59.8% when compared to 79.8% of untreated dogs and this difference was statistically significant ($p=0.0013$) (Table 4.5). With regard to parasite species, dewormed dogs had significantly less hookworm positive diagnoses ($p=0.0002$) than untreated dogs. The difference in the hookworm prevalence was pronounced in dogs treated with pyrantel compared to untreated dogs ($p=0.0002$) (Table 4.6). Upon comparison of fecal egg counts for dogs treated with pyrantel and untreated dogs, there was no statistical difference, although the treated group had a lower mean EPG (Table 4.7). There was no statistical difference with regard to the other helminths.

Table 4.5. Parasitism in shelter dogs based on history of deworming.

	Sample size	Helminth Positive (%)	Hookworm (%)	<i>Trichuris</i> (%)	<i>Toxocara</i> (%)	<i>D. caninum</i> (%)
Dewormed	125	75 (59.8)	54 (43.2)	36 (28.8)	6 (4.8)	6 (66.7)
Not dewormed	84	67 (79.8)	58 (69.0)	23 (27.4)	7 (8.3)	3 (33.3)
<i>p</i> value		0.0013	0.0002	0.823	0.300	

Table 4.6. Prevalence of parasites in dogs with a history of anthelmintic treatment at shelters.

Anthelmintic	Dewormed dogs	Helminth positive (%)	Hookworm (%)	<i>T. vulpis</i> (%)	<i>T. canis</i> (%)	<i>D. caninum</i> (%)
Pyrantel	84	57 (67.9)	45 (53.6)	31 (36.9)	3 (3.6)	4 (4.8)
Fenbendazole	14	4 (28.6)	2 (14.3)	0	2 (14.3)	2 (14.3)
Ivermectin	1	1 (100.0)	0	0	1 (100.0)	0
Combinations*	26	10 (38.5)	6 (23.1)	5 (19.2)	0	0
Total	125	72 (56.7)	53 (41.7)	36 (28.3)	6 (4.7)	6 (4.7)

*Combinations included: pyrantel + ivermectin (9); pyrantel + moxidectin (Advantage multi®) (2); fenbendazole + ivermectin + moxidectin (Advantage multi®) (13); moxidectin (Advantage multi®) + ivermectin (2)

Table 4.7. Occurance of hookworm and *Trichuris* in dogs treated with pyrantel.

	Samples	Helminths %	Hookworm %	<i>Trichuris</i> %	Mean HW epg
Dewormed	84	50 (59.1)	46 (54.8)	31 (36.9)	2034.1
Untreated	82	67 (81.7)	58 (70.7)	23 (28.1)	4528.2
<i>p</i> value		0.059	0.033	0.223	0.057

4.3.5 Speciation of Hookworm by PCR

DNA was successfully amplified with NC5/NC2 primers in 41 out of 88 (Table 4.8) samples and resulted in a ~860 base pair (bp) size amplicon (Figure 4.3). RFLP resulted in five main fragment sizes which matched those of *A. caninum* (e Silva et al., 2006). The

five fragments were ~ 222, 189, 146, 129 and 89 bp in size. Sequencing of the undigested PCR product from seven samples with similar PCR-RFLP products confirmed that the species was *A. caninum*. The rest of the samples tested resulted in non-specific amplification.

PCR using the NC1/NC2 primer set resulted in an amplicon ~300 bp in size. Of the 88 samples that were tested, DNA was successfully amplified in 75 of them and RFLP resulted in three main fragment sizes (128, 87 and 73 bp) (Figure 4.3) which matched *A. caninum* control samples. The *A. braziliense* DNA was successfully amplified with both primer sets and resulted in similar size amplicons as *A. caninum*. Digestion with endonuclease *Hinf*I of the PCR product from the NC5/NC2 primer set reaction, resulted in five clear bands (396, 308, 222, 145 and 125 bp)(Figure 4.2) while digestion of the NC1/NC2 PCR product resulted in two clear bands or fragments (176 and 127 bp) (Figure 4.2 and 4.4). However, the PCR product from the NC5/NC2 primer set was not always consistent both with *A. caninum* and *A. braziliense*, sometimes resulting in additional bands (Figure 4.4). RFLP of the PCR product with additional bands using NC5/NC2 primers produced different results (Figure 4.4) than when a single band amplicon was digested.

Table 4.8. Comparison of PCR-RFLP protocols for speciation of *A. caninum* from 88 samples by size of PCR amplicon and number of fragments generated from endonuclease digestion.

Primer sets	Amplicon (bp)	Positive samples (%)	<i>Hinf</i>I-RFLP fragments
NC5/NC2	~860	41 (46.6)	5
NC1/NC2	~280	75 (85.2)	3




Figure 4.2. PCR-RFLP results for the *A. braziliense* using primer sets NC1/NC2 and NC5/NC2. Lane 1, PCR product using primer set NC5/NC2; Lane 2, PCR product using primer set NC1/NC2; Lane 3 RFLP product from primer set NC5/NC2; Lane 4, RFLP product from primer set NC1/NC2; Lane M, 100bp DNA ladder.




Figure 4.3. PCR-RFLP results for the *A. caninum* using primer sets NC1/NC2 and NC5/NC2. Lane 5, PCR product using primer set NC5/NC2; Lane 2, PCR product using primer set NC1/NC2; Lane 3 RFLP product from primer set NC1/NC2; Lane 4, RFLP product from primer set NC5/NC2; Lane M, 100bp DNA ladder.




Figure 4.4. PCR-RFLP results for *A. braziliense* showing non-specific amplification using NC5/NC2 primer set in Lane 9; Lane 10, PCR product using primer set NC1/NC2; Lane 11, RFLP product with unrecognizable fragment sizes from PCR product of primer set NC5/NC2; Lane 12, RFLP result from primer set NC1/NC2; Lane M, 100bp DNA ladder.

4.4 Discussion

The prevalence of parasites in South Louisiana shelter dogs was higher than what has been reported in the past in both the 1982 Louisiana survey and a 1996 national survey of shelter dogs (Blagburn et al., 1996; Hoskins et al., 1982). However the trend of finding hookworm in the highest prevalence, followed by *T. vulpis*, was comparable to previous reports. The high prevalence of parasites raises concern for the risk of infection or re-infection of pets and humans due to environmental contamination, particularly from dogs adopted from shelters but also in the homes from which the shelter animals were surrendered. The lack of a significant difference in parasitism between dewormed and untreated dogs indicates that the deworming protocols currently used by the shelters were inadequate and ineffective. This calls for a change in how the deworming is done or the drugs used. Most shelters only treated once and often only adoptable dogs were treated. It

is advisable that the deworming be repeated as per manufacturer's instructions for the medication to be effective i.e. treatment with pyrantel should be repeated after 10-14 days in order to be effective against newly matured adult worms in the intestine. Helminths with migratory larvae or larvae that undergo hypobiosis are a challenge to treat because commonly used dewormers, such as pyrantel, have no effect on larvae stages and require that the dog be retreated 2-4 weeks after the first treatment to eliminate all adult worms (Bowman, 2009).

A previous Louisiana study (Hoskins et al., 1982) found that hookworm prevalence was significantly higher in male dogs but this difference was not observed in this study. The difference between young and older dogs with regard to *T. canis* and *Cystoisospora* spp. was expected considering the life cycle of *T. canis* where infection can be acquired transplacentally and possible acquired immunity to *Cystoisospora* spp. with age. This difference has been observed in other studies (Blagburn et al., 1996; Hoskins et al., 1982; Little et al., 2009).

There was a significant difference in hookworm prevalence between the pyrantel-treated group and the untreated dogs. However, the fecal egg counts were not significantly different but highly suggestive that pyrantel may have been effective. It has been observed in controlled experiments on efficacy against *A. caninum* that when adult worms occur in large numbers, they produce fewer eggs than when adults are in lower numbers in the intestine (Kopp et al., 2007) and it was suggested that a reduction of adult hookworm in the intestines would thus be a better indication of treatment efficacy. Kopp et al. (2007) in Australia also observed an increase in anthelmintic resistance to pyrantel

pamoate by *A. caninum*, a finding that requires confirmatory investigation in other regions of the world.

Alaria spp. were found in three dogs from three parishes. Diagnosis requires demonstration of eggs by sedimentation examination because the eggs, like many other trematode eggs, do not float in common fecal flotation solutions. *Alaria* is acquired through ingestion of intermediate or paratenic hosts like snails and frogs (Taylor et al., 2007). Adults of *Alaria* live in the intestine attached to the mucosa but causes little pathology or clinical signs, and is therefore usually an incidental finding at necropsy. Human infection has been reported and a fatal case of *Alaria* in man was reported due to possible consumption of undercooked frog legs (Fernandes et al., 1976; Freeman et al., 1976). *Alaria* mesocercaria, the infective stage for mammals, have previously been recovered from snakes and frogs in Louisiana and from experimentally infected alligators (Shoop and Corkum, 1981). Therefore precautions should be taken when handling raw meat from wildlife sources and in food preparation using meat from these species, since the potential for human infection does exist.

The prevalence of parasites in the all shelters studied were over 72% except for East Feliciana parish for which only 31% of the dogs were parasitized. This can be attributed to the high level of care that animals in that shelter received as well as being housed in very sanitary conditions. The East Feliciana shelter was also a ‘no-kill’ shelter and administered a combination of anthelmintics, including heartworm preventatives, regularly. They also performed *Giardia* ELISA tests on site. The other shelter (Point Coupee) with a ‘no-kill’ policy had a very high prevalence of parasites even though they too administered a combination of anthelmintics regularly. Dogs in this shelter were let

out in a dirt run where feces were never picked up and this could explain the high number of dogs with hookworm. The St. Martin shelter, which reported good management practices and use of fenbendazole on all the dogs, had a low prevalence of hookworm (18.2%). By contrast, five of the shelters reported deworming only puppies or adoptable dogs, and only upon intake, thereby putting them at risk of infection from dogs that were not dewormed. Other than in the East Feliciana shelter, treatment for coccidiosis or *Giardia* was only given upon diagnosis by the visiting veterinarian. Although overcrowding was not a major finding among the shelters, it can contribute to the prevalence of coccidiosis and *Giardia* infections particularly among puppies and should therefore be avoided. Findings from these shelters indicate that a combination of appropriate and regular deworming as well as maintaining sanitary conditions, like picking up feces in common areas, are essential in preventing intestinal parasite infection in dogs kept in shelters.

It is essential and highly recommended that dogs are dewormed on adoption and the adopting individual be advised to take the dog for a full examination, including a fecal exam, by a veterinarian as soon as possible to prevent parasitic infections from posing a risk to humans and other pets. It was not possible to properly determine drug efficacy as this was not a controlled study, and there was great variation in period between deworming and sample collection as well as inconsistencies in dosage. Further studies are therefore required to determine if drug efficacy is a factor in the high level of parasitism found in shelter dogs. In order to prevent possible development of anthelmintic resistance, it is advisable to deworm based on fecal exams and give appropriate anthelmintic in the recommended dosage (Palmer et al., 2007).

A. caninum was the only hookworm species found in this study by analysis of the 75 of 88 DNA samples that were successfully amplified with the NC5/NC2 primer set. The NC5/NC2 primer set was only able to amplify *Ancylostoma* specific DNA in 41 samples and resulted in much non-specific amplification. DNA amplification may have failed in the other samples due to possible DNA degradation or presence of PCR inhibitors, since the DNA was from eggs in fecal samples. In the samples that did have amplification of the ~860bp target sequence when using the NC5/NC2 primer set, many of them also had additional amplifications that distorted the results of the RFLP, although other researchers have not reported this problem. The results from the NC1/NC2 primer set PCR were consistent and the resultant RFLP fragments, three for *A. caninum* and two for *A. braziliense*, were also consistent, suggesting this method of differentiating *A. caninum* and *A. braziliense* is more optimal. Further investigation should be done to determine if the NC1/NC2 PCR-RFLP can be used to differentiate other *Ancylostoma* species.

4.4.1 Conclusion

- The prevalence of gastrointestinal parasites in Louisiana shelter dogs is higher than what has been reported previously.
- Hookworm is the most prevalent intestinal parasite and *A. caninum* is the predominant hookworm species in shelter dogs.
- *T. canis* and *Cystoisospora* spp. prevalence were significantly higher in dogs <6 months old.
- *Alaria* spp. was reported in dogs from three parishes.
- Anthelmintic protocols for shelters were found to be inadequate and needed to be revised to prevent the spread of gastrointestinal parasites.

- PCR-RFLP using primer set NC1/NC2 was more reliable and efficient than using primer set NC5/NC2.

4.5 References

- Baker, D.G. 2007. Flynn's Parasites of Laboratory Animals, In: Blackwell Publishing, Ames, Iowa, 1-10.
- Blagburn, B.L., Lindsay, D.S., L., V.J., S., R.N., C., W.J., 1996. Prevalence of canine parasites based on fecal flotation. The compendium on continuing education of the practising vet 18, 483-508.
- Blagburn, B.L., Little, S.E., Butler, J.M., Land, T.M., Mount, J.D., Wright, J.C., Griffin, B., Hostetler, J. 2012. National prevalence of parasites in shelter dogs based on centrifugal fecal flotation. In: American Association of Veterinary Parasitologists 57th annual meeting, San Diego, CA, August 4-7, 76.
- Bowman, D.D., 2009. Georgis' Parasitology for Veterinarians, 9th Edition. W. B. Saunders Elsevier, St. Louis, MO.
- de Gruijter, J.M., van Lieshout, L., Gasser, R.B., Verweij, J.J., Brienens, E.A.T., Ziem, J.B., Yelifari, L., Polderman, A.M., 2005. Polymerase chain reaction-based differential diagnosis of *Ancylostoma duodenale* and *Necator americanus* infections in humans in northern Ghana. Tropical Medicine & International Health 10, 574-580.
- e Silva, L.M.C., Miranda, R.R.C., Santos, H.A., Rabelo, É.M.L., 2006. Differential diagnosis of dog hookworms based on PCR-RFLP from the ITS region of their rDNA. Veterinary Parasitology 140, 373-377.
- Fernandes, B., Cooper, J., Cullen, J., Freeman, R., Ritchie, A., Scott, A., Stuart, P., 1976. Systemic infection with *Alaria americana* (Trematoda). Canadian Medical Association Journal 115, 1111.
- Freeman, R., Stuart, P., Cullen, S., Ritchie, A., Mildon, A., Fernandes, B., Bonin, R., 1976. Fatal human infection with mesocercariae of the trematode *Alaria americana*. The American Journal of Tropical Medicine and Hygiene 25, 803.
- Gasser, R.B., Chilton, N.B., Hoste, H., Beveridge, I., 1993. Rapid sequencing of rDNA from single worms and eggs of parasitic helminths. Nucleic Acids Research 21, 2525-2526.
- Gasser, R.B., Stewart, L.E., Speare, R., 1996. Genetic markers in ribosomal DNA for hookworm identification. Acta Tropica 62, 15-21.
- Harada, U., Mori, O., 1955. A new method for culturing hookworm. Yonago Acta Med 1, 177-179.

- Hoskins, J.D., Malone, J.B., Smith, P.H., Uhl, S.A., 1982. Prevalence of parasitism diagnosed by fecal examination in Louisiana dogs. *American Journal of Veterinary Research* 43, 1106-1109.
- Kopp, S.R., Kotze, A.C., McCarthy, J.S., Coleman, G.T., 2007. High-level pyrantel resistance in the hookworm *Ancylostoma caninum*. *Veterinary Parasitology* 143, 299-304.
- Liccioli, S., Catalano, S., Kutz, S.J., Lejeune, M., Verocai, G.G., Duignan, P.J., Fuentealba, C., Ruckstuhl, K.E., Massolo, A., 2012. Sensitivity of double centrifugation sugar fecal flotation for detecting intestinal helminths in coyotes (*Canis latrans*). *Journal of Wildlife Diseases* 48, 717-723.
- Liotta, J.L., Youn, H., Aksel, S., Bienhoff, S.E., Bowman, D.D., 2012. Prevalence of *Ancylostoma braziliense* in Dogs From Alachua and Marion Counties, Florida, United States. *Journal of Parasitology* 98, 1039-1040.
- Little, S.E., Johnson, E.M., Lewis, D., Jaklitsch, R.P., Payton, M.E., Blagburn, B.L., Bowman, D.D., Moroff, S., Tams, T., Rich, L., Aucoin, D., 2009. Prevalence of intestinal parasites in pet dogs in the United States. *Veterinary Parasitology* 166, 144-152.
- Lucio-Forster, A., Liotta, J.L., Yaros, J.P., Briggs, K.R., Mohammed, H.O., Bowman, D.D., 2012. Morphological differentiation of eggs of *Ancylostoma caninum*, *Ancylostoma tubaeforme*, and *Ancylostoma braziliense* from dogs and cats in the United States. *Journal of Parasitology* 98, 1041-1044.
- Palmer, C.S., Traub, R.J., Robertson, I.D., Hobbs, R.P., Elliot, A., While, L., Rees, R., Thompson, R.C.A., 2007. The veterinary and public health significance of hookworm in dogs and cats in Australia and the status of *A. ceylanicum*. *Veterinary Parasitology* 145, 304-313.
- Rehbein, S., Lindner, T., Visser, M., Winter, R., 2011. Evaluation of a double centrifugation technique for the detection of *Anoplocephala* eggs in horse faeces. *Journal of Helminthology* 85, 409-414.
- Shoop, W.L., Corkum, K.C., 1981. Epidemiology of *Alaria marcianae* mesocercariae in Louisiana. *The Journal of Parasitology* 67, 928-931.
- Sloss, M.W., Kemp, R.L., Zajac, A.M., 1994. *Veterinary Clinical Parasitology*, 6th Edition. Iowa State University Press, Ames, IA.
- Taylor, M.A., Coop, R.L., Wall, R.L., 2007. *Veterinary Parasitology*, 3rd Edition. Blackwell Publishing, Oxford, UK.

APPENDIX A

NATURE PUBLISHING GROUP LICENSE TERMS AND CONDITIONS

Nov 07, 2013

This is a License Agreement between Ntombi Mudenda ("You") and Nature Publishing Group ("Nature Publishing Group") provided by Copyright Clearance Center ("CCC"). The license consists of your order details, the terms and conditions provided by Nature Publishing Group, and the payment terms and conditions.

All payments must be made in full to CCC. For payment instructions, please see information listed at the bottom of this form.

License Number	3231490976711
License date	Sep 17, 2013
Licensed content publisher	Nature Publishing Group
Licensed content publication	Nature Reviews Microbiology
Licensed content title	Focus: Soil-transmitted helminthiasis
Licensed content author	L. Savioli and M. Albonico
Licensed content date	Aug 1, 2004
Volume number	2
Issue number	8
Type of Use	reuse in a thesis/dissertation
Requestor type	academic/educational
Format	print and electronic
Portion	figures/tables/illustrations
Number of figures/tables /illustrations	1
High-res required	no
Figures	Figure 1 : Global distribution and control status of soil-transmitted helminthiasis
Author of this NPG article	no
Your reference number	
Title of your thesis / dissertation	Epidemiology and molecular characterization of human and canine hookworm
Expected completion date	Dec 2013
Estimated size (number of pages)	150
Total	0.00 USD
Terms and Conditions	

Terms and Conditions for Permissions

Nature Publishing Group hereby grants you a non-exclusive license to reproduce this

material for this purpose, and for no other use, subject to the conditions below:

1. NPG warrants that it has, to the best of its knowledge, the rights to license reuse of this material. However, you should ensure that the material you are requesting is original to Nature Publishing Group and does not carry the copyright of another entity (as credited in the published version). If the credit line on any part of the material you have requested indicates that it was reprinted or adapted by NPG with permission from another source, then you should also seek permission from that source to reuse the material.
2. Permission granted free of charge for material in print is also usually granted for any electronic version of that work, provided that the material is incidental to the work as a whole and that the electronic version is essentially equivalent to, or substitutes for, the print version. Where print permission has been granted for a fee, separate permission must be obtained for any additional, electronic re-use (unless, as in the case of a full paper, this has already been accounted for during your initial request in the calculation of a print run). NB: In all cases, web-based use of full-text articles must be authorized separately through the 'Use on a Web Site' option when requesting permission.
3. Permission granted for a first edition does not apply to second and subsequent editions and for editions in other languages (except for signatories to the STM Permissions Guidelines, or where the first edition permission was granted for free).
4. Nature Publishing Group's permission must be acknowledged next to the figure, table or abstract in print. In electronic form, this acknowledgement must be visible at the same time as the figure/table/abstract, and must be hyperlinked to the journal's homepage.
5. The credit line should read:
Reprinted by permission from Macmillan Publishers Ltd: [JOURNAL NAME]
(reference citation), copyright (year of publication)
For AOP papers, the credit line should read:
Reprinted by permission from Macmillan Publishers Ltd: [JOURNAL NAME],
advance online publication, day month year (doi: 10.1038/sj.[JOURNAL
ACRONYM].XXXXXX)

Note: For republication from the British Journal of Cancer, the following credit lines apply.

Reprinted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication) For AOP papers, the credit line should read:
Reprinted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME], advance online publication, day month year (doi: 10.1038/sj.[JOURNAL ACRONYM].XXXXXX)

6. Adaptations of single figures do not require NPG approval. However, the adaptation

should be credited as follows:

Adapted by permission from Macmillan Publishers Ltd: [JOURNAL NAME]
(reference citation), copyright (year of publication)

Note: For adaptation from the *British Journal of Cancer*, the following credit line applies.

Adapted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication)

7. Translations of 401 words up to a whole article require NPG approval. Please visit <http://www.macmillanmedicalcommunications.com> for more information. Translations of up to a 400 words do not require NPG approval. The translation should be credited as follows:

Translated by permission from Macmillan Publishers Ltd: [JOURNAL NAME]
(reference citation), copyright (year of publication).

Note: For translation from the *British Journal of Cancer*, the following credit line applies.

Translated by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication)

We are certain that all parties will benefit from this agreement and wish you the best in the use of this material. Thank you.

Special Terms:

v1.1

If you would like to pay for this license now, please remit this license along with your payment made payable to "COPYRIGHT CLEARANCE CENTER" otherwise you will be invoiced within 48 hours of the license date. Payment should be in the form of a check or money order referencing your account number and this invoice number RLNK501114947.

Once you receive your invoice for this order, you may pay your invoice by credit card. Please follow instructions provided at that time.

**Make Payment To:
Copyright Clearance Center
Dept 001
P.O. Box 843006
Boston, MA 02284-3006**

For suggestions or comments regarding this order, contact RightsLink Customer Support: customercare@copyright.com or +1-877-622-5543 (toll free in the US) or +1-978-646-2777.

Gratis licenses (referencing \$0 in the Total field) are free. Please retain this printable license for your reference. No payment is required.

APPENDIX B



Ntombi Mudenda
PhD Candidate
Department of Pathobiological Sciences
School of Veterinary Medicine
Louisiana State University
Skip Bertman Drive
Baton Rouge, LA 70802
USA

Dear Dr. Mudenda, I am writing to confirm our authorization from Geospatial Health journal to use your publication titled “Ecological niche modelling of hookworm in Brazil based on climate” (Vol 6 No. 3, Sept 2012, pg 111-123) as part of your dissertation.

Best regards

Laura Rinaldi, *Associate Editor Geospatial Health*

A handwritten signature in black ink, appearing to read 'Laura Rinaldi', is located at the bottom right of the page.

APPENDIX C



CENTER FOR CLINICAL RESEARCH
AND TECHNOLOGY
11100 Euclid Avenue, Lakeside 1400
Cleveland, Ohio 44106

Exemption approval for IRB# EM-13-39

October 14, 2013

Ronald E Blanton,

Upon administrative review of your project entitled "Hookworm and Gastrointestinal Nematodes Among Patients in Mutuipe Municipality in Brazil" (IRB # EM-13-39), the IRB has determined that this protocol meets the criteria for exemption under Federal regulation 45 CFR 46.101 (b) (4).

Please note the expiration date for this exemption approval is **10/14/2019**.

Continuing review is not required for research for your exempt project. However, if the research is ongoing for more than six years, you must resubmit the project for re-review as a new submission.

Please see the UHCMC IRB policy regarding Exempt Research for more information:

http://www.uhhospitals.org/Portals/6/docs/research/irb/forms/May_07/IRB_Policy_Exempt_Human_Research_5_2007.doc

Please feel free to contact the Office of Institutional Review at (216) 844-1529 if you have questions.

Thank You,

A handwritten signature in black ink, appearing to read 'Ronald E Blanton'.

UHCMC IRB Chairperson
(Signature was applied by the IRB Administration Office)

*The UHCMC IRB operates under the HHS Federal Wide Assurance of Compliance number 00003937 and
IRB registration numbers 00000694 and 00001691*

p. 1 of 1 for Ronald E Blanton iRIS Reference # 029135

THE VITA

Ntombi B. Mudenda was born and raised in Zambia. She graduated from the University of Zambia, School of Veterinary Medicine with a Bachelor of Veterinary Medicine. She worked for the School of Veterinary Medicine as a house surgeon before pursuing a Master of Science degree in wild animal health at the Royal Veterinary College in London, UK under the sponsorship of the Beit Trust. Ntombi then returned to Zambia and lectured at the Veterinary School. She became a Fulbright foreign exchange student and joined the doctoral program at Louisiana State University, School of Veterinary Medicine in 2010. Ntombi is anticipating graduation with a Doctor of Philosophy in parasitology in December 2013.