

5-3-2007

Pair correlations in nuclei involved in neutrinoless double β decay: Ge76 and Se76

S. J. Freeman
The University of Manchester

J. P. Schiffer
Argonne National Laboratory

A. C.C. Villari
IN2P3 Institut National de Physique Nucleaire et de Physique des Particules

J. A. Clark
Yale University

C. Deibel
Yale University

See next page for additional authors

Follow this and additional works at: https://digitalcommons.lsu.edu/physics_astronomy_pubs

Recommended Citation

Freeman, S., Schiffer, J., Villari, A., Clark, J., Deibel, C., Gros, S., Heinz, A., Hirata, D., Jiang, C., Kay, B., Parikh, A., Parker, P., Qian, J., Rehm, K., Tang, X., Werner, V., & Wrede, C. (2007). Pair correlations in nuclei involved in neutrinoless double β decay: Ge76 and Se76. *Physical Review C - Nuclear Physics*, 75 (5) <https://doi.org/10.1103/PhysRevC.75.051301>

This Article is brought to you for free and open access by the Department of Physics & Astronomy at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Authors

S. J. Freeman, J. P. Schiffer, A. C.C. Villari, J. A. Clark, C. Deibel, S. Gros, A. Heinz, D. Hirata, C. L. Jiang, B. P. Kay, A. Parikh, P. D. Parker, J. Qian, K. E. Rehm, X. D. Tang, V. Werner, and C. Wrede

Pair Correlations in Nuclei Involved in Neutrinoless Double Beta Decay: ^{76}Ge and ^{76}Se .

S.J. Freeman,¹ J.P. Schiffer,^{2,*} A.C.C. Villari,³ J.A. Clark,⁴ C. Deibel,⁴
S. Gros,² A. Heinz,⁴ D. Hirata,^{3,5} C.L. Jiang,² B.P. Kay,¹ A. Parikh,⁴
P.D. Parker,⁴ J. Qian,⁴ K.E. Rehm,² X.D. Tang,² V. Werner,⁴ and C. Wrede⁴

¹*Schuster Laboratory, the University of Manchester, Manchester M13 9PL, UK*

²*Argonne National Laboratory, Argonne, Illinois 60439, USA*

³*GANIL (IN2P3/CNRS - DSM/CEA),*

B.P. 55027 14076 Caen Cedex 5, France

⁴*A.W. Wright Nuclear Structure Laboratory,*

Yale University, New Haven, CT 06520, USA

⁵*The Open University, Dept. of Physics and Astronomy, Milton Keynes, MK7 6AA, UK*

(Dated: September 17, 2021)

Abstract

Precision measurements were carried out to test the similarities between the ground states of ^{76}Ge and ^{76}Se . The extent to which these two nuclei can be characterized as consisting of correlated pairs of neutrons in a BCS-like ground state was studied. The pair removal (p,t) reaction was measured at the far forward angle of 3° . The relative cross sections are consistent (at the 5% level) with the description of these nuclei in terms of a correlated pairing state outside the N=28 closed shells with no pairing vibrations. Data were also obtained for ^{74}Ge and ^{78}Se .

PACS numbers: 25.40.-Hs, 27.50.+e, 23.40.-Hc

*Electronic address: schiffer@anl.gov

I. INTRODUCTION

Interest in the possibility of observing neutrinoless double beta decay ($0\nu 2\beta$) is considerable. If this decay were to be definitively observed, it would show that the neutrinos are their own antiparticles. In addition, the rate of the decay would be a measure of the neutrino rest mass, if the nuclear matrix element were known. Unfortunately, theoretical calculations of this do not agree well with each other [1]. It seems appropriate to determine additional properties of the ground states of the possible $0\nu 2\beta$ systems by experiment, and thus help constrain and test theoretical calculations of this exotic decay mode. One of the likely candidate nuclei is ^{76}Ge decaying to the ground state of ^{76}Se . We have started with a study of the properties of the ground states of these nuclei, and especially the similarities and differences between them, using transfer reactions. One part of this study is an accurate measurement of one-nucleon transfer in order to probe the occupation numbers of valence orbits for both neutrons and protons, with particular attention to changes in these occupations. The other part is to study pair-correlations in these nuclei by nucleon pair transfer. Here we report on a comparison of neutron pair transfer from the (p,t) reaction. We hope to obtain similar data on proton pair correlations from ($^3\text{He},n$) reactions in a future experiment.

Pair transfer between 0^+ states in the (p,t) or (t,p) reaction proceeds via $L=0$ transfer and the angular distribution, for energies above the Coulomb barrier, is sharply forward peaked. This feature was recognized early [2] and was crucial in exploring the importance of such correlations and their excitations, the so-called pairing vibrations [3]. The latter are an indication of deviations from the simplest pairing picture and can occur in regions of changing shapes, or when there is a gap in single-particle states, such as near a shell closure.

We have carried out measurements of the neutron pair-removal (p,t) reaction on targets of $^{74,76}\text{Ge}$ and $^{76,78}\text{Se}$. The reaction on ^{78}Se was measured because the $^{78}\text{Se}(p,t)^{76}\text{Se}$ leads to ^{76}Se , while $^{76}\text{Se}(p,t)^{74}\text{Se}$ starts from the ground state of ^{76}Se . Thus both reactions are relevant to the pairing structure of this ground state. The ^{74}Ge target was included as a check.

There are two relevant aspects to these measurements. The first is the matter of pairing vibrations. The even Ge and Se isotopes are well studied and evidence for excited 0^+ states has been established [4, 5, 6]. In some of the lighter isotopes of Ge and Se, two-neutron transfer reactions have shown significant strength populating excited 0^+ states.

These pairing vibrations indicate that there are significant BCS-like pair correlations in connecting the target ground state in an even initial nucleus to excited 0^+ states in the final. If there were significant differences in this regard between reactions leading to or from ^{76}Ge and ^{76}Se , this would be an indication that the pair correlations in the initial and final states in double-beta decay differ. Any reliable calculations of the process would presumably have to reproduce such differences in order to obtain a reasonable matrix element for the $0\nu 2\beta$ decay. Secondly, if accurate cross sections were available for pair transfer further checks could be made of the similarity between the initial and final wave functions used in such calculations.

II. EXPERIMENTAL CONSIDERATIONS

Given that the $L=0$ (p,t) transitions are the strongest in the spectrum of final states at very forward angles, which is also the region where the approximations inherent in the distorted wave Born approximation (DWBA) are best satisfied, it is desirable to carry out these measurements as close to 0° as feasible. The (p,t) reaction had been studied previously on Ge [4] and Se [6] isotopes; relevant (t,p) measurements have also been made [7, 8, 9]. Both the experimental methodology and the data analysis methods in the two (p,t) studies were different (the target thickness was estimated from high-energy elastic scattering where optical model predictions are ambiguous, the angular distributions start at 7.5 degrees, and only angle-integrated cross sections are quoted, etc.), making a reliable systematic comparison of Ge and Se data difficult. Our purpose here was to measure the cross sections at as far forward angles as feasible and to obtain a consistent set of accurate cross sections with particular care taken to reduce relative systematic uncertainties.

The choice of energy was governed by the desirability of having both protons and tritons well above the Coulomb barrier. For the present measurement we therefore chose 23-MeV protons from the Yale ESTU tandem Van de Graaf accelerator. This energy is similar to that used in earlier experiments, both for (p,t) and (t,p) reactions. Because one of the objectives of the present measurement was to obtain accurate cross sections, we chose to measure the thickness of the evaporated germanium and selenium targets in situ by simply lowering the proton beam energy to 6 MeV, where the elastic scattering cross sections are very close to Rutherford values. The angle should not be so far forward that small

uncertainties in angle would become significant and 30° was chosen because calculations with several optical potentials showed that the deviation from Rutherford scattering was less than 2%. The $^{76,74}\text{Ge}$ and $^{78,76}\text{Se}$ target thicknesses were found to range between 160 and $400 \mu\text{g}/\text{cm}^2$. Since Se can sublime at a relatively low temperature, this low-energy target thickness measurement was made at the beginning and then at the end of the experiment; no significant differences ($<3\%$) were observed. The highest beam currents used were about 35 nA, though considerably lower (2-3 nA) for the 3° measurements.

The Yale Enge split-pole spectrograph was used for the measurements with a focal-plane detector that cleanly separates tritons from other reaction products. As monitors, two Si surface barrier detectors at $\pm 32^\circ$ were used. They were calibrated in terms of beam intensity using a current integrator connected to a Faraday cup. The beam integrator was set to the same scale that was used for the low-energy measurements and the solid-angle setting for the aperture of the spectrograph was also the same, thus establishing a relationship between an absolute cross section scale in terms of the monitor yields, instead of the beam integrator, for each target.

The 3° setting for the spectrograph required that the Faraday cup be retracted so that the beam-current measurement had to rely on the previously calibrated monitor counters. Removal of the Faraday cup meant that the beam entered the spectrometer. The magnetic rigidity of the tritons from the reaction is such that, with the magnetic field set for observing tritons, protons from the target cannot directly reach the focal plane and are intercepted inside the spectrometer. Protons scattered at this point can enter the focal plane and, whilst they can be distinguished from tritons by their ionization density, they do impose a counting rate limit. As a result, the furthest forward angle where measurements could be made was 3° . Distorted-wave calculations indicate that the cross section at this angle is lower than that at 0° by about 8%. Spectra were also measured at the laboratory angle of 22° , which is close to the minimum for $L=0$ angular distributions, though the location and depth of the minimum is very sensitive to the Q-value and the distorting parameters as is seen in Figure 1. Nevertheless, the ratio of the 3° to the 22° cross sections is huge compared to that for the other L values, and is therefore an excellent identifier of $L=0$ transitions, though the precise value of this ratio will depend on the exact location and depth of the sharp minimum in the angular distribution.

Representative spectra from the 3° measurements are shown in Figure 2 where the

ground-state transitions are clearly seen to dominate. The results of the cross-section measurements are shown in Table I below, with ratios to the ground-state cross sections given only for states with yields, at 3° , larger than 1% of the ground state yield. The transitions where the ratio of cross sections between 3° and 22° is consistent with $L=0$ are shown in bold. More complete data, though at further back angles and with less attention to accurate relative cross sections had been reported, but the emphasis in these previous measurements was on angular distributions starting at 7.5° , and only angle-integrated cross sections are quoted [4, 5]. The uncertainty in the present experimental cross sections is believed to be $\pm 10\%$, while that in the relative values is estimated at $\pm 5\%$. These uncertainties are dominated by estimates of systematic errors (constancy of the beam spot on target, accuracy of angle determinations in the monitors, possible small drifts in monitor calibration, possible inefficiency in the focal plane detector, uniformity of target thickness etc.) while the statistical contribution is of the order of 1%.

The (p,t) differential cross sections at 3° for populating the 0^+ ground states for the four targets are very similar: 6.4, 6.7, 6.0, and 7.1 mb/sr for $^{74,76}\text{Ge}$ and $^{76,78}\text{Se}$ respectively. Excited 0^+ states stand out in the ratio between the 3° and 22° yields, which is an order of magnitude larger than for any other excited state. With the exception of the ^{74}Ge target, none of these excited 0^+ states is populated with a cross section at 3° that is more than 2% of that leading to the ground states. In the $^{74}\text{Ge}(p,t)^{72}\text{Ge}$ reaction the cross section to the first-excited 0^+ state is 1.9 mb/sr. This feature is well known [4] as an example of a pairing vibration. The case of ^{74}Ge is illustrative of effects that can be problematic; however the context of the current work is related only to the $^{76}\text{Ge}/^{76}\text{Se}$ double-beta-decay system.

III. DWBA CALCULATIONS

The calculations were carried out with the DWBA program PTOLEMY [10] to correct the dependence of the reaction on Q-values. The consideration of the details of nuclear structure is beyond the scope of this study, even though ^{76}Ge and ^{78}Se have 6 neutron vacancies in the $N=50$ shell, ^{74}Ge and ^{76}Se have 8. The form factor for the neutron pair was calculated assuming a mass-2, $\ell = 0$ di-neutron bound in a Woods-Saxon potential with the appropriate binding energy and having 3 nodes in its wave function. The proton potentials were those of Ref. [11] and the triton potential that of Ref. [12]. The measured cross sections

at 3 degrees are given in Table II together with the ratio of the experimental cross sections to the calculated values. The absolute magnitude of the DWBA cross section is very sensitive to the choice of the distorting potentials as is the location of the first minimum in the angular distribution. However, all calculations indicate that the relative values between these targets of the calculated cross sections at 3° remain the same (within 10%) with various reasonable potentials for protons and tritons.

With the proton potential of Ref. [11], the average ratio changes from 136 to 217. The experimental cross sections by themselves are remarkably constant; without any correction for reaction dynamics they vary by $\pm 6\%$. Dividing the experimental numbers by the calculated reaction cross sections, reduced the difference between ^{76}Ge and ^{76}Se even further. However, this improvement is probably not significant, in view of the estimated 5% relative error and the neglect of the different numbers of neutrons. The peak cross sections and DWBA trends are also shown in Figure 3.

IV. PAIR-ADDING REACTIONS

The pair-adding (t,p) reaction had been studied previously, for both the $^{74}\text{Ge}(t,p)^{76}\text{Ge}$ [7, 8] and the $^{76}\text{Se}(t,p)^{78}\text{Se}$ [9] reactions. While the ground-state transitions are the same as the ones studied here, transitions to excited states could, in principle, show up and indicate pairing vibrations. While the angular distributions were not measured at as far forward angles as in the present work, some estimate can be obtained by comparing integral cross sections. No excited 0^+ states were seen in these (t,p) studies with a strength greater than 4% of the ground-state transition, confirming the dominance of pair correlations in the ground states of these nuclei, with no splitting into pairing vibrations.

V. CONCLUSIONS

The experimental cross sections measured in this experiment are remarkably constant for ground-state transitions with the various targets. The difference between ^{76}Ge and ^{76}Se , in the ratio of the experimental ground-state transition strengths divided by the appropriate DWBA calculations, is less than the 5% estimated accuracy of the measurements. Transitions to excited 0^+ states from ^{76}Ge and ^{76}Se targets are no more than a few percent of

the ground-state transitions indicating no sign of the pairing vibrations that appear in some of the lighter isotopes. Had such admixtures been present, this would have complicated a simple comparison of the ground states. The constancy of the ground-state strength in pair correlations seems to be as true for neutron-pair-adding transfers leading to these nuclei as it is for pair removal from them. The present results suggest that the ground states of ^{76}Ge and ^{76}Se exhibit quantitatively very similar neutron pair-correlations. Changes in pairing are thus unlikely to be a significant complicating factor in the wave functions of these states for calculations of neutrinoless double β decay.

We wish to acknowledge helpful discussions with S.C. Pieper, Ben Bayman, and M.H. Macfarlane. The help of John Greene in meticulous work on target preparation is also acknowledged. The work was supported by the U.S. Department of Energy, Office of Nuclear Physics, under contracts DE-FG02-91ER-40609 and DE-AC02-06CH11357, the UK Engineering and Physical Sciences Research Council, and IN2P3/CNRS-France.

-
- [1] J.N. Bahcall, H. Murayama, and C. Peña-Garay, *Phys. Rev. D* **70**, 033012 (2004).
 - [2] S. Yoshida, *Nucl. Phys.* **33**, 685 (1962).
 - [3] D.R. Bes and R.A. Broglia, *Nucl. Phys* **80**, 289 (1966).
 - [4] D. Guilbault et al., *Phys. Rev. C* **16**, 1840 (1977).
 - [5] D. Ardouin et al., *Phys. Rev. C* **18**, 1201 (1978).
 - [6] M. Borsaru et al., *Nucl. Phys.* **284**, 379 (1977).
 - [7] S. Mordechai et al., *Phys Rev. C* **18**, 2498 (1978).
 - [8] C. Lebrun et al., *Phys. Rev. C* **19**, 1224 (1979).
 - [9] D.L. Watson and H.T. Fortune, *Phys. Rev. C* **35**, 430 (1987).
 - [10] M.H. Macfarlane and Steven C. Pieper, ANL-76-11 Rev. 1 Argonne National Laboratory Report (1978) unpublished.
 - [11] F.D. Becchetti and G.W. Greenlees, *Phys. Rev.* **182**, 1190 (1969).
 - [12] R. Perry, *Phys. Rev. C* **24**, 1471 (1981).
 - [13] F.G. Perey, *Phys. Rev.* **131**, 745 (1963).




FIG. 1: (Color online) DWBA calculations of ground-state angular distributions at 23 MeV for different targets, using the proton optical potentials of Ref. [12] and the triton potential from Ref. [7]. The intent of the figure is to illustrate the sensitivity of the shapes of the angular distributions to the Q-values, while the peak cross sections remain relatively stable. The relative variation in the peak cross sections for different choices of potentials is very similar.




FIG. 2: (Color online) Spectra of tritons at 3° measured with the Yale split-pole spectrograph, normalized to 100 for the ground-state peak, and labeled in each case by the target nuclide. The peaks corresponding to $L=0$ transitions are identified by a pointer. Peaks due to isotopic impurities are marked by an x. While there is evidence in $^{74}\text{Ge}(p,t)^{72}\text{Ge}$ of substantial strength in a low-lying excited 0^+ state, there are no large admixtures seen for the ^{76}Ge and ^{76}Se targets.




FIG. 3: (Color online) The ground-state 0^+ to 0^+ cross sections at 3° are plotted as a function of Q-value, for convenience in display. Also shown are the DWBA cross sections multiplied by one average normalization factor for each proton potential. The estimated relative errors are shown on the experimental points.

TABLE I: Summary of (p,t) cross sections at 3° and the ratio (in %) of these to the 22° values. Transitions consistent with L=0 are shown in boldface.

$^{74}\text{Ge}(p,t)^{72}\text{Ge}$		
$\sigma_{\text{gs}}(\text{lab}) = 6.4 \text{ mb/sr}$		
Excitation Energy (keV)	$(\sigma/\sigma_{\text{gs}})_{3^\circ}$	Ratio(3°/22°)
0	100	86
691	29	280
834	2.8	0.9
1464	0.5	1.5
2024	0.5	4
2762	0.9	130

$^{76}\text{Ge}(p,t)^{74}\text{Ge}$		
$\sigma_{\text{gs}}(\text{lab}) = 6.7 \text{ mb/sr}$		
Excitation Energy (keV)	$(\sigma/\sigma_{\text{gs}})_{3^\circ}$	Ratio(3°/22°)
0	100	50
596	3.2	1.0
1204	1.1	1.6
1463	2.2	0.8
2198	2.9	3
2833	1.7	6

$^{76}\text{Se}(p,t)^{74}\text{Se}$		
$\sigma_{\text{gs}}(\text{lab}) = 6.0 \text{ mb/sr}$		
Excitation Energy (keV)	$(\sigma/\sigma_{\text{gs}})_{3^\circ}$	Ratio(3°/22°)
0	100	115
635	1.0	0.4
854	1.4	80

$^{78}\text{Se}(p,t)^{76}\text{Se}$		
$\sigma_{\text{gs}}(\text{lab}) = 7.1 \text{ mb/sr}$		
Excitation Energy (keV)	$(\sigma/\sigma_{\text{gs}})_{3^\circ}$	Ratio(3°/22°)
0	100	150
559	1.2	0.4
1121	0.8	4
1220	0.7	1.0

TABLE II: 3° laboratory cross sections and ratios to DWBA. Cross sections are for the ground-state to ground-state transitions.

Target	$\sigma_{\text{exp}}(lab)$ (mb/sr)	σ_{DWBA} (mb/sr)	$\sigma_{\text{exp}}/\sigma_{\text{DWBA}}$
^{74}Ge	6.4	0.0438	147
^{76}Ge	6.7	0.0499	135
^{76}Se	6.0	0.0437	137
^{78}Se	7.1	0.0431	164