

5-1-2014

APeg3: Regulation of Peg3 through an evolutionarily conserved ncRNA

Wesley D. Frey
Louisiana State University

Joomyeong Kim
Louisiana State University

Follow this and additional works at: https://digitalcommons.lsu.edu/biosci_pubs

Recommended Citation

Frey, W., & Kim, J. (2014). APeg3: Regulation of Peg3 through an evolutionarily conserved ncRNA. *Gene*, 540 (2), 251-257. <https://doi.org/10.1016/j.gene.2014.02.056>

This Article is brought to you for free and open access by the Department of Biological Sciences at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Published in final edited form as:

Gene. 2014 May 1; 540(2): 251–257. doi:10.1016/j.gene.2014.02.056.

APeg3: regulation of *Peg3* through an evolutionarily conserved ncRNA

Wesley D. Frey and Joomyeong Kim

Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, USA

Joomyeong Kim: jkim@lsu.edu

Abstract

Mammalian *APeg3* is an antisense gene that is localized within the 3'-untranslated region of the imprinted gene, *Peg3*. *APeg3* is expressed only in the vasopressinergic neurons of the hypothalamus, thus is predicted to play significant roles in this specific area of the brain. In the current study, we investigate the functions of *APeg3* with comparative genomics and cell line-based functional approaches. The transcribed region of *APeg3* displays high levels of sequence conservation among placental mammals, but without any obvious open reading frame, suggesting that *APeg3* may have been selected as a ncRNA gene during eutherian evolution. This has been further supported by the detection of a conserved local RNA secondary structure within *APeg3*. RNA secondary structure analyses indicate a single conserved hairpin-loop structure towards the 5' end of the transcript. The results from cell line-based transfection experiments demonstrate that *APeg3* has the potential to down-regulate the transcription and protein levels of *Peg3*. The observed down-regulation by *APeg3* is also somewhat orientation-independent. Overall, these results suggest that *APeg3* has evolved as a ncRNA gene and controls the function of its sense gene *Peg3* within specific neuronal cells.

Keywords

Peg3; *APeg3*; ncRNA; Vasopressin neuron; Genomic imprinting

1. Introduction

APeg3 is an antisense gene that is located within the 3'-untranslated region of an imprinted gene, *Peg3* (paternally expressed gene 3). The expression of *APeg3* is detected only in vasopressinergic neurons of the hypothalamus, suggesting a very specialized role in the neuronal cells controlling blood pressure and the volume of bodily fluid in mammals (Glasgow et al., 2005). In fact, *APeg3* was initially identified as a gene that is highly up-regulated in response to osmotic challenges in rat brains (Glasgow et al., 2005). Studies on *Peg3* also demonstrate up-regulation against osmotic stress in this cell type, suggesting that

© 2014 Elsevier B.V. All rights reserved.

Correspondence to: Joomyeong Kim, jkim@lsu.edu.

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.gene.2014.02.056>.

Conflict of interest: The authors declare no conflict of interest.

both *APeg3* and *Peg3* may play important roles in the functions of vasopressin-expressing neurons (Yamashita et al., 2002). Earlier studies from rat brain suggest that *APeg3* might code for a small-sized Open Reading Frame (ORF), but were not substantiated by later studies as the observed ORF is not conserved in other mammals, such as humans and mice. Interestingly, *APeg3* is also maternally imprinted as seen in *Peg3*: only the paternal allele is expressed while the maternal allele is repressed (Choo et al., 2008). Nevertheless, the functional impetus for *APeg3* imprinting is currently not well understood.

APeg3 is one of several ncRNA genes that have been identified from mammalian imprinted domains. The list of the imprinted ncRNA genes includes *H19*, *IPW*, *Kcnq1ot1*, *Nespas*, *Airn*, *Copg2*, and *Gtl2* (Koerner et al., 2009). *APeg3* is very unique compared to the other imprinted ncRNA genes in the following aspects. First and foremost, *APeg3* is the only imprinted antisense transcript expressed from the same (paternal) allele as its sense counterpart. Similar antisense ncRNA counterparts to imprinted genes, such as *anti-Rtl1* and *Copg2-AS*, are selectively expressed from the opposite allele as the sense genes (Ferguson-Smith, 2011; Yamasaki et al., 2000). Secondly, the size of *APeg3* is relatively small, 1.5 kb in length, whereas the size of the other imprinted non-coding genes range up to several hundreds of kb in length (Glasgow et al., 2005; Koerner et al., 2009). *H19* is the only other imprinted ncRNA of similar length at ~1.9 kb (Brannan et al., 1990). Thirdly, *APeg3* lies antisense to the protein-coding gene, *Peg3*. Furthermore, the transcribed region overlaps with the 3'UTR of *Peg3*. This is quite different from other imprinted ncRNA genes, such as *Gtl2*, which tend to overlap with the entire imprinted region. Finally, the transcript of *APeg3* is detected as an intronless mRNA with poly-A tails in vivo, suggesting that *APeg3*'s mRNA goes through normal processes as Pol II transcripts (Choo et al., 2008). By contrast, other long ncRNA genes, such as *Airn*, *Kcnqot1*, and *Rian* reside inside the nucleus and do not go through normal processes as Pol II transcripts. Despite these unique features, the biological roles of *APeg3* are currently unknown. However, given the antisense/sense relationship between *APeg3* and *Peg3*, it is reasonable to predict that the main function of *APeg3* may be controlling the transcription and protein levels of *Peg3*. To investigate this possibility, we performed a series of comparative genomics and cell-line based functional assays in the current study. The results suggest that *APeg3* may have evolved as a ncRNA gene controlling *Peg3* mRNA and protein levels.

2. Materials and methods

2.1. Database search and gene prediction

The intronless human antisense transcript, *PEG3-ASI* (NR_023847.2), was used as a reference sequence in the BLAST program offered through the UCSC genome browser (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Using this sequence, homologous nucleotide sequences were obtained from the available genome sequences of multiple mammalian species. After our initial inspection of the retrieved sequences, one representative sequence from each order of placental mammals was used for sequence alignment with the ClustalW multiple sequence alignment (<http://www.genome.jp/tools/clustalw/>) as well as CLC Bio Workbench. The 1.5-kb genomic region for each species' *APeg3* gene is as follows: *Mus musculus* for Rodentia (GRCm38/mm10 chr7: 6,706,295–6,707,624), *Homo sapiens* for

Primates (GRCh37/hg19 chr19: 57,323,893–57,325,161), *Equus caballus* for Perissodactyla (Broad/EquCab2 chr10: 25,780,355–25,781,617), *Oryctolagus cuniculus* for Lagomorpha (Broad/oryCun2 chrUn0113: 622,308–623,587), *Loxodonta africana* for Proboscidea (Broad/loxAfr3 scaffold_4: 18,505,516–18,506,810), *Tursiops truncatus* for Cetacea (Baylor Ttru_1.4/turTru2 JH478484: 13,610–14,893), *Myotis lucifugus* for Chiroptera (Broad Institute Myoluc2.0/myoluc2 GL430552: 92,812–94,034), *Dasypus novemcinctus* for Cingulata (Baylor/dasNov3 JH562679: 85,036–86,329) and *Trichechus manatus* for Sirenia (Broad v1.0/triMan1 JH594782: 4,734,304–4,735,606). These sequences were further analyzed to identify several sequence features for the transcribed region of *APeg3*. Transcription Start Site (TSS) and Poly-A site were identified based on switchgear analyses. A potential TATA box was identified 25-bp upstream of the TSS site as an evolutionarily conserved element. A target site for miR-124 was also identified based on TargetScan (<http://www.targetscan.org/>) data. The results of these analyses are readily available on the UCSC genome browser. All the sequences used for the current study are available (Supplementary materials 2–3).

2.2. Secondary structure prediction of *APeg3*

The primary sequence alignment of *APeg3* was used as an input for covariant and minimal fold energy (MFE) analysis using Vienna Suite software (Washietl et al., 2005). RNA folding was predicted using the following two programs: RNA AliFold (<http://rna.tbi.univie.ac.at/cgi-bin/RNAalifold.cgi>) to predict the macro-secondary structures (global folds) and RNAz (<http://www.tbi.univie.ac.at/~wash/RNAz/>) to predict the micro-secondary structures (local folds) of *APeg3* (Washietl et al., 2005; Gruber et al., 2007). The programs used the primary sequence alignments to generate a consensus fold structure for *APeg3*. This series of analyses used two controls. The sequence of the H19 gene was included as a positive control, which has been previously shown to have significantly conserved secondary structures important for its function as a lncRNA (Juan et al., 2000), and a shuffled sequence of *APeg3* was also included as a negative control.

2.3. Construction and transfection of *APeg3* overexpression vectors

To construct the Short- and Full-length *APeg3* expression vectors, the cDNA prepared from adult mouse brains was amplified with artificial BamHI and NotI sites using the following primers:

APeg3 Full-length-APeg3Full_BamHI CGCGGATCCGGGAATTAA
 GTCTGGAGACACAAAGATCTAAGG and APeg3Short_NotI ATAA
 GAATGCGGCCGCGCACCAGTGCAGGTGGTGGCGGA. APeg3 Short-
 APeg3Short_BamHI CGCGGATCCCAATCAGTCTCAAGGGTCTGGGT and
 APeg3Short_NotI ATAAGAATGCGGCCGCGCACCAGTGCAGGTGG
 TGCGGA.

(-)APeg3 Full-length-(-)APeg3Short_BamHI CGCGGATCCGCACC
 AGTGCAGGTGGTGGCGGA and (-)APeg3Full_NotI
 ATAAGAATGCGGCCGCGGAATTAAGTCTGGAGACACAAAGATCTAAG
 G (-)APeg3 Short-length-(-)APeg3Short_NotI

ATAAGAATGCGGCCGCAATCA GTCTCAAGGGGTCTGGGT
 (–)APeg3Short_BamHI CGCGGATCCGCA CCAGTGCAGGTGGTGC GGA

The amplified products were digested with BamHI and *NotI* and cloned into pCDNA3.1Hygro(–) vector (Invitrogen). In this vector system, the expression of *APeg3* was driven by the CMV promoter and terminated by the Poly-A signal of BGH. The constructed vectors were transfected into Neuro2A cells with each well containing 0.15×10^6 cells using 4 μ L lipofectamine 2000 (Invitrogen). Transfections were carried out in triplicate with 3 wells designated for subsequent RNA isolation and 3 wells for protein extraction. The transfected cells were selected using 500 μ g/mL hygromycin B 24 h post-transfection. The cells were harvested for RNA isolation using TRIzol (Invitrogen) 48 h post-transfection, and also for protein extraction in lysis buffer (0.25 M Tris–HCl, pH 7.8, plus 0.1% NP-40) supplemented with fresh β -mercaptoethanol and the proteinase inhibitor cocktail (Calbiotech).

2.4. cDNA synthesis and quantitative Reverse-Transcription PCR (qRT-PCR)

Total RNA was isolated from transfected Neuro2A cells using a commercial kit (Trizol, Invitrogen). The isolated RNA was first reverse-transcribed using the MLV First-Strand Synthesis System (Invitrogen) with random and oligo dT primers. Subsequent cDNA was used as a template for qRT-PCR. This analysis was performed with the iQ SYBR green supermix (Bio-Rad) using the iCycler iQTM multicolor real-time detection system (Bio-Rad). All qRT-PCR reactions were carried out for 40 cycles under standard PCR conditions. The expression levels of *Peg3* were first normalized to those of β -Actin, then compared to those of the empty vector (pCDNA) controls. The primers for *Peg3* and β -Actin were as follows: *Peg3*-1a (5′-GGTTCAGTGTGGGTGCACTAGA CT-3′), *Peg3*-1b.1 (5′-GCTCACACCCAAGGGCTTGAGCGT-3′), *Peg3*-1b.3 (5′-TCCCTAGTGTGCATGATCTGGT-3′), *bactin*-1a (5′-GAGCACCCCTGTGCTGCTCACCGA-3′) and *bactin*-1b (5′-CTCTTTGATGTCACGCACGATTTTC-3′).

2.5. Protein isolations and Western blots

The transfected Neuro2A cells were first rinsed with 1 \times PBS, and lysed with the lysis buffer (0.25 M Tris–HCl, pH 7.8, plus 0.1% NP-40). The subsequent crude lysates were collected and snap-frozen in liquid nitrogen. Later, the lysate was then thawed at 4 °C to prevent protein degradation. The cellular debris was removed by centrifugation for 10 min at 4 °C. Protein concentrations were determined by the Bradford assay kit (Pierce), using diluted BSA as protein standards. Sixty micrograms of each lysate was separated on 10% SDS-PAGE gels and transferred to PVDF membranes (Hybond-P, Amersham) using a Mini Trans-Blot wet transfer cell (Bio-Rad). Membranes were blocked for 1 h in the Tris-buffered saline containing 1% skim milk and 0.05% Tween100, and incubated overnight at 4 °C with the custom-made anti-PEG3 antibody (Kim et al., 2012). These blots were incubated for an additional 1 h with the secondary antibody linked to horseradish peroxidase (Sigma). The blots were developed using the Western blot detection system according to the manufacturer's protocol (Intron Biotech). Densitometry was measured using Jmol: an open-source Java viewer for chemical structures in 3D.

3. Results

3.1. APeg3 as an evolutionarily conserved ncRNA in placental mammals

To analyze the evolutionary conservation of *APeg3*, human *PEG3-AS1* sequence (NR 023847.2) was used as a probe to search the UCSC genome browser, yielding genomic sequences for each of the 9 eutherian orders (Supplementary material 1). Using these sequences, pairwise sequence comparison analyses were conducted, and the results indicated high levels of sequence conservation between different mammals: 76 to 84% nucleotide sequence identity in the 1.5-kb transcribed region of *APeg3* (Table 1). Multiple sequence alignments were also performed using the 9 representative sequences, and the results were visualized as a graph in the bottom panel of Fig. 1. The UCSC PhyloP-derived placental mammal sequences used for the zoomed-out conservation include multiple species corresponding mainly to primate lineages as these are the most complete sequenced genomes and do not account for gaps in sequences. In this way, the conservation analysis could be biased. However, our custom zoomed-in conservation was derived from only one representative species of each order within the class eutheria, offering an unbiased characterization of the region. Differences in conservation level between the zoomed-out view and the zoomed-in view are the result of hand-picking genomic regions of *APeg3* to enhance the accuracy of the initial PhyloP conservation. High levels of sequence conservation are again observed throughout the 1.5-kb region with many small-size insertions/deletions specific to individual species, which can be seen as sudden drop-offs in the graph. Several sequence motifs, important for Pol II transcription, are also found within or around this 1.5-kb genomic region, including a TATA box, Transcription Start Site (TSS) and two Poly-Adenylation sites. However, there are no ORFs or Kozak sequences that are conserved among placental mammals. These results suggest that the 1.5-kb genomic region of *APeg3* is a gene template for a poly-adenylated non-coding RNA (ncRNA). The high levels of sequence conservation detected within the 1.5-kb region are also unusual, and can be contrasted with the 3'-UTR of an adjacent gene, *Zim1*, which shows no sequence conservation (Mammalian Cons Plot in Fig. 1). Overall, the high level of primary sequence conservation further supports that the region has not evolved simply as an UTR for the sense gene, *Peg3*. Instead, this region demonstrates signs of selection as a ncRNA gene during mammalian evolution.

3.2. Global and local secondary structure of APeg3 with evolutionary conservation

To determine potential functions of *APeg3* as a structural ncRNA, secondary structure predictions were performed using RNAalifold and RNAz programs. First, RNAalifold uses sequence alignments and determines folding free energies of relatively large RNA sequences (>200 bp) (Bernhart et al., 2008; Ge and Zhang, 2013). This program produces a consensus fold structure based on the lowest folding energy modeled and weights the conservation of folded structures among individual sequences. Both *APeg3* and the corresponding reverse complement (-)*APeg3* alignment were analyzed using RNAalifold to detect potentially large-scale secondary structures in both orientations. This series of analyses also used two controls: H19 as a positive control, an imprinted ncRNA previously seen to harbor functional characteristics of similar length to *APeg3*, and a shuffled *APeg3* alignment as a negative control. According to the outcomes of the prediction (Fig. 2), the sense and

antisense *APeg3* exhibit similar conservation levels of secondary structure with different overall structures.

Second, the RNAz program was used to detect any conserved micro-secondary structures within *APeg3*. The program analyzes multiple sequence alignments and recognizes conserved fold structures in small (200 bp) sliding-windows (Gruber et al., 2007). These folds are based on the lowest MFE of each short sequence. Covariant analysis was then performed on each fold and the resulting folds were scored by the program. In this program, a class probability (P) is used to measure the likelihood of a fold. The program uses a cutoff of $P = 0.9$, which can accurately predict 75% of known ncRNAs and classify folds with 99% specificity (Washietl et al., 2005). According to the outcomes, a single fold was predicted for *APeg3* in the 1.5-kb interval (Fig. 2C). The fold was conserved at the 5'-side of the transcript (nucleotide positions 145–185 in NR 023846.1) ($P = 0.99$). Interestingly, the antisense of *APeg3* also displayed one conserved stem-loop structure, located at the 3'-side of the transcript ($P = 0.91$) (Fig. 2D). Overall, these secondary structure fold predictions indicate the presence of a conserved hairpin loop structure in both sense and antisense orientations of *APeg3*. A series of predictions were also conducted using two controls, H19 and the shuffled *APeg3*. This series of analyses predicted a known small fold, miR675, within H19 ($P = 1.0$), and only one insignificant fold within the shuffled *APeg3* ($P = 0.76$); confirming the sensitivity of the RNAz program.

3.3. *APeg3* down-regulates *Peg3* mRNA

As seen in many antisense transcripts (Osato et al., 2007; Katayama et al., 2005), the potential function of *APeg3* may be to regulate the expression of the sense gene, *Peg3*. To determine whether *APeg3* has the potential to regulate *Peg3*, we performed a series of in vitro experiments as described below. We have constructed 4 expression vectors containing the different portions of mouse *APeg3*: *APeg3*-Full and -Short in both sense and antisense directions (the extent of *APeg3*-Short is indicated in Fig. 1). After transfections of these vectors to Neuro2A cells, we measured *Peg3* mRNA levels using qRT-PCR (Fig. 3). This series of expression analyses used an empty vector as a control (pCDNA). Overexpression of *APeg3*-Short resulted in down-regulation of *Peg3* by $57.60 \pm 0.4\%$ ($P = 0.0034$), whereas overexpression of (-) *APeg3*-Short resulted in a much smaller change in *Peg3* expression ($27.80 \pm 0.16\%$) ($P = 0.0536$). On the other hand, overexpression of *APeg3*-Full caused a much greater down-regulation of *Peg3* ($81.32 \pm 0.03\%$) ($P = 0.0008$), which was an ~20% lower expression than those from *APeg3*-Short. Interestingly, however, overexpression of (-) *APeg3*-Full also caused down-regulation of *Peg3* ($49.30 \pm 0.054\%$) ($P = 0.0271$). Given the results from both directions of *APeg3*-Full, the observed down-regulation by *APeg3*-Full may be orientation-independent. Furthermore, it is possible that the potential region causing this down-regulation might be located within the 5'-side of the 1.5-kb region of *APeg3* since this down-regulation is much more obvious in the Full-length constructs than in the Short-length constructs lacking the 5'-side of *APeg3*. To account for any minor transcriptional variants of *Peg3*, we also performed the previous qRT-PCR with primers amplifying the entire *Peg3* transcript (*Peg3*-1a, *Peg3*-1b.3) and observed a similar trend (Supplementary Fig. 2). *Peg3* mRNA levels were decreased in transfections of *APeg3*-Short ($71.43 \pm 0.056\%$) ($P = 0.0118$), *APeg3*-Full ($11.13 \pm 0.0635\%$) ($P = 0.0229$) and (-)*APeg3*-Full

($24.61 \pm 0.237\%$) ($P = 0.0332$), while changes in *Peg3* mRNA were not statistically significant in the (-) APeg3-Short ($54.18 \pm 0.425\%$) ($P = 0.2135$) transfections. However, these alternative forms of Peg3 have minimal contribution as compared to the major Peg3 transcript (Kim et al., 1997). Taken together, these results demonstrate that APeg3 has the potential to down-regulate the level of *Peg3* mRNA in vitro.

3.4. APeg3 down-regulates the protein levels of PEG3

To confirm whether the observed down-regulation of *Peg3* transcript resulted in reduced levels of the PEG3 protein, we performed Western blots with protein extracts prepared from the transfected Neuro2A cells (Fig. 4). The protein extracts from the two samples transfected with the sense and antisense APeg3-Full, respectively, were analyzed and compared with those from the sample transfected with the empty vector (pCDNA). The protein levels of PEG3 in each sample were first quantified using densitometric assays through the Jmol application <http://www.jmol.org>, and later normalized with those of β -Actin. The normalized values were finally compared with serial dilutions (100–40%) of the control sample transfected with pCDNA. According to the results, the protein levels of PEG3 were lowered by 85% in the transfectants with APeg3-Full, and by 74% in the transfectant with (-)APeg3-Full (lanes 2 and 3 in Fig. 4). The reduced protein levels observed from these transfections with APeg3-Full are consistent with the down-regulation observed in *Peg3* mRNA (Fig. 3). In conclusion, the results from both qRT-PCR and Western blotting demonstrate that the overexpression of APeg3 indeed results in reduction in the mRNA and protein levels of *Peg3*.

4. Discussion

In the current study, the unknown functions of APeg3 have been investigated using two different approaches, comparative genomic and cell line-based functional analyses. First, comparison of the sequences from different mammals revealed unusually high levels of conservation, but without any obvious ORF in the transcribed region of APeg3, suggesting that APeg3 may have been selected as a ncRNA gene during mammalian evolution (Fig. 1). This has been further supported by the detection of a conserved RNA secondary structure within APeg3 (Fig. 2). Second, the results from cell line-based analyses further demonstrated that APeg3 has the potential to down-regulate the transcription and protein levels of *Peg3* (Figs. 3 & 4). Thus, these results suggest that APeg3 has likely evolved as a ncRNA gene controlling the function of the sense gene, *Peg3*.

The high level of sequence conservation within the transcribed region of APeg3 is very unusual given the following reasoning (Fig. 1): The transcribed region of APeg3 does not harbor any obvious ORF and Kozak consensus motifs that are conserved among mammals, thus suggesting that APeg3 is a ncRNA gene. This is further supported by the presence of many insertions/deletions in the transcribed region of APeg3 between different mammals. At the same time, the sense direction of APeg3 is part of the 3'UTR of *Peg3*. In terms of functional selection during evolution, both ncRNA and 3'UTRs are expected to have much less constraints than the protein-coding region of individual genes (Li and Graur, 1991). As a result, these non-coding regions should have much less sequence conservation than the coding regions of genes. This can be easily observed in the 3'UTR of many genes, including

Zim1, which shows no conservation at all (Fig. 1). Nevertheless, the transcribed region of *APeg3* has maintained high levels of sequence identity during mammalian evolution (Table 1), which is quite comparable to the conservation levels of protein-coding regions. This might be related to the fact that this region harbors sequence elements for both genes, *APeg3* and *Peg3* with opposite transcriptional orientations. Given this special situation, this is the most likely explanation for the unusual sequence conservation observed from *APeg3*, which needs further investigation in the near future.

According to the results derived from in vitro transfection experiments, *APeg3* has the potential to down-regulate expression of *Peg3* (Figs. 3 & 4). The actual mechanism by which *APeg3* down-regulates *Peg3* is currently unknown, but the following scenario can be envisioned with the limited evidence presented in the current study. The most dramatic down-regulation by *APeg3* was observed with the full-length *APeg3* constructs, yet this down-regulation appears to be somewhat orientation-independent. The complementarity between *Peg3* and *APeg3* could allow for RNA:RNA interactions at the proposed hairpin-loop which could be causing transcriptional or translational interference of *Peg3* via *APeg3* regardless of *APeg3* orientation. However, in silico predictions do not correlate this interaction to be the most likely at the 5' hairpin-loop structure (Mueckstein et al., 2006).

Another possibility for the orientation-independent down-regulation of *Peg3* is through small RNA-mediated silencing. There are currently four known mechanisms for small RNA to down-regulate protein-coding mRNA, miRNA, snoRNA, pi-RNA and siRNA (Meister and Tuschl, 2004). Of these four, the miRNA-mediated mechanism is mainly known to be associated with imprinted domains, and functions independent of orientation (Royo and Cavaille, 2008; Meister and Tuschl, 2004). Thus, the orientation-independent down-regulation by *APeg3* might be mediated through some unknown miRNA embedded within the *APeg3* locus. In that regard, it is relevant to point out the fact that the 5'-side of *APeg3* contains a potential target site for miR-124 (Fig. 1), further supporting this possibility.

If this is the case, is this potential miRNA from *APeg3* mainly targeting the transcript from *Peg3*? Given the similar expression profiles in vasopressin-expressing neurons, we believe that *Peg3* is most likely the main target of *APeg3*. However, we cannot rule out the possibility that other genes in those specific neuronal cells could be targeted by a potential miRNA from *APeg3*. In conclusion, the observed down-regulation of *Peg3* by *APeg3* is very intriguing given the crucial roles played by *Peg3* in eutherians, thus it would be of great interest to further investigate this observed down-regulation by *APeg3* in terms of molecular mechanisms as well as potential targets.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We would like to thank Corey Bretz, Arundhati Bakshi and Dr. Hana Kim for their careful reading and discussion of the manuscript. We would like to thank Dr. David Donze and Asawari Korde for providing the facilities and guidance in northern blotting for confirmation of *APeg3* over-expression and Michelle Thiaville for her mentorship in the *APeg3* transfection experiments. We would also like to thank Rui Lu, Ryoichi Teruyama and James Moroney

for their intellectual input towards experimental designs. Finally, another acknowledgement should go out to HK for her help with data analysis and comparative genomics. We would also like to thank the reviewers for their insightful comments. This research was supported by the National Institutes of Health (J.K. R01-GM066225 and R01-GM097074).

References

- Bernhart SH, Hofacker IL, Will S, Gruber AR, Stadler PF. RNAalifold: improved consensus structure prediction for RNA alignments. *BMC Bioinformatics*. 2008; 9:474. [PubMed: 19014431]
- Brannan CI, Dees EC, Ingram RS, Tilghman SM. The product of the H19 gene may function as an RNA. *Molecular and Cellular Biology*. 1990; 10(1):28–36. [PubMed: 1688465]
- Choo JH, Kim JD, Kim J. Imprinting of an evolutionarily conserved antisense transcript gene APeg3. *Gene*. 2008; 409(1–2):28–33. [PubMed: 18166281]
- Ferguson-Smith, Anne C. Genomic imprinting: the emergence of an epigenetic paradigm. *Nature Reviews. Genetics*. 2011; 12(8):565–575. [PubMed: 21765458]
- Ge P, Zhang S. Incorporating phylogenetic-based covarying mutations into RNAalifold for RNA consensus structure prediction. *BMC Bioinformatics*. 2013; 14:142. [PubMed: 23621982]
- Glasgow E, Ryu SL, Yamashita M, Zhang BJ, Mutsuga N, Gainer H. APeg3, a novel paternally expressed gene 3 antisense RNA transcript specifically expressed in vasopressinergic magnocellular neurons in the rat supraoptic nucleus. *Molecular Brain Research*. 2005; 137(1–2):143–151. [PubMed: 15950772]
- Gruber AR, Neubock R, Hofacker IL, Washietl S. The RNaz web server: prediction of thermodynamically stable and evolutionarily conserved RNA structures. *Nucleic Acids Research*. 2007; 35:W335–8. Web Server issue. [PubMed: 17452347]
- Juan V, Crain C, Wilson C. Evidence for evolutionarily conserved secondary structure in the H19 tumor suppressor RNA. *Nucleic Acids Research*. 2000; 28(5):1221–1227. [PubMed: 10666466]
- Katayama S, et al. Antisense transcription in the mammalian transcriptome. *Science*. 2005; 309(5740):1564–1566. [PubMed: 16141073]
- Kim J, Ashworth L, Branscomb E, Stubbs L. The human homolog of a mouse-imprinted gene, Peg3, maps to a zinc finger gene-rich region of human chromosome 19q13.4. *Genome Research*. 1997; 7(5):532–540. [PubMed: 9149948]
- Kim J, Ekram MB, Kim H, Faisal M, Frey WD, Huang JM, Tran K, Kim MM, Yu S. Imprinting control region (ICR) of the Peg3 domain. *Human Molecular Genetics*. 2012; 21(12):2677–2687. [PubMed: 22394678]
- Koerner MV, Pauler FM, Huang R, Barlow DP. The function of non-coding RNAs in genomic imprinting. *Development*. 2009; 136(11):1771–1783. [PubMed: 19429783]
- Li, WH.; Graur, D. *Fundamentals of Molecular Evolution*. Vol. 1. Sinauer Associates; Sunderland, Mass: 1991. p. 73
- Meister G, Tuschl T. Mechanisms of gene silencing by double-stranded RNA. *Nature*. 2004; 431(2006):343–349. [PubMed: 15372041]
- Mueckstein U, Tafer H, Hackermueller J, Bernhart SH, Stadler PF, Hofacker IL. Thermodynamics of RNA–RNA binding. *Bioinformatics*. 2006; 22(10):1177–1182. [PubMed: 16446276]
- Osato N, Suzuki Y, Ikeo K, Gojobori T. Transcriptional interferences in cis natural antisense transcripts of humans and mice. *Genetics*. 2007; 176(2):1299–1306. [PubMed: 17409075]
- Royo H, Cavaillie J. Non-coding RNAs in imprinted gene clusters. *Biology of the Cell*. 2008; 100(3):149–166. [PubMed: 18271756]
- Washietl S, Hofacker IL, Stadler PF. Fast and reliable prediction of noncoding RNAs. *Proceedings of the National Academy of Sciences of the United States of America*. 2005; 102(7):2454–2459. [PubMed: 15665081]
- Yamasaki K, Hayashida S, Miura K, Masuzaki H, Ishimaru T, Niikawa N, Kishino T. The novel gene, gamma2-COP (COPG2), in the 7q32 imprinted domain escapes genomic imprinting. *Genomics*. 2000; 68(3):330–335. [PubMed: 10995575]
- Yamashita M, Glasgow E, Zhang BJ, Kusano K, Gainer H. Identification of cell-specific messenger ribonucleic acids in oxytocinergic and vasopressinergic magnocellular neurons in rat supraoptic

nucleus by single-cell differential hybridization. *Endocrinology*. 2002; 143(11):4464–4476.
[PubMed: 12399444]

Abbreviations

ORF	Open Reading Frame
MFE	minimal fold energy
ncRNA	non-coding RNA
TSS	transcriptional start site
Peg3	paternally expressed gene 3
APeg3	antisense paternally expressed gene 3
CMV	Cytomegalovirus
BGH	Bovine Growth Hormone
mm9	<i>Mus musculus</i> assembly 9




Fig. 1. Evolutionary conservation of *Apeg3* among placental mammals. UCSC Genome Browser *Mus musculus* assembly 9 (mm9) displays the 100-kb genomic region surrounding mouse *Apeg3*. The transcribed and exon regions of *Peg3* and *Zim1* are also presented along with the transcribed region of *Apeg3*. Evolutionary conservation of this genomic interval among placental mammals is shown with a graphical output of the PhyloP analysis. PhyloP conservation analysis of *Zim1* 3' UTR is designated in a red box. In a zoom-in view, two black boxes represent the different-length known cDNAs for *Apeg3*. *Apeg3*-Full is based on the deposited cDNA sequences from human and rat, while *Apeg3*-Short is based on the cDNA sequences from mouse. The evolutionary conservation of *Apeg3* was further analyzed using the genomic sequences obtained from 9 representative placental mammals (Supplementary Fig. 1). This analysis identified a highly conserved TATA box (TATA), transcriptional start site (TSS) and a poly-adenylation signal (pA1). This analysis also confirmed the conservation of a potential target site for miRNA (miRNA-124). Areas of high conservation within these regions are underscored in red.




Fig. 2. Predicted secondary structures of *APEG3* mRNA. The 1.5-kb nucleotide sequences from 9 representative mammals were used to predict the global secondary structures of *APEG3* using the Alifold prediction program: sense (A) and antisense (B) strands of *APEG3*. Conserved bases are color-coded based on their conservation level (legend inset). Local secondary structures were also modeled using the lowest fold energy as well as covariant analysis on small (100 bp) windows within *APEG3*. The RNAz program predicted a single conserved hairpin-loop structure within the sense (P = 0.91) and antisense (P = 0.99) *APEG3* transcripts (C, D) with the 98.93% folding accuracy. The predicted hairpin-loop structure is located in the 5' side of the *APEG3* transcript (nucleotide positions 145–185 in mouse *APEG3*).




Fig. 3.

Effects of *APeg3* overexpression on the transcription levels of *Peg3*. A series of constructs transcribing *APeg3* were transfected into Neuro2A cells. *APeg3* transcription was driven by the Cytomegalovirus (CMV) promoter, and terminated and polyadenylated by a Bovine Growth Hormone (BGH) terminator. Transfected cells were selected with hygromycin antibiotic (500 mg/mL). Potential effects of *APeg3* overexpression were measured by determining the transcription levels of *Peg3* using the total RNA isolated from the transfected cells with qRT-PCR. The transcription levels of *Peg3* were first normalized to β -*Actin*, then compared to a control (pCDNA, empty vector). The results derived from multiple transfections (n = 6) with each trial representing three individual wells were summarized and presented as graphs with standard deviations. Asterisks represent statistically significant changes in the transcription levels of *Peg3* between two constructs.




Fig. 4. Effects of *APeg3* overexpression on the protein levels of PEG3. A set of transfected Neuro2A were further analyzed using Western blotting to measure the protein levels of PEG3. The density of the detected PEG3 bands by anti-PEG3 antibodies was normalized to that of the β -Actin band, then compared with serial dilutions of pCDNA-transfected Neuro2A as standards. Numbers below the bands describe the protein levels of PEG3 and β -Actin. Numbers within the red box indicate protein levels PEG3 relative to β -Actin.

Table 1

Pairwise sequence comparison of *APeg3*.

Organism	<i>E. caballus</i>	<i>T. truncatus</i>	<i>M. lucifugus</i>	<i>L. africana</i>	<i>T. manatus</i>	<i>D. novemcinctus</i>	<i>M. musculus</i>	<i>O. cuniculus</i>	<i>H. Sapiens</i>
<i>Equus caballus</i>	100	84.73	82.41	77.93	79.82	78.56	75.5	80.18	83.2
<i>Tursiops truncatus</i>	84.73	100	79.78	75.12	76.47	75.23	71.66	77.13	80.99
<i>Myotis lucifugus</i>	82.41	79.78	100	74.03	74.92	73.96	72.64	77.18	79.22
<i>Loxodonta africana</i>	77.93	75.12	74.03	100	89.01	77.34	68.61	73.15	75.53
<i>Trichechus manatus</i>	79.82	76.47	74.92	89.01	100	79.1	70.19	74.41	77.78
<i>Dasylops novemcinctus</i>	78.56	75.23	73.96	77.34	79.1	100	69.5	74.17	76.42
<i>Mus musculus</i>	75.5	71.66	72.64	68.61	70.19	69.5	100	79.18	77.33
<i>Oryzologus cuniculus</i>	80.18	77.13	77.18	73.15	74.41	74.17	79.18	100	83.84
<i>Homo sapiens</i>	83.2	80.99	79.22	75.53	77.78	76.42	77.33	83.84	100