

1-2-2003

Microscopic description of the scissors mode in odd-mass heavy deformed nuclei

Carlos E. Vargas
GANIL Grand Accelérateur National d'Ions Lourds

Jorge G. Hirsch
Universidad Nacional Autónoma de México

Jerry P. Draayer
Louisiana State University

Follow this and additional works at: https://digitalcommons.lsu.edu/physics_astronomy_pubs

Recommended Citation

Vargas, C., Hirsch, J., & Draayer, J. (2003). Microscopic description of the scissors mode in odd-mass heavy deformed nuclei. *Physics Letters, Section B: Nuclear, Elementary Particle and High-Energy Physics*, 551 (1-2), 98-102. [https://doi.org/10.1016/S0370-2693\(02\)03009-5](https://doi.org/10.1016/S0370-2693(02)03009-5)

This Article is brought to you for free and open access by the Department of Physics & Astronomy at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.



ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Physics Letters B 551 (2003) 98–102

PHYSICS LETTERS B

www.elsevier.com/locate/npe

Microscopic description of the scissors mode in odd-mass heavy deformed nuclei

Carlos E. Vargas^a, Jorge G. Hirsch^b, Jerry P. Draayer^c

^a *Grand Accélérateur National d'Ions Lourds, BP 5027, F-14076 Caen cedex 5, France*

^b *Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Apartado Postal 70-543, México 04510 DF, Mexico*

^c *Department of Physics and Astronomy, Louisiana State University, Baton Rouge, LA 70803, USA*

Received 4 October 2002; accepted 31 October 2002

Editor: J.-P. Blaizot

Abstract

Pseudo-SU(3) shell-model results are reported for M1 excitation strengths in ^{157}Gd , ^{163}Dy and ^{169}Tm in the energy range between 2 and 4 MeV. Non-zero pseudo-spin couplings between the configurations play a very important role in determining the M1 strength distribution, especially its rapidly changing fragmentation pattern which differs significantly from what has been found in neighboring even–even systems. The results suggest one should examine contributions from intruder levels.

© 2002 Elsevier Science B.V. Open access under [CC BY license](https://creativecommons.org/licenses/by/4.0/).

PACS: 21.60.Fw; 23.20.Js; 27.70.+q

Keywords: Pseudo-SU(3) shell model; ^{157}Gd ; ^{163}Dy ; ^{169}Tm ; Scissors mode; Odd-mass nuclei

The scissors mode in nuclei refers to a pictorial image of deformed proton and neutron distributions oscillating against one another [1]. A description of this mode within the framework of the IBM [2] led to its detection in ^{156}Gd using high-resolution inelastic electron scattering [3]. Systematic studies employing nuclear resonance fluorescence scattering (NRF) measurements [4] followed. The non-observation of these low-energy M1 excitations in inelastic proton scattering (IPS) [5] confirmed its orbital character [6]. Over the past two decades an impressive wealth of informa-

tion about the scissors mode in even–even nuclei has been obtained and analyzed [7].

Low-energy M1 transitions in odd-mass nuclei were first observed in ^{163}Dy in 1993 [8]. Unambiguous spin and parity assignments of excited states in these nuclei are difficult to make due to the half-integer character of the angular momentum of the states [9]. Furthermore, the M1 strengths in odd-mass nuclei are highly fragmented. Since the intensities are far smaller than in even–even nuclei, their identification against the background [10], which is complicated by the presence of a small amount of impurities in the target [7], requires much higher experimental resolution [11].

Theoretical descriptions of scissors modes in odd-mass nuclei have been offered within the context of the IBFM [12,13], the particle–core-coupling model [14]

E-mail addresses: vargas@ganil.fr (C.E. Vargas), hirsch@nuclecu.unam.mx (J.G. Hirsch), draayer@lsu.edu (J.P. Draayer).

and the QPNM [15]. While the different models agree in relating the presence of the uncoupled nucleon with the observed fragmentation, the detailed description of this mode, with a nearly flat spectrum in some nuclei and has well-defined peaks in others is still not understood. Recently, the interplay between the spin and orbital M1 channels was examined [16] in the energy range between 4–10 MeV [17].

In the present Letter we analyze scissors-like M1 transitions in ^{157}Gd , ^{163}Dy and ^{169}Tm . These nuclei have been studied experimentally by a number of researchers [8,9,18]. A fully microscopic description of M1 transitions strengths between 2 and 4 MeV in these rare-earth nuclei was carried out using the pseudo-SU(3) shell model. Good qualitative descriptions of the fragmentation of the M1 transition strength is obtained by including, for the first time, states with pseudo-spin 1 (in addition to $\tilde{S} = 0$) and 3/2 (in addition to $\tilde{S} = 1/2$). For normal parity levels our findings suggest that while orbital couplings are important, in odd–even mass nuclei it is spin-flip type couplings that dominate M1 strengths in the low-energy domain. These spin-flip type transitions were also found to be essential for describing the rapidly changing fragmentation patterns found in neighboring odd- A nuclei. Freezing the unique parity orbitals, which is the usual assumption, prevents the theory from giving a quantitative description of the M1 strength, a result that is not surprising since intruder states have the largest l values and therefore contribute maximally to orbital-type M1 transitions.

The pseudo-SU(3) model [19,20] capitalizes on the existence of pseudo-spin symmetry, which refers to the experimental fact that single-particle orbitals with $j = l - 1/2$ and $j = (l - 2) + 1/2$ in the shell η lie very close in energy and can therefore be labeled as pseudo-spin doublets with quantum numbers $\tilde{j} = j$, $\tilde{\eta} = \eta - 1$, and $\tilde{l} = l - 1$. Its origin has been traced back to the relativistic Dirac equation [21]. In the present version of the pseudo-SU(3) model, the intruder level with opposite parity in each major shell is removed from active consideration [22] and pseudo-orbital and pseudo-spin quantum numbers are assigned to the remaining single-particle states. This assumption represents the strongest limitation of the present model.

Many-particle states of n_α active nucleons ($\alpha = p, n$) in a given (N) normal parity shell η_α^N are

classified by the following group chain [23–25]:

$$\begin{aligned} \{1^{n_\alpha}\} \quad \{\tilde{f}_\alpha\} \quad \{f_\alpha\} \gamma_\alpha \quad (\lambda_\alpha, \mu_\alpha) \quad \tilde{S}_\alpha K_\alpha \\ U(\Omega_\alpha^N) \supset U(\Omega_\alpha^N/2) \times U(2) \supset SU(3) \times SU(2) \\ \tilde{L}_\alpha \quad J_\alpha \\ \supset SO(3) \times SU(2) \supset SU_J(2), \end{aligned} \quad (1)$$

where above each group the quantum numbers that characterize its irreducible representations (irreps) are given and γ_α and K_α are multiplicity labels of the indicated reductions.

The Hamiltonian used in the calculations includes spherical Nilsson single-particle terms for the protons and neutrons ($H_{\text{sp},\pi[v]}$), the quadrupole–quadrupole ($\tilde{Q} \cdot \tilde{Q}$) and pairing ($H_{\text{pair},\pi[v]}$) interactions, as well as three ‘rotor-like’ terms that are diagonal in the SU(3) basis:

$$\begin{aligned} H = H_{\text{sp},\pi} + H_{\text{sp},\nu} - \frac{1}{2} \chi \tilde{Q} \cdot \tilde{Q} - G_\pi H_{\text{pair},\pi} \\ - G_\nu H_{\text{pair},\nu} + a K_J^2 + b J^2 + A_{\text{sym}} \tilde{C}_2. \end{aligned} \quad (2)$$

A detailed analysis of each term of this Hamiltonian and its parametrization can be found in [25]. The three free parameters a , b , A_{sym} were fixed by the best reproduction of the low-energy spectra; no additional parameters enter into the theory—the calculated M1 transitions reported below were not fit to the data.

A description of the low-energy spectra and B(E2) transition strengths in even–even nuclei [26] and odd-mass heavy deformed nuclei [25,27] have been carried out using linear combinations of SU(3) coupled proton–neutron irreps with largest C_2 values and pseudo-spin 0 and 1/2 (for even and odd number of nucleons, respectively), which are mixed by the single-particle terms in the Hamiltonian.

The large number of states that can decay through M1 transitions to the ground state in odd-mass nuclei, led us to enlarge the basis by including states with pseudo-spin 1 and 3/2. These configurations are necessary to describe excited rotational bands and to account for the strong fragmentation of the M1 strengths between 2 and 4 MeV in odd-mass nuclei.

The inclusion of configurations with pseudo-spin 1 and 3/2 in the Hilbert space allows for a description of highly excited rotational bands in odd-mass nuclei. This is illustrated in Ref. [28], where several rotational bands in ^{157}Gd , ^{163}Dy and ^{169}Tm are described,

including both excitation energies and intra- and inter-band B(E2) transition strengths, and shown to be in close agreement with the experimental data. In contrast, when the configuration space was restricted to the most spatially symmetric configurations, those with pseudo-spin 0 and 1/2, it was only possible to describe in ^{163}Dy the first three low-energy bands [27]. The pseudo-spin symmetry is still approximately preserved in the present case, with these three low-energy bands showing only a small amount of pseudo-spin 1 and 3/2 admixing into predominantly pseudo-spin 0 and 1/2 configurations, respectively.

The M1 transitions are mediated by the operator

$$T_{\mu}^1 = \sqrt{\frac{3}{4\pi}} \mu_N \{ g_{\pi}^o L_{\mu}^{\pi} + g_{\pi}^s S_{\mu}^{\pi} + g_{\nu}^o L_{\mu}^{\nu} + g_{\nu}^s S_{\mu}^{\nu} \} \quad (3)$$

with

$$L^{\pi, [\nu]} = \sum_i^{Z, [N]} l^{\pi, [\nu]}(i),$$

$$S^{\pi, [\nu]} = \sum_i^{Z, [N]} s^{\pi, [\nu]}(i). \quad (4)$$

In Eq. (3) the orbital and ‘quenched’ (by a factor of 0.7) spin g factors for protons and neutrons are used:

$$g_{\pi}^o = 1, \quad g_{\nu}^o = 0,$$

$$g_{\pi}^s = (0.7)5.5857,$$

$$g_{\nu}^s = -(0.7)3.8263. \quad (5)$$

To evaluate the M1 transition operator between eigenstates of the Hamiltonian (2), the pseudo-SU(3) tensorial expansion of the T1 operator (3) [24] was employed.

In what follows, the $B(M1; J_i^{\pi} \rightarrow J_f^{\pi})$ transitions in ^{157}Gd , ^{163}Dy and ^{169}Tm are presented. J_i^{π} refers to the ground states $3/2^{-}$, $5/2^{-}$ and $1/2^{+}$ in these nuclei. In each figure, insert (a) corresponds to the experimental results, while insert (b) represents the theoretical values obtained with the T1 operator of Eq. (3). Insert (c) shows the values with $g_{\pi, \nu}^o$ in Eq. (3) set to zero, i.e., with only the spin part of the T1 operator taken into account, and insert (d) shows the results with $g_{\pi, \nu}^s$ in Eq. (3) set to zero, i.e., including only the orbital part of T1.

The differences between the M1 transition strength distribution in ^{157}Gd , ^{163}Dy and ^{169}Tm , shown Figs. 1,




Fig. 1. Distribution of M1 transitions between 2 and 4 MeV for ^{157}Gd . Insert (a) shows the experimental values [9], insert (b) shows the theoretical values with the complete T1 operator, insert (c) shows the values with $g_{\pi, \nu}^o = 0$ (only the spin channel) and insert (d) with $g_{\pi, \nu}^s = 0$ (only the orbital channel).

2, and 3 respectively (notice the change on the scale), are both striking and well-known [10]. In ^{157}Gd there are 88 known M1 transitions between 2 and 4 MeV, all smaller than $0.05 \mu^2$ and distributed in a nearly flat spectrum. In ^{163}Dy the M1 transition strengths are distributed only among 17 peaks, clustered in three well-defined groups, and most of them have strengths between 0.1 and $0.2 \mu^2$. ^{169}Tm has an intermediate degree of fragmentation, with some clustered structures and many transitions on the order of $0.1 \mu^2$.

Using an enlarged version of pseudo-SU(3) shell-model theory described above, we obtained a microscopic description of these M1 transitions and their fragmentation in the three nuclei. The gross features of the M1 strength distributions in each of the nuclei are clearly reproduced, i.e., the different fragmentation patterns. On the other hand, for ^{157}Gd and ^{163}Dy the M1 strength distributions are displaced toward higher energies by about 0.75 MeV and the total sums are




Fig. 2. M1 transitions for ^{163}Dy . Convention is the same as in Fig. 1. Experimental values taken from Ref. [8].




Fig. 3. M1 transitions for ^{169}Tm . Convention is the same as in Fig. 1. Experimental values were taken from Ref. [18].

underestimated. This effect could be related with the absence of spin dependent terms in the Hamiltonian (2). For ^{169}Tm the distribution in energy of the M1 strengths is correct, but some transition strengths are overestimated by a factor 2 to 3.

The ground state wave functions of the two nuclei with odd number of neutrons, ^{157}Gd and ^{163}Dy , have one important difference. In ^{163}Dy the ground state has only pseudo-spin 0 and 1/2 components, while ^{157}Gd has a 13% mixing with pseudo-spin 1 and 3/2 components. In the M1 transition matrix elements the presence of these components in the later case gives rise to interference and fragmentation, while its absence in the former nuclei is associated with few large M1 transitions. The odd proton number of ^{169}Tm allows orbital proton excitations between half-integer components, building up its large M1 summed transition strength.

Having analyzed the similarities and differences between the experimental data and the theoretical predictions, we proceed to discuss the spin and orbital contributions to the M1 transitions. In insert (c) of

each figure the M1 transition strengths calculated only with the spin operators, i.e., making $g_{\pi,\nu}^o = 0$ in Eq. (3), is presented. Insert (d) shows the M1 strength when only the orbital part of the operator (3) are included ($g_{\pi,\nu}^s = 0$). In all cases the spin coupling is by far the dominant mode, but for ^{169}Tm the orbital contribution is also large.

In the case of ^{163}Dy , there is an almost null contribution from the orbital part of the transition operator ($0.103 \mu^2$), which in fact interferes destructively with the spin channel ($0.543 \mu^2$) to produce a summed M1 strength of $0.483 \mu^2$ in the scissors energy region.

The ‘angle’ between the orbital and spin channels, as defined by Fayache et al. [16] is 110° for ^{163}Dy . For ^{157}Gd , this angle has a value of 83° and for ^{169}Tm it is 96° . From Table 1 it can be seen that below 2 MeV the spin transitions are clearly dominant. Nevertheless, it should be emphasized that contributions of the intruder sector have been neglected.

The pseudo-SU(3) shell model for odd-mass nuclei has been shown to offer a qualitative microscopic description of the scissors mode and its fragmentation. In

Table 1
Summed $B(M1; \uparrow)$ strengths (in μ^2) in the different energy regions

		$E < 2$ MeV	2–4 MeV	4 MeV $< E$
^{157}Gd	Experiment [9]		1.596 ± 0.235	
	Theory	0.232	0.782	0.613
	Spin only	0.138	0.389	0.371
	Orbital only	0.084	0.308	0.385
^{163}Dy	Experiment [8]		1.641 ± 0.338	
	Theory	0.630	0.483	0.030
	Spin only	0.908	0.543	0.026
	Orbital only	0.088	0.103	0.012
^{169}Tm	Experiment [18]	1.912 ± 0.244	2.833 ± 0.812	0.515 ± 0.274
	Theory	2.460	3.769	0.435
	Spin only	2.245	2.332	0.164
	Orbital only	1.483	1.838	0.321

order to successfully reproduce the observed fragmentation of the M1 strength, it was necessary to use realistic values for the single particle energies and to enlarge the Hilbert space to include those pseudo-SU(3) irreps with the largest C_2 values and pseudo-spin 1 and $3/2$. This expansion of the model space allowed the T1 operator to connect the ground state with many excited states ($|J_f - J_i| \leq 1$) in the energy range between 2 and 4 MeV. The transitions are dominated by spin couplings, but interference with the orbital mode is very important.

A fully quantitative treatment of the problem should take into account contribution from the intruder sector. Detailed studies of M1 transitions in other odd-mass nuclei are under investigation and should offer an opportunity to further apply and test the theory.

Acknowledgements

The authors thank S. Pittel, A. Frank and P. Van Isacker for constructive comments. This work was supported in part by CONACyT (México) and the US National Science Foundation.

References

- [1] N. Lo Iudice, F. Palumbo, Phys. Rev. Lett. 41 (1978) 1532.
- [2] A.E.L. Dieperink, Prog. Part. Nucl. Phys. 9 (1983) 121; F. Iachello, Phys. Rev. Lett. 53 (1984) 1427.
- [3] D. Bohle, et al., Phys. Lett. B 137 (1984) 27.
- [4] U.E.P. Berg, et al., Phys. Lett. B 149 (1984) 59; B. Kasten, et al., Phys. Rev. Lett. 63 (1989) 609; H.H. Pitz, et al., Nucl. Phys. A 492 (1989) 411.
- [5] C. Djalali, et al., Phys. Lett. B 164 (1985) 269; D. Frekers, et al., Phys. Lett. B 218 (1989) 429.
- [6] C. Wesselborg, et al., Z. Phys. A 323 (1986) 485.
- [7] U. Kneissl, et al., Prog. Part. Nucl. Phys. 37 (1996) 349; A. Richter, Prog. Part. Nucl. Phys. 34 (1995) 261; D. Zawischa, J. Phys. G 24 (1998) 683.
- [8] I. Bauske, et al., Phys. Rev. Lett. 71 (1993) 975.
- [9] J. Margraf, et al., Phys. Rev. C 52 (1995) 2429.
- [10] J. Enders, et al., Phys. Rev. Lett. 79 (1997) 2010.
- [11] H. Friedrich, et al., Nucl. Phys. A 567 (1994) 266.
- [12] P. Van Isacker, A. Frank, Phys. Lett. B 225 (1989) 1.
- [13] A. Frank, et al., Nucl. Phys. A 531 (1991) 125.
- [14] A.A. Raduta, N. Lo Iudice, Z. Phys. A 334 (1989) 403; A.A. Raduta, D.S. Delion, Nucl. Phys. A 513 (1990) 11.
- [15] V.G. Soloviev, et al., Phys. Rev. C 53 (1996) 1022.
- [16] M.S. Fayache, et al., Phys. Rev. C 61 (2000) 024308.
- [17] N. Lo Iudice, et al., Phys. Rev. C 64 (2001) 054301.
- [18] N. Huxel, et al., Nucl. Phys. A 645 (1999) 239.
- [19] K.T. Hecht, A. Adler, Nucl. Phys. A 137 (1969) 129.
- [20] A. Arima, et al., Phys. Lett. B 30 (1969) 517.
- [21] A.L. Blokhin, et al., Phys. Rev. Lett. 74 (1995) 4149; J.N. Ginocchio, Phys. Rev. Lett. 78 (1997) 436.
- [22] C. Vargas, J.G. Hirsch, P.O. Hess, J.P. Draayer, Phys. Rev. C 58 (1998) 1488.
- [23] J.P. Draayer, et al., Nucl. Phys. A 381 (1982) 1; J.P. Draayer, K.J. Weeks, Ann. Phys. 156 (1984) 41.
- [24] O. Castaños, et al., Ann. Phys. 329 (1987) 290.
- [25] C. Vargas, J.G. Hirsch, T. Beuschel, J.P. Draayer, Phys. Rev. C 61 (2000) 031301; C.E. Vargas, J.G. Hirsch, J.P. Draayer, Nucl. Phys. A 673 (2000) 219.
- [26] G. Popa, et al., Phys. Rev. C 62 (2000) 064313.
- [27] C.E. Vargas, et al., Phys. Rev. C 64 (2001) 034306.
- [28] C.E. Vargas, J.G. Hirsch, J.P. Draayer, Phys. Rev. C 66 (2002), in press.