

10-1-2020

The case for re-examining glycosylation inhibitors, mimetics, primers and glycosylation decoys as antivirals and anti-inflammatories in covid19

Roger A. Laine
Louisiana State University

Follow this and additional works at: https://digitalcommons.lsu.edu/biosci_pubs

Recommended Citation

Laine, R. (2020). The case for re-examining glycosylation inhibitors, mimetics, primers and glycosylation decoys as antivirals and anti-inflammatories in covid19. *Glycobiology*, 30 (10), 763-767. <https://doi.org/10.1093/glycob/cwaa083>

This Letter to the Editor is brought to you for free and open access by the Department of Biological Sciences at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Letter to the Glyco-Forum

The case for re-examining glycosylation inhibitors, mimetics, primers and glycosylation decoys as antivirals and anti-inflammatories in COVID19

Roger A Laine¹

Departments of Biological Sciences and Chemistry, Louisiana State University and A&M College, Baton Rouge, LA 70803, USA

¹To whom correspondence should be addressed: Tel: +1 225 268 3052; e-mail: rlaine@lsu.edu, rogerlaine@gmail.com

Received 20 August 2020; Accepted 20 August 2020

Introduction

The SARS COV2 and other SARS viruses employ a heavily glycosylated spike S protein (Watanabe et al. 2020; Shajahan et al. 2020a) to bind to the ACE2 cellular receptor, also heavily glycosylated (Shajahan et al. 2020b), on host cells. Historically, there is abundant information of glycosylation inhibitors for antiviral activity to stimulate their investigation for COVID19 treatment. While there are certain to be side effects by interfering systematically with glycosylation, a short-term dose of primers, decoys or glycosylation inhibitors might be sufficient to stop or slow the COVID19 infection cycle with minimal or temporary side effects. For example, castanospermine derivative celgosivir (6-O-butanoylcastanospermine) (Sels et al. 1999) and Miglitol, Miglustat (N-butyl –1-deoxynojirimycin) (Campo et al. 2013; Zamoner et al. 2019) are already FDA approved for type 2 diabetes and Gaucher's disease (Rosenbloom and Weinreb 2013), passing safety and effectivity phases. This article is to stimulate repurpose testing of these already-approved compounds and others for COVID19 therapeutics.

Previously reported efficacy of antiviral activities

Copious literature suggests that glycosylation inhibitors are potent antivirals for enveloped viruses with a coat or spike protein. In the 1970s and 1980s, a number of investigators examined interference with glycan synthesis on viral coat proteins as a mode for viral inhibition. Compans showed in 1971, *D*-glucosamine and 2-deoxy-*D*-glucose inhibited glycoprotein maturation in Sindbis virus (Compans 1971). Several other studies used these two compounds for inhibition of Semliki Forest virus (Kaluza et al. 1972), enveloped RNA viruses (Kaluza et al. 1973), Influenza (Klenk et al. 1972), VSV, Newcastle disease (Scholtissek et al. 1974) and enveloped viruses in general (Scholtissek et al. 1975). In 1973, Okayama et al. showed

that β -xylosides could stimulate free GAG chains and inhibit GAG formation on proteoglycan core proteins (Okayama et al. 1973). Later, a number of researchers showed that GAGs were important as cellular receptors in HSV and HCV and that β -xylosides could inhibit viral infectivity by sidetracking GAG synthesis to soluble saccharides (see below).

Schwartz reported, in 1974, tunicamycin inhibition of glycosylation of influenza virus hemagglutinin (Schwartz et al. 1974) and in 1976 showed that tunicamycin, which inhibits the formation of N-acetylglucosamine-lipid intermediates in N-linked glycan synthesis (Lennarz 1975), suppressed glycoprotein synthesis in Semliki Forest, influenza and avian sarcoma virus (Schwarz et al. 1976). Kornfeld et al. (1977) examined the effect of tunicamycin (Leavitt et al. 1977a) on growth of Sindbis and VSV in BHK cells and found at 500 ng/ml, 99.9% inhibition of replication without detection of non-infectious particles (Leavitt et al. 1977b). Viral proteins were synthesized, but glycoproteins were under-glycosylated (Leavitt et al. 1977b) and temperature sensitive (Gibson et al. 1979). Their conclusion was that in enveloped viruses, glycosylation was essential for normal viral particle assembly. Morrison et al. (1978) found that tunicamycin had no effect on glycoprotein attachment to intracellular membranes or protein transport to the endoplasmic reticulum lumen, but instead prevented glycoprotein migration from the rough to smooth. Kornfeld et al. (1979) found that VSV non-glycosylated glycoprotein underwent intracellular aggregation upon tunicamycin treatment (Gibson et al. 1979), an indication of improper folding or exposure of hydrophobic peptide due to pauciglycosylation. Diggelman (1979) showed biosynthesis of an unglycosylated envelope glycoprotein of Rous sarcoma virus in the presence of tunicamycin. Pizer et al. (1980) showed that tunicamycin inhibited production of infectious herpes simplex virus-1 (HSV1). Blocking addition of carbohydrate to glycoprotein precursors with tunicamycin resulted in disappearance

of proteins, postulated due to degradation of unglycosylated polypeptides. They concluded that the glycans either stabilize the envelope proteins or provide proper structure for processing of the proteins necessary for infectivity. Stallcup and Fields (1981) inhibited measles virus with tunicamycin (24).

Holmes et al. (1981) showed that a corona virus, mouse hepatitis virus, MHV, had a glycoprotein E1 glycosylated normally in the presence of tunicamycin. This indicated that E1 bore O-linked instead of N-linked glycans. The “peplomeric glycoprotein E2 was not detectable upon tunicamycin treatment,” indicating its synthesis was interdicted or its degradation was facilitated by lack of N-linked glycosylation and was improperly processed (Holmes et al. 1981). E2 may be analogous to the S-spike glycoprotein. Viral particles were produced with tunicamycin treatment but were noninfective, lacking E2, which was required for attachment to viral receptors on cells, probably analogous ACE2 by today’s knowledge. At that time, no drug was known to inhibit O-glycosylation, such as α -benzyl-GalNAc, in use today. Because of these observations, either N-linked or O-linked glycosylation interference may potentially interdict SARS-Cov2 infectivity.

Tunicamycin was employed in antiviral tests and was very effective, but it was considered too toxic by most investigators to be developed as a drug at the concentrations used. However, Banerjee et al. (2011), in cancer treatment, found a safe concentration of a purified tunicamycin and dosage in mice (250 μ g/kg oral twice a week for 4 weeks) that inhibited MDA-MB0231 triple negative breast-cancer tumor formation in live mice, without overt toxicity and involved the unfolded protein response. Nanoformulations were effective (Banerjee et al. 2013). Recent articles define the detailed basis for tunicamycin’s activity (Hakulinen et al. 2017; Yoo et al. 2018). Tunicamycins should be tested now for infectivity assays with SARS COV2 alone or in combinations with other inhibitors.

Elbein et al. (1982) showed swainsonine, an α -mannosidase inhibitor, to inhibit processing of oligosaccharides on influenza viral hemagglutinin, and in 1983 showed its effect on vesicular stomatitis virus (Kang and Elbein 1983). In 1983, he showed castanospermine, an α -glucosidase inhibitor, inhibited influenza hemagglutinin expression (Pan et al. 1983). Both inhibited N-glycan processing. Tyms et al. (1987) showed castanospermine and other plant alkaloid glucosidase inhibitors to block HIV growth. Sunkara et al. (1987) tested castanospermine and 1-deoxynojirimycin (Duvoglustat) showing anti-retroviral activity. Dwek et al. (1998) later suggested α -glucosidase inhibitors as general antiviral agents (Mehta et al. 1998). Chapel et al. (2006) showed antiviral effects of α -glucosidase inhibitors on hepatitis C virus. Chang et al. (2013a) reported that iminosugar α -glucosidase inhibitors related to 1-deoxynojirimycin (that interfered with N-linked glycan maturation in the calnexin-mediated folding pathway) were also effective as antivirals for *Arenaviridae*, *Bunyaviridae*, *Filoviridae* and *Flaviviridae* including hemorrhagic fever Marburg and Ebola in mice. 1-Deoxynojirimycin (Duvoglustat) suppresses postprandial blood glucose and is used for treatment of diabetics in doses of 20-110 mg/kg (Gao et al. 2016, Tong et al. 2018). Qu et al. (2011), inhibitors of endoplasmic reticulum α -glucosidases potently suppress hepatitis C virus virion assembly and release. Howe et al. (2013) applied similar compounds to HCV and BVDV. Castanospermine and its O- and N-butyl derivatives were tested in many viral systems with positive results and FDA approval for type 2 diabetes and Gaucher’s treatments as described below. Off-label, repurposing treatments for COVID19 should be considered.

Celgosivir (6-O-butanoyl castanospermine), an approved α -glucosidase inhibitor drug for type 2 diabetes and Gaucher’s disease, inhibits all four Dengue, DENV serotypes (Sayce et al. 2010; Rathore et al. 2011). “Fluorescence microscopy showed that the antiviral mechanism of Celgosivir, is in part, due to misfolding and accumulation of DENV non-structural protein 1 (NS1) in the endoplasmic reticulum” (Lachmann 2003). Moreover, Celgosivir modulates the host’s unfolded protein response (UPR) for its antiviral action. Significantly, Celgosivir is effective in lethal challenge mouse models that recapitulate primary or secondary antibody-dependent enhanced DENV infection. Celgosivir-treated mice showed enhanced survival, reduced viremia and robust immune response, as reflected by serum cytokine analysis. Importantly, survival increased even after treatment was delayed till 2 days post-infection.” (Rathore et al. 2011). Together, this suggests that Celgosivir, which has been clinically determined to be safe in humans, may be a valuable candidate for clinical testing in COVID19 patients. (Ibid).

Safety

Celgosivir has passed safety studies in humans (Sels et al. 1999; Rathore et al. 2011). N-butyl-deoxynojirimycin (Miglustat) has been approved for clinical use since 2002 for Gaucher’s disease (Cox et al. 2000; Lachmann 2003), after tests in 2000 to show its activity was to decrease substrate biosynthesis. This safety data suggests that both should be investigated repurposed off-label for COVID19 infectivity, since safety, PK and dosage has already been established. Also, long-term, 24-month studies of Miglustat have established safety and efficacy (Pastores et al. 2005), where COVID19 treatment may require very brief application. Chang et al. (2013b) and Pérez-García et al. (2017) reviewed and recommended the use of ER glycosylation inhibitors as potential targets for viral therapeutics. Main side effects were GI, manageable by diet and anti-propulsives. Duvoglustat, (1-deoxymannojirimycin) (Fuhrmann et al. 1984) amphotycin and related compounds should also be tested. Considering the seriousness of this pandemic, researchers with access to these and other glycosylation inhibitors should collaborate with virology labs that can test the infectivity cycle of SARS COV2.

Following Okayama’s earlier work (1973), Schwartz et al. (1974) showed stimulation of synthesis of free chondroitin sulfate chains by β -D-xylosides in cultured cells. Esko and Montgomery (1995), following Okayama’s and Schwartz’s earlier work, described xyloside “primers” of glycans that could be employed in tissue culture to prevent sugars from attaching to proteins, proteoglycans and glycolipids. They found β -D-xylosides initiate glycosaminoglycan (GAG) synthesis by substituting for endogenous xylosylated core proteins. Xylosides will also prime oligosaccharides that resemble glycolipids. N-acetyl- α -D-galactosaminides initiate O-linked oligosaccharide synthesis found on mucins and other glycoproteins and can be used to disrupt O-glycosylation. Disaccharides, (e.g. peracetylated N-acetyllactosaminide), can act as primers. Competing with endogenous substrates, they interdict proteoglycan and glycoprotein glycosylation. Esko used acetylated xylose derivatives where PNP-Xyl treatment decreased heparan sulfate expression on the cell surface of Syndecan 4 cells and abrogated the HCV transmission in a concentration-dependent manner (Shieh et al. 1992; Fritz et al. 1994; Fritz and Esko 2001). Xylosides and thioxylosides should be tested in SARS COV2 infectivity studies.

E-selectin/ligand interdiction for inflammation in the ARDS symptoms in COVID19

E-selectin is a key to the first step in inflammation where E-selectin ligand Sialyl LeX binds E-selectin on capillary endothelium for neutrophil extravasation. Mulligan *et al.* (1993) early showed in a mouse lung inflammation model of ARDS that oligosaccharides containing Sialyl LeX inhibited neutrophils from causing the pneumonic symptoms. Glycomimetics, Inc., under John Magnani's scientific direction, has developed pan-selectin inhibitors and follow-on compounds that may also be effective in interdiction of the ARDS pathology (Chang *et al.* 2010). Magnani *et al.* developed a potent anti-inflammatory Sialyl LeX analog that interdicted inflammation in sickle cell anemia (Chang *et al.* 2010), and this and follow-on compounds from glycomimetics could be employed to test COVID19 lung and cardiac inflammation. Neelamegham, Matta *et al.* (Wang *et al.* 2018) reported using thioglycoside N-glycosylation decoys to hijack E-selectin ligand glycosylation, such that neutrophils treated with acetylated GlcNAc-S-NAP are 90% inhibited in rolling and tethering, and 90% inhibited in extravasation in a mouse ex-vivo model. Experiments with lung inflammation mouse models such as used by Mulligan *et al.* (1993) should be pursued toward finding drugs for the deadly pneumonic phase of COVID19 that is responsible for most fatalities.

It seems a reasonable approach that interfering with host glycosylation systems hijacked by SARS COV2, plus interfering with the ACE2 receptor glycosylation may combine to 1) interdict infectivity and 2) glycosylation interference with Sialyl LeX can inhibit E-selectin-based inflammatory responses, mitigating the Covid19 ARDS pathology. Important here is that decoys of glycosylation address the host enzyme systems; therefore, the virus cannot perform a simple mutation to overcome the interdiction, as in vaccines, like influenza. In addition, several of the compounds are FDA approved for other indications, which could be repurposed and tested off-label.

Heparinoid efficacy

Coagulopathies and heparin treatment are the subject of several articles regarding the pathology of COVID19 (Becker 2020; Porfidia and Pola 2020), and heparan sulfate derivatives and sulfated oligosaccharides appear from several laboratories to play a role in the binding of spike protein to the ACE2 receptor (Clausen *et al.* 2020; Kim *et al.* 2020; Kwon *et al.* 2020; Mycroft-West *et al.* 2020; Tandon *et al.* 2020; Zhang *et al.* 2020). Thus, heparan sulfate derived or sulfated nonanticoagulant and anticoagulant oligosaccharides and derivatives may be effective interdictors of both viral infectivity and coagulopathies, and FDA-approved heparinoid products should be considered for repurposing for COVID19.

Summary

Glycobiologists with access to any of the compounds mentioned herein or related to glycosylation interdictors are encouraged to send their samples for testing to virology labs conducting SARS COV2/ACE2-binding studies, spike glycoprotein fusion tests, or infectivity assays. Repurposing of already approved drugs, and testing others with low toxicity may uncover other avenues from a solid glycobiology antiviral background for a much-needed development of COVID19 therapeutics. There are also outstanding opportunities for selectin anti-inflammatory and heparinoid anticoagulation approaches for COVID19 ARDS.

Conflict of interest statement

R.A. Laine is a managing partner of TumorEnd, LLC, a company working on thioglycoside anti-inflammatory (ref. 56) and anti-viral compounds (www.tumorend.com).

References

- Banerjee A, Lang JY, Hung MC *et al.* 2011. Unfolded protein response is required in nu/nu mice microvasculature for treating breast tumor with tunicamycin. *J Biol Chem.* 286:29127–29138.
- Banerjee A, Johnson KT, Banerjee IA, Banerjee DK. 2013. Nanoformulation enhances anti-angiogenic efficacy of tunicamycin. *Transl Cancer Res.* 2:240–255.
- Becker RC. 2020. COVID-19 update: Covid-19-associated coagulopathy. *J Thromb Thrombolysis.* 50:54–67.
- Campo V.L., Aragão-Leoneti V., Carvalho I. 2013. Glycosidases and diabetes: Metabolic changes, mode of action and therapeutic perspectives. In: Pilar AR, Thisbe L, editors. *Carbohydrate chemistry.* Volume 9. Royal Society of Chemistry; UK: Cambridge. p. 181–203.
- Chang J, Block TM, Guo JT. 2013b. Antiviral therapies targeting host ER alpha-glucosidases: Current status and future directions. *Antiviral Res.* 99:251–260.
- Chang J, Warren TK, Zhao X, Gill T, Guo F, Wang L, Comunale MA, Du Y, Alonzi DS, Yu W *et al.* 2013a. Small molecule inhibitors of ER α -glucosidases are active against multiple hemorrhagic fever viruses. *Antiviral Res.* 98:432–440.
- Chapel C, Garcia C, Roingard P, Zitzmann N, Dubuisson J, Dwek RA, Trepo C, Zoulim F, Durantel D. 2006. Antiviral effect of alpha-glucosidase inhibitors on viral morphogenesis and binding properties of hepatitis C virus-like particles. *J Gen Virol.* 87:861–871.
- Chang J, Patton JT, Sarkar A, Ernst B, Magnani JL, Frenette PS. 2010. GMI-1070, a novel pan-selectin antagonist, reverses acute vascular occlusions in sickle cell mice. *Blood.* 116:1779–1786.
- Clausen TM, Sandoval DR, Spliid CB, Pihl J, Painter CD, Thacker BE, Glass CA, Narayanan A, Majowicz SA, Zhang Y *et al.* 2020. SARS-CoV-2 infection depends on cellular HEPARAN SULFATE and ACE2. *bioRxiv.* 2020.07.14.201616.
- Compans RW. 1971. Location of the glycoprotein in the membrane of Sindbis virus. *Nature New Biol.* 229:114–116.
- Cox T, Lachmann R, Hollak C, Aerts J, van Weely S, Hrebicek M, Platt F, Butters T, Dwek R, Moyses C *et al.* 2000. Novel oral treatment of Gaucher's disease with N-butyldeoxyjirimycin (OGT 918) to decrease substrate biosynthesis. *Lancet.* 355:1481–1485.
- Diggelman H. 1979. Biosynthesis of an unglycosylated envelope glycoprotein of Rous sarcoma virus in the presence of tunicamycin. *J Virol.* 30:799–804.
- Elbein AD, Dorling PR, Vosbeck K, Horisberger M. 1982. Swainsonine prevents the processing of the oligosaccharide chains of influenza virus hemagglutinin. *J Biol Chem.* 257:1573–1576.
- Elstein D, Hollak C, Aerts JM, van Weely S, Maas M, Cox TM, Lachmann RH, Hrebicek M, Platt FM, Butters TD *et al.* 2004. Sustained therapeutic effects of oral miglustat (Zavesca, N-butyldeoxyjirimycin, OGT 918) in type I Gaucher disease. *J Inherit Metab Dis.* 27: 757–766.
- Esko JD, Montgomery RI. 1995. Synthetic glycosides as primers of oligosaccharide biosynthesis and inhibitors of glycoprotein and proteoglycan assembly. *Curr Protoc Mol Biol.* 32: Chapter 17: Unit17.11.-17.16 PMID: 18265139.
- Fritz TA, Lugemwa FN, Sarkar AK, Esko JD. 1994. Biosynthesis of heparan sulfate on beta-D-xylosides depends on aglycone structure. *J Biol Chem.* 269:300–307.
- Fritz TA, Esko JD. 2001. Xyloside priming of glycosaminoglycan biosynthesis and inhibition of proteoglycan assembly. *Meth Mol Biol.* 171:317–323.
- Fuhrmann U, Bause E, Legler G, Ploegh H. 1984. Novel mannosidase inhibitor blocking conversion of high mannose to complex oligosaccharides. *Nature.* 307:755–758.

- Gao K, Zheng C, Wang T, Zhao H, Wang J, Wang Z, Zhai X, Jia Z, Chen J, Zhou Y *et al.* 2016. 1-Deoxynojirimycin: Occurrence, extraction, chemistry, oral pharmacokinetics, biological activities and in Silico target fishing. *Molecules*. 21:1600.
- Gibson R, Schlesinger S, Kornfeld S. 1979. The nonglycosylated glycoprotein of vesicular stomatitis virus is temperature-sensitive and undergoes intracellular aggregation at elevated temperatures. *J Biol Chem*. 254:3600–3607.
- Hakulinen JK, Hering J, Branden G, Chen H, Snijder A, Ek M *et al.* 2017. MraY-antibiotic complex reveals details of tunicamycin mode of action. *Nat Chem Biol*. 13:265–267.
- Holmes KV, Doller EW, Sturman LS. 1981. Tunicamycin resistant glycosylation of a coronavirus glycoprotein: Demonstration of a novel type of viral glycoprotein. *Virology*. 115:334–344.
- Howe JD, Smith N, Lee MJ, Ardes-Guisot N, Vauzeilles B, Desire J, Baron A, Bleriot Y, Sollogoub M, Alonzi DS *et al.* 2013. Novel imino sugar α -glucosidase inhibitors as antiviral compounds. *Bioorg Med Chem*. 21:4831–4838.
- Kaluza G, Schmidt MFG, Scholtissek C. 1973. Effect of 2-deoxy-D-glucose on the multiplication of Semliki forest virus and the reversal of the block of mannose. *Virology*. 54:179–189.
- Kaluza G, Scholtissek C, Rott R. 1972. Inhibition of the multiplication of enveloped RNA viruses by glucosamine and 2-deoxy-D-glucose. *J Gen Virol*. 14:251–259.
- Kang MS, Elbein AD. 1983. Alterations in the structure of the oligosaccharide of vesicular stomatitis virus G protein by swainsonine. *J Virol*. 46:60–69.
- Kim SY, Jin W, Sood A, Montgomery DW, Grant OC, Fuster MM, Fu L, Dordick JS, Woods RJ, Zhang F *et al.* 2020. Glycosaminoglycan binding motif at S1/S2 proteolytic cleavage site on spike glycoprotein may facilitate novel coronavirus (SARS-CoV-2) host cell entry. *bioRxiv*. 2020.04.14.041459.
- Klenk H-D, Scholtissek C, Rott R. 1972. Inhibition of glycoprotein biosynthesis of influenza virus by D-glucosamine and 2-deoxy-D-glucose. *Virology*. 49:251–259.
- Kwon PS, Oh H, Kwon S, Jin W, Zhang F, Fraser K, Hong JJ, Linhardt RJ, Dordick JS. 2020. Sulfated polysaccharides effectively inhibit SARS-CoV-2 in vitro. *Cell Discov*. 6:50.
- Lachmann RH. 2003. Miglustat: Oxford GlycoSciences/Actelion NB-DNJ has been approved for clinical use since 2002 as a treatment for Gaucher's disease – A lysosomal storage disorder. *Curr Opin Investig Drugs*. 4:472–479.
- Leavitt R, Schlesinger S, Kornfeld S. 1977a. Tunicamycin inhibits glycosylation and multiplication of Simbis and vesicular Stomatitis viruses. *J. Virol*. 21:375–385.
- Leavitt R, Schlesinger S, Kornfeld S. 1977b. Impaired intracellular migration and altered solubility of nonglycosylated glycoproteins of vesicular stomatitis virus and Sindbis virus. *J Biol Chem*. 252: 9018–9025.
- Lennarz WJ. 1975. Lipid linked sugars in glycoprotein synthesis. *Science*. 188:986–991.
- Mehta A, Zitzmann N, Rudd PM, Block TM, Dwek RA. 1998. Alpha-glucosidase inhibitors as potential broad based anti-viral agents. *FEBS Lett*. 430:17–22.
- Morrison TG, McQuain CO, Simpson D. 1978. Assembly of viral membranes: Maturation of the vesicular stomatitis virus glycoprotein in the presence of tunicamycin. *J Virol*. 28:368–374.
- Mulligan MS, Lowe JB, Larsen RD, Paulson J, Zheng Z, DeFrees S, Maemura K, Fukuda M, Ward PA. 1993. Protective effects of sialylated oligosaccharides in immune complex-induced acute lung injury. *J. Exp. Med*. 178:623–631.
- Mycroft-West CJ, Su D, Pagani I, Rudd TR, Elli S, Guimond SE, Miller G, Meneghetti MCZ, Nader HB, Li Y *et al.* 2020. Heparin inhibits cellular invasion by SARS-CoV-2: Structural dependence of the interaction of the surface protein (spike) S1 receptor binding domain with heparin. *bioRxiv*. 2020.04.28.066761.
- Okayama M, Kimata K, Suzuki K. 1973. β -xylosides stimulate free GAG chains and inhibit GAG formation on proteoglycan core proteins, *Biochem. J. (Tokyo)*. 74:1069–1073.
- Pan YT, Hori H, Saul R, Sanford BA, Molyneux RJ, Elbein AD. 1983. Castanospermine inhibits the processing of the oligosaccharide portion of the influenza viral hemagglutinin. *Biochemistry*. 22: 3975–3984.
- Pastores GM, Barnett NL, Kolodny EH. 2005. An open-label, noncomparative study of miglustat in type I Gaucher disease: Efficacy and tolerability over 24 months of treatment. *Clin Ther*. 27:1215–1227.
- Pérez-García LA, Martínez-Duncker I, Mora Montes HM. 2017. The endoplasmic reticulum alpha-Glycosidases as potential targets for virus control. *Curr Protein Pept Sci*. 18:1090–1097.
- Pizer LI, Cohen GH, Eisenberg RJ. 1980. Effect of tunicamycin on herpes simplex virus glycoproteins and infectious virus production. *J Virol*. 34:142–153.
- Porfida A, Pola R. 2020. Venous thromboembolism and heparin use in COVID-19 patients: Juggling between pragmatic choices, suggestions of medical societies and the lack of guidelines [published correction appears in]. *J Thromb Thrombolysis*. 50:68–71.
- Qu X, Pan X, Weidner J *et al.* 2011, 2011. Inhibitors of endoplasmic reticulum α -glucosidases potently suppress hepatitis C virus virion assembly and release. *Antimicrob Agents Chemother*. 55(3):1036–1044.
- Rathore AP, Paradkar PN, Watanabe S, Tan KH, Sung C, Connelly J, Low J, Ooi EE, Vasudevan SG. 2011. Celgosivir treatment misfolds dengue virus NS1 protein, induces cellular pro-survival genes and protects against lethal challenge mouse model. *Antiviral Res*. 92:453–460.
- Rosenbloom BE, Weinreb NJ. 2013. Gaucher disease: A comprehensive review. *Crit Rev Oncog*. 18:163–175.
- Sayce AC, Miller JL, Zitzmann N. 2010. Targeting a host process as an antiviral approach against dengue virus. *Trends Microbiol*. 18:323–330.
- Scholtissek C, Rott R, Hau G, Kaluza G. 1974. Inhibition of the multiplication of VSV and Newcastle disease virus by 2-deoxy-D-glucose. *J Virol*. 13:1186–1193.
- Scholtissek C, Rott R, Klenk HD. 1975. Two different mechanisms of the inhibition of the multiplication of enveloped viruses by glucosamine. *Virology*. 63:191–200.
- Schwarz RT, Klenk HD. 1974. Inhibition of glycosylation of the influenza virus hemagglutinin. *J Virol*. 14:1023–1034.
- Schwartz NB, Galligani L, Ho P-L, Dorfman A. 1974. Stimulation of synthesis of free chondroitin sulfate chains by β -xylosides in cultured cells. *Proc Natl Acad Sci USA*. 71:4047–4051.
- Schwarz RT, Rohrschneider JM, Schmidt MFG. 1976. Suppression of glycoprotein formation of Semliki Forest, influenza and avian sarcoma virus by tunicamycin. *J Virol*. 12:782–779.
- Sels JP, Huijberts MS, Wolffenbuttel BH. 1999. Miglitol, a new alpha-glucosidase inhibitor. *Expert Opin Pharmacother*. 1:149–156.
- Shajahan A, Archer-Hartmann S, Supekar NT, Gleinich AS, Heiss C, Azadi P. 2020b. Comprehensive characterization of N- and O- glycosylation of SARSCoV-2 human receptor angiotensin converting enzyme 2. *bioRxiv*. 2020.05.01.071688.
- Shajahan A, Supekar NT, Gleinich AS, Azadi P. 2020a. Deducing the N- and O-glycosylation profile of the spike protein of novel coronavirus SARS-CoV-2. *bioRxiv*. 2020.04.01.020966.
- Shieh MT, WuDunn D, Montgomery RI, Esko JD, Spear PG. 1992. Cell surface receptors for herpes simplex virus are heparan sulfate proteoglycans. *J Cell Biol*. 116:1273–1281.
- Stallcup KC, Fields BN. 1981. The replication of measles virus in the presence of tunicamycin. *Virology*. 108:391–404.
- Sunkara PS, Bowlin TL, Liu PS, Sjoerdsma A. 1987. Antiretroviral activity of castanospermine and deoxynojirimycin, specific inhibitors of glycoprotein processing. *Biochem Biophys Res Commun*. 148:206–210.
- Tandon R, Sharp JS, Zhang F, Pomin VH, Ashpole NM, Mitra D, Jin W, Liu H, Sharma P, Linhardt RJ. 2020. Effective inhibition of SARS-CoV-2 entry by heparin and enoxaparin derivatives. *bioRxiv*. 2020.06.08.140236.
- Tong TT, Zhao EH, Gao HL, Xu YH, Zhao YJ, Fu G, Cui HJ. 2018. Recent Research Advances of 1-deoxynojirimycin and its derivatives. *Zhongguo Zhong Yao Za Zhi* 43:1990–1997.
- Tyms AS, Berrie EM, Ryder TA, Nash RJ, Hegarty MP, Taylor DL, Mobberley MA, Davis JM, Bell EA, Jeffries DJ *et al.* 1987. Castanospermine and other

- plant alkaloid inhibitors of glucosidase activity block the growth of HIV. *Lancet*. 2:1025–1026.
- Wang SS, Gao X, delSolar V, Yu X, Antonopoulos A, Friedman AE, Matich EK, Atilla-Gokcumen GE, Nasirikenari M, Lau JT *et al.* 2018. Thioglycosides are efficient metabolic decoys of glycosylation that reduce Selectin dependent leukocyte adhesion. *Cell Chemical Biology*. 25:1519–1532.
- Watanabe Y, Allen JD, Wrapp D, McLellan JS, Crispin M. 2020. Site-specific analysis of the SARS-CoV-2 glycan shield. *bioRxiv*. 2020.03.26.010322.
- Yoo J, Mashalidis EH, Kuk ACY, Yamamoto K, Kaeser B, Ichikawa S *et al.* 2018. GlcNAc-1-P-transferase-tunicamycin complex structure reveals basis for inhibition of N-glycosylation. *Nat Struct Mol Biol*. 25: 217–224.
- Zamoner LOB, Aragão-Leoneti V, Carvalho I. 2019. Iminosugars: Effects of stereochemistry, ring size, and N-substituents on Glucosidase activities. *Pharmaceuticals (Basel)*. 12:108.
- Zhang Q, Chen CZ, Swaroop M, Xu M, Wang L, Lee J, Pradhan M, Shen M, Luo XY, Huang Z *et al.* 2020. Targeting heparan sulfate proteoglycan-assisted endocytosis as a COVID-19 therapeutic option. *bioRxiv*. 2020.07.14.202549.