12-14-2007

Search for exclusive $\gamma\gamma$ production in hadron-hadron collisions

T. Aaltonen
Helsingin Yliopisto

A. Abulencia
University of Illinois Urbana-Champaign

J. Adelman
The Enrico Fermi Institute

T. Affolder
University of California, Santa Barbara

T. Akimoto
University of Tsukuba

See next page for additional authors

Follow this and additional works at: https://digitalcommons.lsu.edu/physics_astronomy_pubs

Recommended Citation

Aaltonen, T., Abulencia, A., Adelman, J., Affolder, T., Akimoto, T., Albrow, M., Amerio, S., Amidei, D., Anastassov, A., Anikeev, K., Annovi, A., Antos, J., Aoki, M., Apollinari, G., Arisawa, T., Artikov, A., Ashmanskas, W., Attal, A., Aurisano, A., Azfar, F., Azzi-Bacchetta, P., Azzurri, P., Bacchetta, N., Badgett, W., Barbaro-Galtieri, A., Barnes, V., Barnett, B., & Baroiant, S. (2007). Search for exclusive $\gamma\gamma$ production in hadron-hadron collisions. *Physical Review Letters*, 99 (24) <https://doi.org/10.1103/PhysRevLett.99.242002>

This Article is brought to you for free and open access by the Department of Physics & Astronomy at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Authors

T. Aaltonen, A. Abulencia, J. Adelman, T. Affolder, T. Akimoto, M. G. Albrow, S. Amerio, D. Amidei, A. Anastassov, K. Anikeev, A. Annovi, J. Antos, M. Aoki, G. Apollinari, T. Arisawa, A. Artikov, W. Ashmanskas, A. Attal, A. Aurisano, F. Azfar, P. Azzi-Bacchetta, P. Azzurri, N. Bacchetta, W. Badgett, A. Barbaro-Galtieri, V. E. Barnes, B. A. Barnett, and S. Baroiant

Search for Exclusive $\gamma\gamma$ Production in Hadron-Hadron Collisions

T. Aaltonen,²³ A. Abulencia,²⁴ J. Adelman,¹³ T. Affolder,¹⁰ T. Akimoto,⁵⁵ M. G. Albrow,¹⁷ S. Amerio,⁴³ D. Amidei,³⁵ A. Anastassov,⁵² K. Anikeev,¹⁷ A. Annovi,¹⁹ J. Antos,¹⁴ M. Aoki,⁵⁵ G. Apollinari,¹⁷ T. Arisawa,⁵⁷ A. Artikov,¹⁵ W. Ashmanskas,¹⁷ A. Attal,³ A. Aurisano,⁵³ F. Azfar,⁴² P. Azzi-Bacchetta,⁴³ P. Azzurri,⁴⁶ N. Bacchetta,⁴³ W. Badgett,¹⁷ A. Barbaro-Galtieri,²⁹ V. E. Barnes,⁴⁸ B. A. Barnett,²⁵ S. Baroian,⁷ V. Bartsch,³¹ G. Bauer,³³ P.-H. Beauchemin,⁶¹ F. Bedeschi,⁴⁶ S. Behari,²⁵ G. Bellettini,⁴⁶ J. Bellinger,⁵⁹ A. Belloni,³³ D. Benjamin,¹⁶ A. Beretvas,¹⁷ J. Beringer,²⁹ T. Berry,³⁰ A. Bhatti,⁵⁰ M. Binkley,¹⁷ D. Bisello,⁴³ I. Bizjak,³¹ R. E. Blair,² C. Blocker,⁶ B. Blumenfeld,²⁵ A. Bocci,¹⁶ A. Bodek,⁴⁹ V. Boisvert,⁴⁹ G. Bolla,⁴⁸ A. Bolshov,³³ D. Bortoletto,⁴⁸ J. Boudreau,⁴⁷ A. Boveia,¹⁰ B. Brau,¹⁰ L. Brigliadori,⁵ C. Bromberg,³⁶ E. Brubaker,¹³ J. Budagov,¹⁵ H. S. Budd,⁴⁹ S. Budd,²⁴ K. Burkett,¹⁷ G. Busetto,⁴³ P. Bussey,²¹ A. Buzatu,⁶¹ K. L. Byrum,² S. Cabrera,¹⁶ M. Campanelli,²⁰ M. Campbell,³⁵ F. Canelli,¹⁷ A. Canepa,⁴⁵ S. Carrillo,¹⁸ D. Carlsmith,⁵⁹ B. Caron,³⁴ R. Carosi,⁴⁶ S. Carron,⁶¹ B. Casal,¹¹ M. Casarsa,⁵⁴ A. Castro,⁵ P. Catastini,⁴⁶ D. Cauz,⁵⁴ M. Cavalli-Sforza,³ A. Cerri,²⁹ L. Cerrito,³¹ S. H. Chang,²⁸ Y. C. Chen,¹ M. Chertok,⁷ G. Chiarelli,⁴⁶ G. Chlachidze,¹⁷ F. Chlebana,¹⁷ I. Cho,²⁸ K. Cho,²⁸ D. Chokheli,¹⁵ J. P. Chou,²² G. Choudalakis,³³ S. H. Chuang,⁵² K. Chung,¹² W. H. Chung,⁵⁹ Y. S. Chung,⁴⁹ M. Ciljak,⁴⁶ C. I. Ciobanu,²⁴ M. A. Ciocci,⁴⁶ A. Clark,²⁰ D. Clark,⁶ M. Coca,¹⁶ G. Compostella,⁴³ M. E. Convery,⁵⁰ J. Conway,⁷ B. Cooper,³¹ K. Copic,³⁵ M. Cordelli,¹⁹ G. Cortiana,⁴³ F. Crescioli,⁴⁶ C. Cuenca Almenar,⁷ J. Cuevas,¹¹ R. Culbertson,¹⁷ J. C. Cully,³⁵ S. DaRonco,⁴³ M. Datta,¹⁷ S. D'Auria,²¹ T. Davies,²¹ D. Dagenhart,¹⁷ P. de Barbaro,⁴⁹ S. De Cecco,⁵¹ A. Deisher,²⁹ G. De Lentdecker,⁴⁹ G. De Lorenzo,³ M. Dell'Orso,⁴⁶ F. Delli Paoli,⁴³ L. Demortier,⁵⁰ J. Deng,¹⁶ M. Deninno,⁵ D. De Pedis,⁵¹ P. F. Derwent,¹⁷ G. P. Di Giovanni,⁴⁴ C. Dionisi,⁵¹ B. Di Ruzza,⁵⁴ J. R. Dittmann,⁴ M. D'Onofrio,³ C. Dörr,²⁶ S. Donati,⁴⁶ P. Dong,⁸ J. Donini,⁴³ T. Dorigo,⁴³ S. Dube,⁵² J. Efron,³⁹ R. Erbacher,⁷ D. Errede,²⁴ S. Errede,²⁴ R. Eusebi,¹⁷ H. C. Fang,²⁹ S. Farrington,³⁰ I. Fedorko,⁴⁶ W. T. Fedorko,¹³ R. G. Feild,⁶⁰ M. Feindt,²⁶ J. P. Fernandez,³² R. Field,¹⁸ G. Flanagan,⁴⁸ R. Forrest,⁷ S. Forrester,⁷ M. Franklin,²² J. C. Freeman,²⁹ I. Furic,¹³ M. Gallinaro,⁵⁰ J. Galyardt,¹² J. E. Garcia,⁴⁶ F. Garbersson,¹⁰ A. F. Garfinkel,⁴⁸ C. Gay,⁶⁰ H. Gerberich,²⁴ D. Gerdes,³⁵ S. Giagu,⁵¹ P. Giannetti,⁴⁶ K. Gibson,⁴⁷ J. L. Gimmell,⁴⁹ C. Ginsburg,¹⁷ N. Giokaris,¹⁵ M. Giordani,⁵⁴ P. Giromini,¹⁹ M. Giunta,⁴⁶ G. Giurgiu,²⁵ V. Glagolev,¹⁵ D. Glenzinski,¹⁷ M. Gold,³⁷ N. Goldschmidt,¹⁸ J. Goldstein,⁴² A. Golossanov,¹⁷ G. Gomez,¹¹ G. Gomez-Ceballos,³³ M. Goncharov,⁵³ O. González,³² I. Gorelov,³⁷ A. T. Goshaw,¹⁶ K. Goulianos,⁵⁰ A. Gresele,⁴³ S. Grinstein,²² C. Grosso-Pilcher,¹³ R. C. Group,¹⁷ U. Grundler,²⁴ J. Guimaraes da Costa,²² Z. Gunay-Unalan,³⁶ C. Haber,²⁹ K. Hahn,³³ S. R. Hahn,¹⁷ E. Halkiadakis,⁵² A. Hamilton,²⁰ B.-Y. Han,⁴⁹ J. Y. Han,⁴⁹ R. Handler,⁵⁹ F. Happacher,¹⁹ K. Hara,⁵⁵ D. Hare,⁵² M. Hare,⁵⁶ S. Harper,⁴² R. F. Harr,⁵⁸ R. M. Harris,¹⁷ M. Hartz,⁴⁷ K. Hatakeyama,⁵⁰ J. Hauser,⁸ C. Hays,⁴² M. Heck,²⁶ A. Heijboer,⁴⁵ B. Heinemann,²⁹ J. Heinrich,⁴⁵ C. Henderson,³³ M. Herndon,⁵⁹ J. Heuser,²⁶ D. Hidas,¹⁶ C. S. Hill,¹⁰ D. Hirschbuehl,²⁶ A. Hocker,¹⁷ A. Holloway,²² S. Hou,¹ M. Houlden,³⁰ S.-C. Hsu,⁹ B. T. Huffman,⁴² R. E. Hughes,³⁹ U. Husemann,⁶⁰ J. Huston,³⁶ J. Incandela,¹⁰ G. Introzzi,⁴⁶ M. Iori,⁵¹ A. Ivanov,⁷ B. Iyutin,³³ E. James,¹⁷ D. Jang,⁵² B. Jayatilaka,¹⁶ D. Jeans,⁵¹ E. J. Jeon,²⁸ S. Jindariani,¹⁸ W. Johnson,⁷ M. Jones,⁴⁸ K. K. Joo,²⁸ S. Y. Jun,¹² J. E. Jung,²⁸ T. R. Junk,²⁴ T. Kamon,⁵³ P. E. Karchin,⁵⁸ Y. Kato,⁴¹ Y. Kemp,²⁶ R. Kephart,¹⁷ U. Kerzel,²⁶ V. Khotilovich,⁵³ B. Kilminster,³⁹ D. H. Kim,²⁸ H. S. Kim,²⁸ J. E. Kim,²⁸ M. J. Kim,¹⁷ S. B. Kim,²⁸ S. H. Kim,⁵⁵ Y. K. Kim,¹³ N. Kimura,⁵⁵ L. Kirsch,⁶ S. Klimenko,¹⁸ M. Klute,³³ B. Knuteson,³³ B. R. Ko,¹⁶ K. Kondo,⁵⁷ D. J. Kong,²⁸ J. Konigsberg,¹⁸ A. Korytov,¹⁸ A. V. Kotwal,¹⁶ A. C. Kraan,⁴⁵ J. Kraus,²⁴ M. Kreps,²⁶ J. Kroll,⁴⁵ N. Krumnack,⁴ M. Kruse,¹⁶ V. Krutelyov,¹⁰ T. Kubo,⁵⁵ S. E. Kuhlmann,² T. Kuhr,²⁶ N. P. Kulkarni,⁵⁸ Y. Kusakabe,⁵⁷ S. Kwang,¹³ A. T. Laasanen,⁴⁸ S. Lai,⁶¹ S. Lami,⁴⁶ S. Lammel,¹⁷ M. Lancaster,³¹ R. L. Lander,⁷ K. Lannon,³⁹ A. Lath,⁵² G. Latino,⁴⁶ I. Lazzizzera,⁴³ T. LeCompte,² J. Lee,⁴⁹ J. Lee,²⁸ Y. J. Lee,²⁸ S. W. Lee,⁵³ R. Lefèvre,²⁰ N. Leonardo,³³ S. Leone,⁴⁶ S. Levy,¹³ J. D. Lewis,¹⁷ C. Lin,⁶⁰ C. S. Lin,¹⁷ M. Lindgren,¹⁷ E. Lipeles,⁹ A. Lister,⁷ D. O. Litvintsev,¹⁷ T. Liu,¹⁷ N. S. Lockyer,⁴⁵ A. Loginov,⁶⁰ M. Loreti,⁴³ R.-S. Lu,¹ D. Lucchesi,⁴³ P. Lujan,²⁹ P. Lukens,¹⁷ G. Lungu,¹⁸ L. Lyons,⁴² J. Lys,²⁹ R. Lysak,¹⁴ E. Lytken,⁴⁸ P. Mack,²⁶ D. MacQueen,⁶¹ R. Madrak,¹⁷ K. Maeshima,¹⁷ K. Makhoul,³³ T. Maki,²³ P. Maksimovic,²⁵ S. Malde,⁴² S. Malik,³¹ G. Manca,³⁰ A. Manousakis,¹⁵ F. Margaroli,⁵ R. Marginean,¹⁷ C. Marino,²⁶ C. P. Marino,²⁴ A. Martin,⁶⁰ M. Martin,²⁵ V. Martin,²¹ M. Martínez,³ R. Martínez-Ballarín,³² T. Maruyama,⁵⁵ P. Mastrandrea,⁵¹ T. Masubuchi,⁵⁵ H. Matsunaga,⁵⁵ M. E. Mattson,⁵⁸ R. Mazini,⁶¹ P. Mazzanti,⁵ K. S. McFarland,⁴⁹ P. McIntyre,⁵³ R. McNulty,³⁰ A. Mehta,³⁰ P. Mehtala,²³ S. Menzemer,¹¹ A. Menzione,⁴⁶ P. Merkel,⁴⁸ C. Mesropian,⁵⁰ A. Messina,³⁶ T. Miao,¹⁷ N. Miladinovic,⁶ J. Miles,³³ R. Miller,³⁶ C. Mills,¹⁰ M. Milnik,²⁶ A. Mitra,¹ G. Mitselmakher,¹⁸ A. Miyamoto,²⁷ S. Moed,²⁰ N. Moggi,⁵ B. Mohr,⁸ C. S. Moon,²⁸ R. Moore,¹⁷ M. Morello,⁴⁶ P. Movilla Fernandez,²⁹ J. Mülmenstädt,²⁹ A. Mukherjee,¹⁷ Th. Muller,²⁶

R. Mumford,²⁵ P. Murat,¹⁷ M. Mussini,⁵ J. Nachtman,¹⁷ A. Nagano,⁵⁵ J. Naganoma,⁵⁷ K. Nakamura,⁵⁵ I. Nakano,⁴⁰ A. Napier,⁵⁶ V. Necula,¹⁶ C. Neu,⁴⁵ M. S. Neubauer,⁹ J. Nielsen,²⁹ L. Nodulman,² O. Norniella,³ E. Nurse,³¹ S. H. Oh,¹⁶ Y. D. Oh,²⁸ I. Oksuzian,¹⁸ T. Okusawa,⁴¹ R. Oldeman,³⁰ R. Orava,²³ K. Osterberg,²³ C. Pagliarone,⁴⁶ E. Palencia,¹¹ V. Papadimitriou,¹⁷ A. Papaikonomou,²⁶ A. A. Paramonov,¹³ B. Parks,³⁹ S. Pashapour,⁶¹ J. Patrick,¹⁷ G. Pauletta,⁵⁴ M. Paulini,¹² C. Paus,³³ D. E. Pellett,⁷ A. Penzo,⁵⁴ T. J. Phillips,¹⁶ G. Piacentino,⁴⁶ J. Piedra,⁴⁴ L. Pinera,¹⁸ J. Pinfold,³⁴ K. Pitts,²⁴ C. Plager,⁸ L. Pondrom,⁵⁹ X. Portell,³ O. Poukhov,¹⁵ N. Pounder,⁴² F. Prakoshyn,¹⁵ A. Pronko,¹⁷ J. Proudfoot,² F. Ptohos,¹⁹ G. Punzi,⁴⁶ J. Pursley,²⁵ J. Rademacker,⁴² A. Rahaman,⁴⁷ V. Ramakrishnan,⁵⁹ N. Ranjan,⁴⁸ I. Redondo,³² B. Reisert,¹⁷ V. Rekovic,³⁷ P. Renton,⁴² M. Rescigno,⁵¹ S. Richter,²⁶ F. Rimondi,⁵ L. Ristori,⁴⁶ A. Robson,²¹ T. Rodrigo,¹¹ E. Rogers,²⁴ S. Rolli,⁵⁶ R. Roser,¹⁷ M. Rossi,⁵⁴ R. Rossin,¹⁰ P. Roy,⁶¹ A. Ruiz,¹¹ J. Russ,¹² V. Rusu,¹³ H. Saarikko,²³ A. Safonov,⁵³ W. K. Sakumoto,⁴⁹ G. Salamanna,⁵¹ O. Saltó,³ L. Santi,⁵⁴ S. Sarkar,⁵¹ L. Sartori,⁴⁶ K. Sato,¹⁷ P. Savard,⁶¹ A. Savoy-Navarro,⁴⁴ T. Scheidle,²⁶ P. Schlabach,¹⁷ E. E. Schmidt,¹⁷ M. P. Schmidt,⁶⁰ M. Schmitt,³⁸ T. Schwarz,⁷ L. Scodellaro,¹¹ A. L. Scott,¹⁰ A. Scribano,⁴⁶ F. Scuri,⁴⁶ A. Sedov,⁴⁸ S. Seidel,³⁷ Y. Seiya,⁴¹ A. Semenov,¹⁵ L. Sexton-Kennedy,¹⁷ A. Sfyrla,²⁰ S. Z. Shalhout,⁵⁸ M. D. Shapiro,²⁹ T. Shears,³⁰ P. F. Shepard,⁴⁷ D. Sherman,²² M. Shimojima,⁵⁵ M. Shochet,¹³ Y. Shon,⁵⁹ I. Shreyber,²⁰ A. Sidoti,⁴⁶ P. Sinervo,⁶¹ A. Sisakyan,¹⁵ A. J. Slaughter,¹⁷ J. Slaunwhite,³⁹ K. Sliwa,⁵⁶ J. R. Smith,⁷ F. D. Snider,¹⁷ R. Snihur,⁶¹ M. Soderberg,³⁵ A. Soha,⁷ S. Somalwar,⁵² V. Sorin,³⁶ J. Spalding,¹⁷ F. Spinella,⁴⁶ T. Spreitzer,⁶¹ P. Squillacioti,⁴⁶ M. Stanitzki,⁶⁰ A. Staveris-Polykalas,⁴⁶ R. St. Denis,²¹ B. Stelzer,⁸ O. Stelzer-Chilton,⁴² D. Stentz,³⁸ J. Strologas,³⁷ D. Stuart,¹⁰ J. S. Suh,²⁸ A. Sukhanov,¹⁸ H. Sun,⁵⁶ I. Suslov,¹⁵ T. Suzuki,⁵⁵ A. Taffard,²⁴ R. Takashima,⁴⁰ Y. Takeuchi,⁵⁵ R. Tanaka,⁴⁰ M. Tecchio,³⁵ P. K. Teng,¹ K. Terashi,⁵⁰ J. Thom,¹⁷ A. S. Thompson,²¹ E. Thomson,⁴⁵ P. Tipton,⁶⁰ V. Tiwari,¹² S. Tkaczyk,¹⁷ D. Toback,⁵³ S. Tokar,¹⁴ K. Tollefson,³⁶ T. Tomura,⁵⁵ D. Tonelli,⁴⁶ S. Torre,¹⁹ D. Torretta,¹⁷ S. Tourneur,⁴⁴ W. Trischuk,⁶¹ S. Tsuno,⁴⁰ Y. Tu,⁴⁵ N. Turini,⁴⁶ F. Ukegawa,⁵⁵ S. Uozumi,⁵⁵ S. Vallecorsa,²⁰ N. van Remortel,²³ A. Varganov,³⁵ E. Vataha,³⁷ F. Vazquez,¹⁸ G. Velev,¹⁷ C. Vellidis,⁴⁶ G. Veramendi,²⁴ V. Veszpremi,⁴⁸ M. Vidal,³² R. Vidal,¹⁷ I. Vila,¹¹ R. Vilar,¹¹ T. Vine,³¹ M. Vogel,³⁷ I. Vollrath,⁶¹ I. Volobouev,²⁹ G. Volpi,⁴⁶ F. Würthwein,⁹ P. Wagner,⁵³ R. G. Wagner,² R. L. Wagner,¹⁷ J. Wagner,²⁶ W. Wagner,²⁶ R. Wallny,⁸ S. M. Wang,¹ A. Warburton,⁶¹ D. Waters,³¹ M. Weinberger,⁵³ W. C. Wester III,¹⁷ B. Whitehouse,⁵⁶ D. Whiteson,⁴⁵ A. B. Wicklund,² E. Wicklund,¹⁷ G. Williams,⁶¹ H. H. Williams,⁴⁵ P. Wilson,¹⁷ B. L. Winer,³⁹ P. Wittich,¹⁷ S. Wolbers,¹⁷ C. Wolfe,¹³ T. Wright,³⁵ X. Wu,²⁰ S. M. Wynne,³⁰ A. Yagil,⁹ K. Yamamoto,⁴¹ J. Yamaoka,⁵² T. Yamashita,⁴⁰ C. Yang,⁶⁰ U. K. Yang,¹³ Y. C. Yang,²⁸ W. M. Yao,²⁹ G. P. Yeh,¹⁷ J. Yoh,¹⁷ K. Yorita,¹³ T. Yoshida,⁴¹ G. B. Yu,⁴⁹ I. Yu,²⁸ S. S. Yu,¹⁷ J. C. Yun,¹⁷ L. Zanello,⁵¹ A. Zanetti,⁵⁴ I. Zaw,²² X. Zhang,²⁴ J. Zhou,⁵² and S. Zucchelli⁵

(CDF Collaboration)

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*²*Argonne National Laboratory, Argonne, Illinois 60439, USA*³*Institut de Física d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*⁴*Baylor University, Waco, Texas 76798, USA*⁵*Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy*⁶*Brandeis University, Waltham, Massachusetts 02254, USA*⁷*University of California, Davis, Davis, California 95616, USA*⁸*University of California, Los Angeles, Los Angeles, California 90024, USA*⁹*University of California, San Diego, La Jolla, California 92093, USA*¹⁰*University of California, Santa Barbara, Santa Barbara, California 93106, USA*¹¹*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*¹²*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*¹³*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*¹⁴*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*¹⁵*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*¹⁶*Duke University, Durham, North Carolina 27708*¹⁷*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*¹⁸*University of Florida, Gainesville, Florida 32611, USA*¹⁹*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*²⁰*University of Geneva, CH-1211 Geneva 4, Switzerland*²¹*Glasgow University, Glasgow G12 8QQ, United Kingdom*²²*Harvard University, Cambridge, Massachusetts 02138, USA*

- ²³*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*
- ²⁴*University of Illinois, Urbana, Illinois 61801, USA*
- ²⁵*The Johns Hopkins University, Baltimore, Maryland 21218, USA*
- ²⁶*Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*
- ²⁷*High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan*
- ²⁸*Center for High Energy Physics: Kyungpook National University, Taegu 702-701, Korea;
Seoul National University, Seoul 151-742, Korea;
SungKyunKwan University, Suwon 440-746, Korea*
- ²⁹*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*
- ³⁰*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ³¹*University College London, London WC1E 6BT, United Kingdom*
- ³²*Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain*
- ³³*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*
- ³⁴*Institute of Particle Physics: University of Alberta, Edmonton, Canada T6G 2G7;
McGill University, Montréal, Canada H3A 2T8;
and University of Toronto, Toronto, Canada M5S 1A7*
- ³⁵*University of Michigan, Ann Arbor, Michigan 48109, USA*
- ³⁶*Michigan State University, East Lansing, Michigan 48824, USA*
- ³⁷*University of New Mexico, Albuquerque, New Mexico 87131, USA*
- ³⁸*Northwestern University, Evanston, Illinois 60208, USA*
- ³⁹*The Ohio State University, Columbus, Ohio 43210, USA*
- ⁴⁰*Okayama University, Okayama 700-8530, Japan*
- ⁴¹*Osaka City University, Osaka 588, Japan*
- ⁴²*University of Oxford, Oxford OX1 3RH, United Kingdom*
- ⁴³*University of Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy*
- ⁴⁴*LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France*
- ⁴⁵*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- ⁴⁶*Istituto Nazionale di Fisica Nucleare Pisa, Universities of Pisa, Siena
and Scuola Normale Superiore, I-56127 Pisa, Italy*
- ⁴⁷*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
- ⁴⁸*Purdue University, West Lafayette, Indiana 47907, USA*
- ⁴⁹*University of Rochester, Rochester, New York 14627, USA*
- ⁵⁰*The Rockefeller University, New York, New York 10021, USA*
- ⁵¹*Istituto Nazionale di Fisica Nucleare, Sezione di Roma I, University of Rome “La Sapienza,” I-00185 Roma, Italy*
- ⁵²*Rutgers University, Piscataway, New Jersey 08855, USA*
- ⁵³*Texas A&M University, College Station, Texas 77843, USA*
- ⁵⁴*Istituto Nazionale di Fisica Nucleare, University of Trieste/ Udine, Italy*
- ⁵⁵*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- ⁵⁶*Tufts University, Medford, Massachusetts 02155, USA*
- ⁵⁷*Waseda University, Tokyo 169, Japan*
- ⁵⁸*Wayne State University, Detroit, Michigan 48201, USA*
- ⁵⁹*University of Wisconsin, Madison, Wisconsin 53706, USA*
- ⁶⁰*Yale University, New Haven, Connecticut 06520, USA*
- ⁶¹*Institute of Particle Physics: McGill University, Montréal, Canada H3A 2T8;
and University of Toronto, Toronto, Canada M5S 1A7*

(Received 18 July 2007; published 14 December 2007)

We have searched for exclusive $\gamma\gamma$ production in proton-antiproton collisions at $\sqrt{s} = 1.96$ TeV, using 532 pb^{-1} of integrated luminosity taken by the run II Collider Detector at Fermilab. The event signature requires two electromagnetic showers, each with transverse energy $E_T > 5$ GeV and pseudorapidity $|\eta| < 1.0$, with no other particles detected in the event. Three candidate events are observed. We discuss the consistency of the three events with $\gamma\gamma$, $\pi^0\pi^0$, or $\eta\eta$ production. The probability that other processes fluctuate to ≥ 3 events is 1.7×10^{-4} . An upper limit on the cross section of $p\bar{p} \rightarrow p + \gamma\gamma + \bar{p}$ production is set at 410 fb with 95% confidence level.

DOI: [10.1103/PhysRevLett.99.242002](https://doi.org/10.1103/PhysRevLett.99.242002)

PACS numbers: 13.85.Qk, 13.85.Rm, 14.80.Bn

We have searched for the “exclusive” process $p + \bar{p} \rightarrow p + \gamma\gamma + \bar{p}$ in the Collider Detector at Fermilab, CDF II, at $\sqrt{s} = 1.96$ TeV. *Exclusive* means that no other particles

are produced; in our study the p and $\bar{p}$ emerge intact with small transverse momenta p_T [1] and the two photons are central with pseudorapidity, $|\eta| < 1.0$. An exclusive $\gamma\gamma$

FIG. 1. The dominant diagram for central exclusive $\gamma\gamma$ production in $p\bar{p}$ collisions. The primary process is $gg \rightarrow \gamma\gamma$ through quark loops, with a screening gluon to cancel the exchanged color.

event can be produced via $gg \rightarrow \gamma\gamma$ through a quark loop, with an additional “screening” gluon exchanged to cancel the color of the interacting gluons, and so allow the leading hadrons to stay intact, as shown in Fig. 1. This process offers a novel possibility to test QCD and is closely related [2–5] to exclusive Higgs boson [6] production at the LHC $p + p \rightarrow p + H + p$, where the production mechanism of the Higgs boson is gg fusion through a top quark loop. In both cases the final state, $\gamma\gamma$ or H , is not strongly interacting, and thus the QCD calculation of both diagrams is similar. However, the calculation is difficult as the screening gluon has low Q^2 , and other nonperturbative interactions in the same $p\bar{p}$ collision could produce additional particles. Calculations for exclusive Higgs boson production have been made using a variety of models, but these predictions cover a range of over 2 orders of magnitude [4,5]. Since the QCD part of the calculation is the same for H and $\gamma\gamma$ production, and only the calculable matrix elements $gg \rightarrow \gamma\gamma$ and $gg \rightarrow H$ are different, exclusive $\gamma\gamma$ production provides an excellent test of the theoretical predictions for H production. For exclusive production of two photons, each with transverse energy [1] $E_T^\gamma > 5$ GeV and pseudorapidity $|\eta^\gamma| < 1$, the only predicted cross sections [3] are 36 fb at the Tevatron, at $\sqrt{s} = 1.96$ TeV, and 200 fb at the LHC. The same authors predict $\sigma(p + p \rightarrow p + H + p) = 3$ fb at the LHC for a standard model Higgs boson with $M_H = 120\text{--}140$ GeV/ c^2 , claiming a factor of about three uncertainty for both processes. However, a next-to-leading order calculation has not been done, and so these uncertainties are difficult to estimate.

Processes other than $gg \rightarrow \gamma\gamma$ can produce an exclusive $\gamma\gamma$ final state. Contributions from $q\bar{q} \rightarrow \gamma\gamma$ and $\gamma\gamma \rightarrow \gamma\gamma$ are, respectively, $<5\%$ and $<1\%$ of $gg \rightarrow \gamma\gamma$ [3]. The dominant backgrounds to the observation of exclusive $\gamma\gamma$ events are the production of $\pi^0\pi^0$ or $\eta\eta$, with each meson decaying to two photons. No theoretical calculation of exclusive $\pi^0\pi^0$ or $\eta\eta$ production has been published; however, both cross sections are estimated [7] to be about

25% of the diphoton process, in the kinematic range of this study.

This Letter presents the first search for exclusive $\gamma\gamma$ production in hadronic interactions. We use 532 pb^{-1} integrated luminosity of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV delivered to the CDF II detector at the Tevatron. The CDF II detector is a general purpose detector described elsewhere [8]; here we give a brief summary of the detector components used in this analysis. Surrounding the beam pipe is a tracking system consisting of a silicon microstrip detector, a cylindrical drift chamber (COT), and a solenoid providing a 1.4 Tesla magnetic field. The tracking system has nearly 100% efficiency for reconstructing isolated tracks with $p_T \geq 1$ GeV/ c and $|\eta| < 1$. It is surrounded by the central and end-plug calorimeters covering the range $|\eta| < 3.6$. Both calorimeters have separate electromagnetic and hadronic compartments. A proportional wire chamber (CES) [9] is embedded in the central electromagnetic calorimeter, $|\eta| < 1.1$, at a depth of six radiation lengths. It allows a measurement of the number and shape, in both transverse directions, of electromagnetic showers. The anode wire pitch (in ϕ) is 1.5 cm and the cathode strip pitch varies with η from 1.7 cm to 2.0 cm. The CES provides a means of distinguishing single photon showers from $\pi^0 \rightarrow \gamma\gamma$ and $\eta \rightarrow \gamma\gamma$. The region $3.6 < |\eta| < 5.2$ is covered by a lead-liquid scintillator calorimeter called the miniplug [10]. At higher pseudorapidities, $5.4 < |\eta| < 7.4$, scintillation counters, called beam shower counters (BSC), are located on each side of the CDF detector. Gas Cerenkov detectors covering $3.7 < |\eta| < 4.7$ determine the luminosity with a 6% uncertainty [11].

Exclusive $\gamma\gamma$ production is modeled with the EXHUME Monte Carlo generator [12], based on theoretical calculations [3,13]. Simulated single photons, and photons from π^0 and η decay, are passed through the GEANT [14] based detector simulation [15] to determine their detection efficiencies.

The event signature requires two electromagnetic showers each with transverse energy $E_T > 5$ GeV, with no other particles detected in the full CDF detector, which covers $-7.4 < \eta < +7.4$. The outgoing proton and antiproton are not detected. The event selection here follows closely that described in Ref. [16] where, using the same trigger and a similar analysis, we observed exclusive e^+e^- production. The only differences are the tracking requirements, and we restrict the $|\eta|$ coverage from ± 2.0 to ± 1.0 . The trigger requires two electromagnetic clusters and no BSC counter-activity in the region $5.4 < |\eta| < 5.9$. The measured cross section for $|\eta^e| < 2.0$ and $p_T^e \geq 5$ GeV/ c (for both e^+ and e^-) is $1.6_{-0.3}^{+0.5}(\text{stat}) \pm 0.3(\text{syst})$ pb (16 candidates with 1.9 ± 0.3 background), in agreement with the theoretical QED cross section of 1.71 ± 0.01 pb. Assuming the theoretical cross section to be correct, this agreement is evidence that the efficiency of the cuts we make to define exclusive processes is well understood.

For the diphoton analysis we select events containing two electromagnetic showers, each with $E_T > 5$ GeV and $|\eta| < 1.0$ and with a hadronic-to-electromagnetic energy ratio < 0.058 , consistent with that of a photon. We require either no tracks pointing to the showers or two adjacent tracks consistent with a photon conversion ($\gamma \rightarrow e^+e^-$). The efficiency for triggering, reconstructing, and identifying a $\gamma\gamma$ event with two photons each with $E_T^\gamma > 5$ GeV and $|\eta^\gamma| < 1$ is 0.57 ± 0.07 . Cosmic ray events are rejected by requiring that the time of each shower is consistent with photons coming from a bunch crossing. The efficiency of signal events to pass this cut is 0.93 ± 0.03 . We define “exclusivity” cuts that are designed to reject events having any additional particles in the range $|\eta| < 7.4$ that are not associated with the γ candidates; these cuts require no additional energy deposits (“particle signatures”) above noise thresholds in the calorimeters or the BSC. We do not use track or CES information in this selection. One particle can shower and cause several “signatures.” We define the exclusivity cut efficiency ε_{exc} as the probability that this exclusive requirement is not spoiled by another inelastic interaction in the same bunch crossing. It is measured, as explained in Ref. [16], as the fraction of bunch crossing triggers that pass the exclusivity cuts, which depends on the individual bunch-by-bunch luminosities. We find $\varepsilon_{\text{exc}} = 0.086 \pm 0.001$. The total efficiency is reduced by events that contain a photon conversion or electron bremsstrahlung, which fail the exclusivity requirements, estimated to be 0.87 ± 0.09 using the EXHUME simulation. The probability of the scattered $p(\bar{p})$ depositing energy in the BSC is negligible if their p_T is less than 1.2 GeV/ c .

The total efficiency for all the above event selection criteria, for the $p + \bar{p} \rightarrow p + \gamma\gamma + \bar{p}$ process, for photons each with $E_T^\gamma > 5$ GeV and $|\eta^\gamma| < 1$, is $4.0 \pm 0.7\%$. Three events pass the selection criteria; their properties are given in Table I. The E_T values of the six electromagnetic clusters are all between 5 and 7 GeV, and the azimuthal opening angle between the two photon candidates is $\Delta\phi_{\gamma\gamma} \geq 2.9$ rad, so the $\gamma\gamma$ invariant mass exceeds

TABLE I. Properties of the calorimeter showers (S) of the three candidate events: given are the E_T , the η and ϕ location, the total number of CES clusters inside the same CES chamber, N_{CES} , and the χ_{CES}^2 value (a shower shape variable, explained in the text). Also given are the probabilities that a π^0 and a photon have a χ_{CES}^2 value smaller than that observed.

Event	S	E_T (GeV)	(η, ϕ)	N_{CES}	χ_{CES}^2	$P(\pi^0)$	$P(\gamma)$
A	A1	6.8	(0.44, 6.11)	1	1.0	0.14	0.26
	A2	5.9	(0.19, 2.83)	1	1.3	0.19	0.36
B	B1	5.0	(−0.07, 4.86)	1	1.4	0.21	0.39
	B2	5.4	(0.67, 1.66)	2	...	...	...
C	C1	6.0	(−0.44, 1.66)	1	13.4	0.89	0.98
	C2	5.1	(0.22, 5.05)	2	2.2	0.33	0.57

$10 \text{ GeV}/c^2$. The difference from $\Delta\phi_{\gamma\gamma} = \pi$ may be attributed to the outgoing p and $\bar{p}$ transverse momenta.

Five background sources to exclusive $\gamma\gamma$ production are considered: cosmic rays, exclusive e^+e^- events where both electrons are misidentified as photons, nonexclusive events in which additional particles do not leave a signature in the detector, “quasiexclusive” events where one or both outgoing protons dissociate and the dissociation products are all very forward, beyond the detector coverage, and exclusive $\pi^0\pi^0$ or $\eta\eta$ production.

The cosmic ray background is determined to be negligible from the distribution of the arrival time of electromagnetic showers. Cosmic rays are also expected to give hits in the tracking detectors. However, a visual inspection of the event displays shows only random noise hits in the COT and the silicon detector for 5 of the 6 showers in Table I. In the sixth case, shower B2, an e^+e^- pair from a photon conversion is seen, with the sum of the two momenta consistent with the calorimeter shower energy.

Dielectron events [16] could be misidentified as $\gamma\gamma$ events if both electron tracks are not reconstructed or the electrons undergo energetic bremsstrahlung. This contribution is estimated from Ref. [16] to be 0.02 ± 0.02 events.

Nonexclusive events, i.e., those with central particles in addition to the two photons, may appear to be exclusive if the additional particles are not detected through inefficiency. We study this by selecting events that contain two photon candidates and no tracks in the tracking detectors (other than conversion tracks), without other requirements on the central and end-plug calorimeters. Only four events in the data sample pass these criteria: the three candidates with zero additional particle signatures, and one event with 13 signatures in the calorimeters. This background is estimated to be 0.06 ± 0.03 events by using the same shape for the distribution of additional particle signatures as in exclusive e^+e^- events [16].

The proton dissociation background is small since all the dissociation products must have $|\eta| > 7.4$ to escape detection in the BSC counters. There are also few excitation states available to the proton due to spin restrictions on the final state [13]. In Ref. [3] it was estimated that the dissociation background is not expected to exceed $\sim 0.1\%$ of the exclusive signal sample, which corresponds to ≤ 0.01 events in the three-candidate sample. We take this background to be 0.01 ± 0.01 events.

Backgrounds can arise from exclusive pair production of neutral mesons, i.e., $\pi^0\pi^0$ and $\eta\eta$. One photon from the π^0 or η decay can be undetected or, in the π^0 case, the two photon showers can merge. Exclusive $\pi^0\eta$ is suppressed by isospin conservation, and $\gamma + (\pi^0/\eta)$ is forbidden by C-parity conservation. Production of $\pi^0\pi^0$ and $\eta\eta$ cannot be unambiguously distinguished from $\gamma\gamma$ production on an event-by-event basis. Since the cross sections are not well known, these backgrounds cannot be directly calculated; we discuss them later.

We therefore observe three exclusive $p + \bar{p} \rightarrow p + (\gamma\gamma/\pi^0\pi^0/\eta\eta) + \bar{p}$ candidate events with a background of 0.09 ± 0.04 events. The probability for three or more events to be observed when 0.09 ± 0.04 (assumed to be the mean and standard deviation of a gamma distribution) are expected is 1.7×10^{-4} . We set an upper limit on the cross section for exclusive $\gamma\gamma$ production, taking into account the background and its uncertainty, the signal selection efficiency, and the integrated luminosity. A Bayesian approach is used assuming a flat prior for the cross section and a gamma distribution for the uncertainties. This gives a limit on the production cross section $\sigma(p + \bar{p} \rightarrow p + \gamma\gamma + \bar{p}) < 410$ fb (for $E_T^\gamma > 5$ GeV, $|\eta^\gamma| < 1$) at 95% confidence level.

We now discuss the three candidate events as possible $\gamma\gamma$, $\pi^0\pi^0$, or $\eta\eta$ production. The selection efficiency, including exclusivity cuts, for a photon from an isolated $\pi^0 \rightarrow \gamma\gamma$ is 13% lower than that of a direct photon, while the selection efficiency for an isolated $\eta \rightarrow \gamma\gamma$ is 35% lower. Relative to $\pi^0\pi^0$ production, $\eta\eta$ detection is further suppressed by a factor of 0.15, due to the branching fraction for $\eta \rightarrow \gamma\gamma$. We therefore treat the potential background as being predominantly $\pi^0\pi^0$.

We can only distinguish between single photons, and photons from π^0 decay, using the distribution of signals on the CES strips and wires, in the module covering $\Delta\phi = 15^\circ$ and $|\eta| < 1.1$, which contains the shower. CES clusters are formed using 11 adjacent strips or wires. We may observe two separate clusters, $N_{\text{CES}} = 2$, from π^0 decay. If we observe a single cluster, $N_{\text{CES}} = 1$, it could be from a π^0 if the two photon showers overlap or if one photon shower is not detected. The number of CES clusters in the three candidate events is shown in Table I. While only 12% of photons have a second CES cluster, 28% (46%) of the π^0 (η) do. From simulation the probability that one photon from $\pi^0/\eta \rightarrow \gamma\gamma$ is not detected in the CES, by ranging out or not interacting, is 0.125 ± 0.025 . Single clusters from photons or π^0 can be distinguished statistically using their shape. We use the distribution of pulse heights on the wires and strips to form a variable, χ_{CES}^2 , that compares the lateral shape with that for an electron shower. A simulated distribution of χ_{CES}^2 for photons and π^0 's is shown in Fig. 2; it has a longer tail for π^0 's than for photons, but it does not allow an event-by-event separation. Using the distributions in Fig. 2, the probability ($P(\gamma)$, $P(\pi^0)$) that a shower has a χ_{CES}^2 less than the observed value was calculated for the five nonconversion shower candidates. Calculated values are given in Table I.

In event A both showers are single clusters with a small χ_{CES}^2 , more consistent with originating from photons than from π^0 's. In event B shower B1 also has a very low χ_{CES}^2 , while shower B2 is a photon conversion and the χ_{CES}^2 method cannot be used. Two clusters in the CES are separated in ϕ , but not in η , as expected for a conversion. The sum of the two track momenta is 5.40 GeV/c, and the

FIG. 2 (color online). The simulated distribution for χ_{CES}^2 for prompt photons (solid histogram), and $\pi^0 \rightarrow \gamma\gamma$ decays (dashed histogram). In all cases E_T is required to be between 5 and 7 GeV.

calorimeter energy is 5.45 ± 0.35 GeV, so if there were a second photon from a π^0 or η it would have $E_\gamma < 0.55$ GeV (95% C.L.), with a probability $< 10\%$ that a π^0 or η decay would have such an energy asymmetry. Also, no additional shower is observed. Therefore events A and B clearly favor the $\gamma\gamma$ hypothesis, with three narrow single showers and one photon conversion without an accompanying shower. We cannot give an unbiased value for the $\pi^0\pi^0$ background, since this study was done *a posteriori*. In event C, one shower (C1) has a very large χ_{CES}^2 (only 2% of photon showers have a larger value), and the other shower (C2) has $N_{\text{CES}} = 2$. Both favor the hypothesis that it is a $\pi^0\pi^0$ event. A likelihood ratio calculation, using only the N_{CES} and χ_{CES}^2 distributions, favors the $\pi^0\pi^0$ hypothesis over the $\gamma\gamma$ hypothesis by a factor of 4. For one event in three candidates to be $\pi^0\pi^0$ is compatible with the theoretical estimate [7] (about 1/5).

In conclusion, we have observed three candidate events for exclusive $\gamma\gamma$, $\pi^0\pi^0$, or $\eta\eta$ production with an expected background of 0.09 ± 0.04 events. The probability to observe three or more events when 0.09 ± 0.04 are expected from other processes is 1.7×10^{-4} , corresponding to a statistical significance of 3.7σ . Though two of the candidates are most likely to arise from $\gamma\gamma$ production, the $\pi^0\pi^0/\eta\eta$ hypotheses cannot be excluded. Therefore we report a 95% C.L. upper limit on the exclusive $\gamma\gamma$ production cross section ($E_T^\gamma > 5$ GeV, $|\eta^\gamma| < 1.0$) of 410 fb, approximately a factor of 10 higher than the prediction [3]. We note that the prediction of Ref. [3] of 36^{+72}_{-24} fb would correspond to $0.8^{+1.6}_{-0.5}$ events, compatible with our observations. This result may be used to constrain calculations of exclusive Higgs boson production at the LHC; it disfavors the highest predictions.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian

Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS, the Russian Foundation for Basic Research; the Comisión Interministerial de Ciencia y Tecnología, Spain; the European Community's Human Potential Programme; the Slovak R&D Agency, and the Academy of Finland.

-
- [1] A cylindrical coordinate system is used with the z axis along the proton beam direction; θ is the polar angle and ϕ is the azimuthal angle. We define pseudorapidity as $\eta = -\ln \tan(\theta/2)$, transverse momentum as $p_T = |p| \sin\theta$, and transverse energy as $E_T = E \sin\theta$.
 - [2] M. G. Albrow *et al.*, arXiv:hep-ex/0511057.
 - [3] V. A. Khoze *et al.*, Eur. Phys. J. C **38**, 475 (2005).
 - [4] V. A. Khoze, A. D. Martin, and M. G. Ryskin, Eur. Phys. J. C **26**, 229 (2002), and references therein.

- [5] J. R. Forshaw, arXiv:hep-ph/0508274, and references therein.
- [6] P. W. Higgs, Phys. Lett. **12**, 132 (1964); Phys. Rev. **145**, 1156 (1966); F. Englert and R. Brout, Phys. Rev. Lett. **13**, 321 (1964); G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, Phys. Rev. Lett. **13**, 585 (1964).
- [7] V. A. Khoze and M. G. Ryskin (private communication).
- [8] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005), and references therein; D. Amidei *et al.* (CDF Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **350**, 73 (1994); F. Abe *et al.* (CDF Collaboration), Phys. Rev. D **50**, 2966 (1994).
- [9] L. Balka *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **267**, 272 (1988).
- [10] M. Gallinaro *et al.*, IEEE Trans. Nucl. Sci. **52**, 879 (2005).
- [11] D. Acosta *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **494**, 57 (2002).
- [12] J. Monk and A. Pilkington, Comput. Phys. Commun. **175**, 232 (2006).
- [13] V. A. Khoze *et al.*, Eur. Phys. J. C **23**, 311 (2002).
- [14] GEANT, Detector description and simulation tool, CERN Program Library Long Writeup Report No. W5013, 1993.
- [15] E. Gerchtein and M. Paulini, arXiv:physics-0306031 [Computing in High Energy and Nuclear Physics (to be published)].
- [16] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **98**, 112001 (2007).