

12-24-2009

Auger decay in krypton induced by attosecond pulse trains and twin pulses

Christian Buth
Louisiana State University

Kenneth J. Schafer
Louisiana State University

Follow this and additional works at: https://digitalcommons.lsu.edu/physics_astronomy_pubs

Recommended Citation

Buth, C., & Schafer, K. (2009). Auger decay in krypton induced by attosecond pulse trains and twin pulses. *Journal of Physics: Conference Series*, 194 (2) <https://doi.org/10.1088/1742-6596/194/2/022038>

This Conference Proceeding is brought to you for free and open access by the Department of Physics & Astronomy at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

OPEN ACCESS

Auger decay in krypton induced by attosecond pulse trains and twin pulses

To cite this article: Christian Buth and Kenneth J Schafer 2009 *J. Phys.: Conf. Ser.* **194** 022038

View the [article online](#) for updates and enhancements.

You may also like

- [Time-and-energy-resolved measurement of Auger cascades following Kr 3d excitation by attosecond pulses](#)
A J Verhoef, A V Mitrofanov, X T Nguyen et al.
- [Neon in ultrashort and intense x-rays from free electron lasers](#)
Christian Buth, Randolph Beerwerth, Razib Obaid et al.
- [Conformational and nuclear dynamics effects in molecular Auger spectra: fluorine core-hole decay in CF₄](#)
T Arion, O Takahashi, R Püttner et al.



The Electrochemical Society
Advancing solid state & electrochemical science & technology

242nd ECS Meeting

Oct 9 – 13, 2022 • Atlanta, GA, US

Abstract submission deadline: **April 8, 2022**

Connect. Engage. Champion. Empower. Accelerate.

MOVE SCIENCE FORWARD



Submit your abstract



Auger decay in krypton induced by attosecond pulse trains and twin pulses

Christian Buth¹, Kenneth J. Schafer²

Department of Physics and Astronomy, Louisiana State University, Baton Rouge, Louisiana 70803, USA

Synopsis Using attoscience, we study the electron correlations responsible for Auger decay in krypton atoms. The Auger decay is induced by a pulse train or a twin pulse composed of subpulses of attosecond duration. During the Auger decay an optical dressing laser may be present. Interference effects between the ejected Auger electron wave packets are predicted.

Attosecond science aims at studying and controlling the motion of electrons on their natural time scale. The investigation of electronic motion has so far been restricted mostly to one-electron processes. One of the first applications of attoscience was the measurement of the Auger decay time of $3d$ vacancies in krypton atoms—a well-known datum from frequency-domain spectroscopy—with a single attosecond pulse in the presence of an optical streaking laser.

In the present work, we examine the Auger electron spectrum of krypton induced by an arbitrarily shaped XUV light pulse with and without an optical dressing laser. Specifically, we consider Auger decay after

1. a **train of attosecond pulses** with a set duration between the pulses and
2. an **attosecond twin pulse** with varying time delay between the subpulses: an *archetypical* interference experiment [Fig. 1].

The first point refers to a proposed experiment which will repeat the measurement of the Auger decay lifetime of krypton $3d$ vacancies with pulse trains [1]. A train produces a stronger signal than a single pulse and enhances the signal to noise ratio in an experiment. Further a train of pulses leads to electron wave packet interference of the wave packets launched by different pulses. An additional optical streaking laser facilitates the control and measurement of the resulting signal.

The second point proposes an experiment with a simpler setting compared with a pulse train. Additional flexibility is added by varying the time delay between the two attosecond pulses. Predictions of the second setting are displayed in Fig. 1. The upper panel shows the attosecond twin pulse and the exponential decay of the Auger population from a rate equation model in the time domain. After a pulse, the population of inner-shell ionized atoms decays

exponentially. The lower panel shows the Auger electron spectrum from our quantum dynamical model [2,3], if the photoelectron energy is observed in coincidence with the Auger electrons with the stated energy resolution. We predict interference between the two ejected Auger electron wave packets.




Fig. 1. (Color) Upper panel: intensity of an XUV twin pulse (red) and the krypton $3d$ hole population (black). Lower panel: probability density of Auger electrons with momentum \vec{k}_A in the opposite direction of the linear XUV polarization vector for three energy resolutions of the photoelectron measurement.

In conclusion, our proposed experiments are very fundamental; they represent a crucial test for attoscience: to what degree can we measure and control an ultrafast electronic process in the time domain?

References

- [1] Mauritsson J 2009 private communication.
- [2] Buth C and Schafer K J 2009 manuscript in preparation.
- [3] Smirnova O, Yakovlev V S and Scrinzi A 2003 Phys. Rev. Lett. **91**, 253001.

¹ E-mail: christian.buth@web.de

² E-mail: schaffer@phys.lsu.edu